

References

- 1) Japan Environment Agency (1998) Strategic Programs on Environmental Endocrine Disruptors'98 - SPEED'98.
- 2) Soto, A.M., Justicia, H., Wray, J.W. and Sonnenschein, C. (1991) *p*-Nonyl-phenol: An estrogenic xenobiotic released from "modified" polystyrene. *Environmental Health Perspective*, 92, 167-173.
- 3) Sumpter, J.P. and Jobling, S. (1995) Vitellogenesis as a biomarker of estrogenic contamination in the aquatic environment. *Environmental Health Perspective*, 103, 173-178.
- 4) Japan Society on Water Environment (edit.) (2000) Nonionic surfactant and water environment: uses, measuring technology, ecological effect. Gihodo Shuppan.
- 5) Bhatt, B.D., Prasad, J.V., Kalpana, G. and Ali, S. (1992) Separation and characterization of isomers of *p*-nonylphenols by capillary GC/GC-MS/GC-FTIR techniques. *Journal of Chromatographic Science*, 30, 203-210.
- 6) Wheeler, T.F., Heim, J.R., LaTorre, M.R. and Janes, A.B. (1997) Mass spectral characterization of *p*-nonylphenol isomers using high-resolution capillary GC-MS. *Journal of Chromatographic Science*, 35, 19-30
- 7) European Union (2001) European Union Risk Assessment Report. 4-Nonylphenol (branched) and Nonylphenol (unpublished).
- 8) Talmage, S.S. (1994) Environmental and human safety of major surfactants. Alcohol ethoxylates and alkylphenol ethoxylates. The Soap and Detergent Association.
- 9) Kutzer, T. (2001) E-mail from CAS Helpdesk, help@cas.org
- 10) ChemFinder.com Search Result Page (2001) 4-nonylphenol (branched) [84852-15-3], http://chemfinder.camsoft.com/result.asp?mol_rel_id=84852-15-3
- 11) Japan Surfactant Industry Association (1999) Materials for OECD Expert Meeting on short-chained chlorinated paraffins and nonylphenol ethoxylates.
- 12) Toray Research Center, Research & Study Dept. (edit.) (1998) The present state and challenges on endocrine disrupting substances. Toray Research Center.
- 13) Japan Ministry of Economy, Trade and Industry (1998) Dynamic statistics on mining & industry in 1998.
- 14) Japan Ministry of Economy, Trade and Industry (1999) Dynamic statistics on mining & industry in 1999.
- 15) The Chemical Daily Co., Ltd. (2001) 13901 chemical products.
- 16) Japan Surfactant Industry Association (2001) Statistical yearbook on surfactants, etc. 2000.
- 17) Environment Canada Health Canada (2000) Canadian Environmental Protection Act, Priority Substances List, Assessment Report, Nonylphenol and its ethoxylates. Draft for Public Comments.

- 18) Japan Ministry of Environment (2001) Survey report on environmental load of endocrine disrupting chemical substances in FY2000.
- 19) Tsuda, T., Takino, A., Kojima, M., Harada, H., Muraki, K. and Tsuji, M. (2000) 4-nonylphenols and 4-*tert*-octylphenol in water and fish from rivers flowing into Lake Biwa. *Chemosphere*, 41, 757-762.
- 20) Tsuda, T., Takino, A., Muraki, K., Harada, H. and Kojima, M. (2001) Evaluation of 4-nonylphenols and 4-*tert*-octylphenol contamination of fish in rivers by laboratory accumulation and excretion experiments. *Wat., Res.*, 35, 7, 1786-1792.
- 21) Ahel, M., McEvoy, J. and Giger, W. (1993) Bioaccumulation of the lipophilic metabolites of nonionic surfactants in freshwater organisms. *Environmental Pollution*, 79, 243-248.
- 22) Ahel, M. (1987) Biogeochemical behaviour of alkylphenol polyethoxylates in the aquatic environment. University of Zagreb.
- 23) Sundaram, K.M.S. and Szeto, S. (1981) The dissipation of nonylphenol in stream and pond water under simulated field conditions. *J. Environ. Sci. Health*, B16, 6, 767-776
- 24) Lewis, S.K. and Lech, J.J. (1996) Uptake, disposition, and persistence of nonylphenol from water in rainbow trout (*Oncorhynchus mykiss*). *Xenobiotica*, 26, 8, 813-819.
- 25) Blackburn, M.A., Kirby, S.J. and Waldo, M.J. (1999) Concentrations of alkylphenol polyethoxylates entering UK estuaries. *Marine Pollution Bulletin*, 38, 2, 109-118.
- 26) Chemicals Evaluation and Research Institute, Japan (1992) Collection of safety inspection data of existing chemical substances under the law concerning examination and regulation of manufacture, etc. of chemical substances class I.
- 27) Brooke, L.T. (1993) Accumulation and lethality for two freshwater fishes (fathead minnow and bluegill) to nonylphenol. USEPA Draft Report, EPA Contact No. 68-CI-0034
- 28) Ward, T.J. and Boeri, R.L. (1991) Bioconcentration test with nonylphenol and the fathead minnow (*Pimephales promelas*). Report prepared for Chemical Manufacturers Association by Resource Analysts. Study No. 8975-CMA.
- 29) Chemicals Evaluation and Research Institute (CERI), Japan (1997) OECD chemicals test guideline, a guideline on data interpretation.
- 30) Brooke, L.T. (1993) Acute and chronic toxicity of nonylphenol to ten species of aquatic organisms. USEPA Draft Report, EPA Contact No. 68-CI-0034
- 31) Holcombe, G.W., Phipps, G.L., Knuth, M.L. and Felhaber, T. (1984) The acute toxicity of selected substituted phenols, benzenes and benzoic acid esters to fathead minnows *Pimephales promelas*. *Toxicol. Chem.*, 18, 3, 363-381
- 32) Ward, T.J. and Boeri, R.L. (1990) Acute flow toxicity of nonylphenol to the sheepshead minnow (*Cyprinodon variegatus*). Report prepared for Chemical Manufacturers Association by Resource Analysts. Study No. 8972-CMA.
- 33) Ward, T.J. and Boeri, R.L. (1990) Acute flow through toxicity of nonylphenol to the mysid (*Mysidopsis capricornutum*). Report prepared for Chemical Manufacturers Association by Resource Analysts. Study No. 8974-CMA.
- 34) England, D.E. (1995) Chronic toxicity of nonylphenol to *Ceriodaphnia dubia*. Report prepared for Chemical Manufacturers Association by ABC Laboratories Inc. Report #41756.
- 35) England, D.E. and Bussard, J.B. (1994) Toxicity of nonylphenol to the amphipod *Hyalella azteca* (Sausure). Report prepared for Chemical Manufacturers Association by ABC Laboratories Inc. Report #41569.
- 36) Huls (1992) Determination of the effects of nonylphenol on the swimming behaviour of *Daphnia magna* (in accordance with EC 84/449) Final report DK-522.
- 37) Comber, M.H.I., Williams, T.D. and Stewart, K.M. (1993) The effects of nonylphenol on *Daphnia magna*. *Water Research*, 27, 2, 273-276.
- 38) Kopf, W. (1997) Wirkung endokriner Stoffe in Biotests mit Wasserorganismen. In Stoffe mit endokriner Wirkung in Wasser. Bayerisches Landesamt für Wasserwirtschaft, Institut für Wasserforschung München (ed) Oldenbourg.
- 39) Ward, T.J. and Boeri, R.L. (1990) Acute static toxicity of nonylphenol to the marine alga (*Skeletonema costatum*). Report prepared for Chemical Manufacturers Association by Resource Analysts. Study No. 8970-CMA.
- 40) Ward, T.J. and Boeri, R.L. (1990) Acute static toxicity of nonylphenol to the freshwater alga (*Selenastrum capricornutum*). Report prepared for Chemical Manufacturers Association by Resource Analysts. Study No. 8969-CMA.

- 41) Huls(1996)Determination of the effects of nonylphenol on the growth of *Scenedesmus subspicatus* 86.81.SAG (algal growth inhibition test according to UBA Feb 1984), Report AW-185.
- 42) Ward,T.J. and Boeri,R.L.(1991)Early life stage toxicity of nonylphenol to the fathead minnow (*Pimephales promelas*). Report prepared for Chemical Manufactures Association by Resource Analysts. Study No.8979-CMA.
- 43) Ward,T.J. and Boeri,R.L.(1991)Chronic toxicity of nonylphenol to the mysid (*Mysidopsis bahia*). Report prepared for Chemical Manufactures Association by Resource Analysts. Study No.8977-CMA.
- 44) England,D.E. and Bussard,J.B.(1993)Toxicity of nonylphenol to the midge *Chironomus tentans*. Report prepared for Chemical Manufactures Association by ABC Laboratories Inc. Report #40597.
- 45) Shioda,T. and Wakabayashi,M.(2000)Evaluation of reproductivity of medaka (*Oryzias latipes*)exposed to chemicals using a 2-week reproduction test. Water Research and Technology,42,5-6,53-60.
- 46) Huls(1992)Determination of the effects of nonylphenol on the reproduction of *Daphnia magna* (in accordance with OECD Guideline 202 Part α), Final Report DL-143.
- 47) Huls(1992)Determination of the effects of nonylphenol on the reproduction of *Daphnia magna* (in accordance with OECD Guideline 202 Part α), Final Report DL-143a.
- 48) Loomis,A.K. and Thomas,P.(1999)Binding characteristics of estrogen receptor(ER)in Atlantic croaker(*Micropogonias undulatus*)testis:Different affinity for estrogens and xenobiotics from that of hepatic ER¹. Biology of Reproduction,61,51-60.
- 49) White,R.,Jobling,S.,Hoare,S.A.,Sumpter,J.P. and Parker,M.G.(1994) Environmentally persistent alkylphenolic compounds are estrogenic. Endocrinology, 135,1,175-182.
- 50) Milligan,S.R.,Khan,O. and Nash,M.(1998)Competitive binding of xenobiotic oestrogens to rat alpha-fetoprotein and to sex steroid binding proteins in human and rainbow trout(*Oncorhynchus mykiss*)plasma. Genaral and Comparative Endocrinology,112,89-95.
- 51) Flouriot,G.,Pakdel,F.,Ducouret,B. and Valotaire,Y.(1995)Influence of xenobiotics on rainbow trout liver estrogen receptor and vitellogenin gene expression. Journal of Molecular Endocrinology,15,143-151.
- 52) Islinger,M.,Pawlowski,S.,Hollert,H.,Volkl,A. and Braunbeck,T.(1999)Measurement of vitellogenin-mRNA expression in primary cultures of rainbow trout hepatocytes inAnon-radioactive dot blot/RNase protein-assay. The Science of the Total Environment,233,109-122.
- 53) Celius,T.,Haugen,T.B.,Grotmol,T. and Walther,B.T.(1999)A sensitive zonagenetic assay for rapid *in vitro* assessment for estrogenic potency of xenobiotics and mycotoxins. Environmental Health Perspective,107,1,63-68.
- 54) Jobling,S. and Sumpter,J.P.(1993)Detergent components in sewage effluent are weakly oestrogenic to fish: An *in vitro* study using rainbow trout (*Oncorhynchus mykiss*)hepatocytes. Aquatic Toxicology,27,361-372.(206)
- 55) Arukwe,A.,Forlin,L. and Goksoyr,A.(1997)Xenobiotics and steroid biotransformation enzymes in Atlantic salmon(*Salmo salar*)liver treated with an estrogenic compound, 4-nonylphenol. Environmental Toxicology and Chemistry,16,12,2576-2583.
- 56) Miles-Richardson,S.R.,Pierens,S.L.,Nichola,K.M.,Kramer,V.J.,Snyder,E.M., Snyder,S.A., Render,J.A.,Fitzgerald,S.D. and Giesy,J.P.(1999)Effects of waterborn exposure to 4-nonylphenol and nonylphenol ethoxylate on secondary sex characteristics and gonads of fathead minnow(*Pimephales promelas*). Environmental Research Section A 80,S122-S137.
- 57) Ren,L.,Lewis,S.K. and Lech,J.J.(1996)Effects of estrogen and nonylphenol on the post-transcriptional regulation of vitellogenin gene expression. Chemico-Biological Interactions,100,67-76.
- 58) Jobling,S.,Sheahan,D.,Osborne,J.A.,Matthiessen,P. and Sumpter,J.(1996)Inhibition of testicular growth in rainbow trout(*Oncorhynchus mykiss*)exposed to estrogenic alkylphenolic chemicals. Environmental Toxicology and Chemistry,15,2,194-202.

- 59) Gray, M.A. and Metcalfe, C.D. (1997) Induction of testis-ova in Japanese medaka (*Oryzias latipes*) exposed to *p*-nonylphenol. *Environmental Toxicology and Chemistry*, 16, 5, 1082-1086.
- 60) Pedersen, S.N., Christiansen, L.B., Pedersen, K.L., Korsgaard, B. and Bjerregaard, P. (1999) *In vivo* estrogenic activity of branched and linear alkylphenols in rainbow trout (*Oncorhynchus mykiss*). *The Science of the Total Environment*, 233, 89-96.
- 61) Korsgaard, B. and Pedersen, K.L. (1998) Vitellogenin in *Zoarcetes viviparus*: Purification, quantification by ELISA and induction by estradiol-17 β and 4-nonylphenol. *Comparative Biochemistry and Physiology Part C* 120, 159-166.
- 62) Fent, K., Ackermann, G.E. and Schwaiger, J. (1999) Analysis of vitellogenin mRNA by quantitative reverse transcriptional polymerase chain reaction (RT-PCR) in juvenile fish exposed for twelve months to nonylphenol. In *Proceedings of the 10th International Symposium on Pollutant Responses in marine Organisms (PRIMO 10)*, Williamsburg, Virginia.
- 63) Ashfield, L.A., Pottinger, T.G. and Sumpter, J.P. (1998) Exposure of female juvenile rainbow trout to alkylphenolic compounds results in modifications to growth and ovosomatic index. *Environmental Toxicology and Chemistry*, 17, 3, 679-686.
- 64) Christensen *et al.* (1995) As reported in "Chemicals with estrogen-like effects". *Tema Nord* 1996:580, Nordic Council of Ministers, Copenhagen.
- 65) Nimrod, A.C. and Benson, W.H. (1996) Estrogenic responses to xenobiotics in channel catfish (*Ictalurus punctatus*). *Marine Environment Research*, 42, 155-160.
- 66) Nakamura, S. and Iguchi, T. (1998) Mutation of fish observed in the Tama river, special article: the present state of endocrine disruptors. *Science*, 68, 7, 515-517.
- 67) Kahl, M.D., Makynen, E.A., Kosian, P.A. and Ankley, G.T. (1997) Toxicity of 4-nonylphenol in a life-cycle test with the midge *Chironomus tentans*. *Ecotoxicology and Environmental Safety*, 38, 155-160.
- 68) Gronen, S., Denslow, N., Manning, S., Barnes, S., Barnes, D. and Brouwer, M. (1999) Serum vitellogenin levels and reproductive impairment of male Japanese medaka (*Oryzias latipes*) exposed to 4-*tert*-octylphenol. *Environmental Health Perspective*, 107, 385-390.
- 69) Bayley, M., Nielsen, J.R. and Baatrup, E. (1999) Guppy sexual behavior as an effect biomarker of estrogen mimics. *Ecotoxicology and Environmental Safety*, 43, 68-73.
- 70) Gimeno, S., Komen, H., Jobling, S., Sumpter, J. and Browmer, T. (1998) Demasculinisation of sexually mature male common carp, *Cyprinus carpio*, exposed to 4-*tert*-pentylphenol during spermatogenesis. *Aquatic Toxicology*, 43, 93-109.
- 71) Gimeno, S., Gerritsen, A., Bowmer, T. and Komen, H. (1996) Feminization of male carp. *Nature*, 384, 221-222.
- 72) OSPAR Commission (1999) Assessment of Implementation of PARCOM Recommendation 92/8 on Nonylphenol-Ethoxylates (NPEs).
http://www.ospar.org/asp/ospar/dra_spider.asp
- 73) Japan Surfactant Industry Association (1997) List of surfactants, etc., 1998 edition.
- 74) Japan Surfactant Industry Association (2000) List of surfactants, etc., 2001 edition.
- 75) Knaak, J.B., Eldridge, J.M. and Sullivan, L.J. (1966) Excretion of certain polyethylene glycol adducts of nonylphenol by the rat. *Toxicology and Applied Pharmacology*, 9, 331-340.
- 76) Berol Kemi AB (1982) Nonylphenol acute oral toxicity in rats. Inveresk Research International project no. 230086, report no. 2379.
- 77) Huls AG (1982) Nonylphenol: An acute toxicity study (LD₅₀) in the rat. Hazleton Laboratories Deutschland project no. 222/8.
- 78) ICI Central Toxicology Laboratory (1984) Nonylphenol (ex-oil works and Rohm and Haas): Comparison of acute oral toxicities. CTL report no. CTL/L/708
- 79) Gaworski, C.L., Kimbead, E.R. and Doyle, R.L. (1979) Acute toxicity of a number of chemicals of interest to the Air Force. University of California Extension, Wright Patterson Air Force Base, report ISS AMRL-TR-79-11.
- 80) Smyth, H.F., Carpenter, C.P., Weil, C.S., Pozzani, U., Striegel, J.A. and Nycum, J.S. (1969) Range-finding toxicity data: List ϕ . *Am. Ind. Hyg. Assoc. J.*, 30, 470-476.
- 81) Huls AG (1989) Nonylphenol: 28 day oral (dietary) sub-acute toxicity study in the rat. Hazleton UK report no. 5917-671/1.
- 82) Cunny, H.C., Mayes, B.A., Rosica, K.A., Trutter, J.A. and Van Miller, J.P. (1997) Subchronic toxicity (90-day) study with *para*-nonylphenol in rats. *Regulatory Toxicology and Pharmacology*, 26, 172-178.
- 83) National Toxicology Program (1997) Final Report on the reproductive toxicity of nonylphenol (CAS #84852-15-3) administered by gavage to Sprague-Dawley rats. *R.O.W. Sciences* 8989-30.

- 84) Richards, J.F. (1989) Nonylphenol: 28 day oral (dietary) sub-acute toxicity study in the rat. Hazleton UK, England, for Huls, AG, West Germany (Report No. 5917-671/1).
- 85) Odum, J., Pyrah, I.T.G., Foster, J.R., Van Miller, J.P., Joiner, R.L. and Ashby, J. (1999) Comparative Activities of *p*-nonylphenol and diethylstilbestrol in noble rat mammary gland and uterotrophic assays. *Regulatory Toxicology and Pharmacology*, 29, 184-195.
- 86) Odum, J., Pyrah, I.T.G., Soames, A.R., Foster, J.R., Van Miller, J.P., Joiner, R.L. and Ashby, J. (1999) Effects of *p*-nonylphenol (NP) and diethylstilbestrol (DES) on the Alderley Park (Alpk) rat. Comparison of mammary gland and uterus sensitivity following oral gavage or implanted mini-pumps. *Journal of Applied Toxicology*, 19, 367-378.
- 87) de Jager, C., Bornman, M.S. and der Horst, G. (1999) 1. The effect of *p*-nonylphenol, an environmental toxicant with oestrogenic properties, on fertility potential in adult male rats. *Andrologia*, 31, 2, 99-106.
- 88) Odum, J., Lefevre, P.A., Tittenson, S., Paton, D., Routledge, E.J., Beresford, N.A., Sumpter, J.P. and Ashby, J. (1997) The rodent uterotrophic assay: Critical protocol features, study with nonylphenols, and comparison with a yeast estrogenicity assay. *Regulatory Toxicology and Pharmacology*, 25, 176-188.

- 89) Coldham, N.G., Dave, M., Sivapathasundaram, S., McDonnell, D.P., Connor, C., and Sauer, M.J. (1997) Evaluation of a recombinant yeast cell estrogen screening assay. *Environmental Health Perspective*, 105, 7, 734-742.
- 90) Shelby, M.D., Newbold, R.R., Tully, D.B., Chae, K. and Davis, V.L. (1996) Assessing environmental chemicals for estrogen. Using a combination of *in vitro* and *in vivo* assays. *Environmental Health Perspective*, 104, 12, 1296-1300.
- 91) Lee, P.C. (1998) Disruption of male reproductive tract development by administration of the xenoestrogen, nonylphenol, to male newborn rats. *Endocrine*, 9, 105-111.
- 92) Milligan, S.R., Balasubramanian, A.V. and Kalitam, K.C. (1998) Relative potency of xenobiotic estrogens in an acute *in vivo* mammalian assay. *Environmental Health Perspective*, 106, 1, 23-26.
- 93) Nagao, T., Wada, K., Marumo, H., Yoshimura, S. and Ono, H. (2001) Reproductive effects of nonylphenol in rats after gavage administration: A two-generation study. *Reproductive Toxicology*, 15
- 94) Huls AG (1984) Mutagenitätsuntersuchung von nonylphenol mit hilfe des *Salmonella typhimurium*/Mikrosomen-Mutagenitäts-Test nach Ames. Huls report no.84/19, project X 41.
- 95) Shimizu, H., Suzuki, Y., Takemura, N., Goto, S. and Matsushita, H. (1985) The results of microbial mutation test for forty-three industrial chemicals. *Jpn.J.Ind.Health*, 27, 400-419.
- 96) Nishihara, T., Nishikawa, J., Kanayama, T., Dakeyama, F., Saito, K., Imagawa, M., Takatori, S., Kitagawa, Y., Hori, S. and Utsumi, H. (2000) Estrogenic activities of 517 chemicals by yeast two-hybrid assay. *Journal of Health Science*, 46, 4, 282-298.
- 97) Blair, R.M., Fang, H., Branham, W.S., Hass, B.S., Dial, S.L., Moland, C.L., Tong, W., Shi, L., Perkins, R. and Sheehan, D.M. (2000) The estrogen receptor relative binding affinities of 188 natural and xenochemicals: Structural diversity of ligands. *Toxicological Sciences*, 54, 138-153.
- 98) Tabira, Y., Nakai, M., Asai, D., Yakabe, Y., Tahara, Y., Shinmyozu, T., Noguchi, M., Takatsuki, M. and Shimohigashi, Y. (1999) Structural requirements of *para*-alkylphenols to bind to estrogen receptor. *Eur.J.Biochem.*, 262, 240-245.
- 99) Legler, J., van der Brink, C.E., Brouwer, A., Murk, A.J., van der Saag, P.T., Vethaak, A.D. and van der Burg, B. (1999) Development of a stably transfected estrogen receptor-mediated luciferase reporter gene assay in the human T47D breast cancer cell line. *Toxicological Sciences*, 48, 55-66.
- 100) Balaguer, P., Francois, F., Comunale, F., Fenet, H., Boussioux, A.-M., Pons, M., Nicolas, J.-C. and Casellas, C. (1999) Reporter cell lines to study the estrogenic effects of xenoestrogens. *The Science of the Total Environment*, 233, 47-56.
- 101) Jorgensen, M., Vendelbo, B., Skakkebaek, N.E. and Leffers, H. (2000) Assaying estrgenicity by quantitating the expression levels of endogenous estrogen-regulated genes. *Environmental Health Perspectives*, 108, 5, 403-412.