



"*Daphnia* sp., Acute Immobilisation Test and Reproduction Test"

The following Test Guideline includes two parts:
Part I - the 24h EC 50 acute immobilisation test
Part II - the reproduction test (at least 14 days).

PART I - 24 H EC 50 ACUTE IMMOBILISATION TEST

1. INTRODUCTORY INFORMATION

- Prerequisites

- Water solubility
- Vapour pressure

- Guidance information

- Structural formula
- Purity of the substance
- Methods of analysis for the quantification of the substance in water
- Chemical stability in water and light
- n-Octanol/water partition coefficient
- Results of a test on ready biodegradability (see Test Guideline 301)

- Qualifying statement

For chemicals of low solubility under test conditions, it may not be possible to quantitatively determine the EC 50.

- Standard documents

See references (1) to (6), Part II, Section 4, Literature.

**"Daphnia sp., Acute Immobilisation Test
and Reproduction Test"**

2. M E T H O D

A. INTRODUCTION, PURPOSE, SCOPE, RELEVANCE, APPLICATION AND LIMITS OF TEST

- D e f i n i t i o n s

24h EC 50 is the concentration estimated to immobilise 50 per cent of the Daphnia after 24 hours exposure. (If another definition is used, this must be reported, together with its reference.)

Immobilisation: those animals not able to swim within 15 seconds after gentle agitation of the test container are considered to be immobile. (If another definition is used, this must be reported, together with its reference.)

- R e f e r e n c e s u b s t a n c e s

In the acute immobilisation test a reference substance may be tested for EC 50 as a means of assuring that the test conditions are reliable.

- P r i n c i p l e o f t h e t e s t m e t h o d

In the acute immobilisation test a range of concentrations of the substance investigated exerts different degrees of toxic effects on the swimming capability of Daphnia under otherwise identical test conditions. Certain concentrations result in certain percentages of Daphnia being no longer capable of swimming at 24 hours. The test can be extended to 48 hours if desired.

- C o n d i t i o n s f o r t h e v a l i d i t y o f t h e t e s t

- In the control, not more than 10 per cent of the Daphnia should have been immobilised or trapped at the surface of the water.
- The dissolved oxygen concentration at the end of the test should be ≥ 60 per cent of the air saturation value at the temperature used.

**"*Daphnia* sp., Acute Immobilisation Test
and Reproduction Test"**

B. DESCRIPTION OF THE TEST PROCEDURE

• Preparations

Equipment

Equipment which will come into contact with the test solutions should preferably be all-glass; this glassware should be cleaned with solvents known to remove previously tested chemicals.

Holding and dilution water

Any water suitable for culturing *Daphnia*, either natural or reconstituted water, can be used in this test. To avoid the necessity of adaptation prior to the test, it is recommended that the water used in the test be similar to the culture water. Examples of reconstituted water are given in references (1), (2), (4) and (8).

• Experimental animals

Selection of species

Daphnia magna, or any other suitable *Daphnia* species, not more than 24 hours old at the beginning of the test, laboratory bred, apparently healthy and with a known history (breeding method, pretreatment) are used in this test. It is advisable to use the same species in Part I and Part II of this test.

• Performance of the test

- At least 20 animals, preferably divided into four groups of five animals each, should be used at each test concentration and for the controls.
- The *Daphnia* should not be fed during the test.
- Loading: at least 2 ml of test solution should be provided for each animal.
- The test temperature should be between 18 and 22°C, and for each single test it should be constant within $\pm 1^\circ\text{C}$.

**"*Daphnia* sp., Acute Immobilisation Test
and Reproduction Test"**

- A light-dark cycle is optional: complete darkness is acceptable.
- The concentrations tested are in a geometric series. The highest concentration tested should preferably result in 100 per cent immobilisation. It should, however, not exceed 1 g/l. The lowest concentration tested should preferably give no observable effect. The concentrations may be either measured or nominal, i.e. calculated, based upon the amount of material used in preparing the solution.
- Solutions are preferably made up without the use of a solubilising agent. Solutions of test substances of low aqueous solubility may be prepared by mechanical dispersion or, if necessary, by use of vehicles such as organic solvents, emulsifiers or dispersants of low toxicity to *Daphnia*. When such vehicles are used, one control should be exposed to the concentration of the vehicle used with the highest concentration of the test substance. The concentration of organic solvents, emulsifiers or dispersants should not exceed 100 mg/l. In evaluating the data it must be borne in mind that the results may be due to the combined effects of the substance itself and of the vehicle, which cannot normally be distinguished experimentally.
- The dilution water should be aerated prior to the addition of the test substance, and the oxygen concentration of the controls and the test solutions should be measured at the beginning and the end of the test.
- Volatile substances should be tested in completely filled closed containers, large enough to prevent lack of oxygen.
- The pH of the controls and the test solutions should be measured at the beginning and the end of the test; the pH of the test solutions should not be adjusted.

**"*Daphnia* sp., Acute Immobilisation Test
and Reproduction Test"**

3. DATA AND REPORTING

- Treatment of results

The percentage immobility at 24 hours and, if determined, at 48 hours is plotted against concentration on logarithmic-probability paper. Normal statistical procedures are then employed to calculate the EC 50 for the appropriate exposure period (see references, Part II, Section 4, Literature). Confidence limits ($p = 0.95$) for the calculated EC 50 values can be determined using the standard procedures quoted.

Where the data obtained are inadequate for the use of standard methods of calculating the EC 50, the highest concentration causing no immobility and the lowest concentration producing 100 per cent immobility should be used as an approximation for the EC 50 (this being considered the geometric mean of these two concentrations). In this case, the ratio of the higher to the lower concentration should not exceed 2.

- Test report

See pages 12-14 of this Test Guideline.

PART II - REPRODUCTION TEST AT LEAST 14 DAYS

The results of the acute immobilisation test are to be used to determine, with judgement, the concentration levels to be used in the reproduction test. It is suggested that this reproduction test be carried out using a geometrical concentration series of at least five concentrations with a factor of at most 10, starting at about the 24h EC 50 and ending at about 1/100 of the 24h EC 50. If necessary, lower concentrations should be tested.

**"Daphnia sp., Acute Immobilisation Test
and Reproduction Test"**

1. INTRODUCTORY INFORMATION

- **Prerequisites**

- Water solubility
- Vapour pressure
- Chemical stability in water and light
- Results of a test on ready biodegradability (see Test Guidelines 301 A-E)
- 24h EC 50 in Daphnia

- **Guidance information**

- Structural formula
- Purity of the test substance
- n-Octanol/water partition coefficient
- Method of analysis for the quantification of the test substance in water

- **Qualifying statement**

- For chemicals with low solubility under test conditions, it may not be possible to quantitatively determine the EC 50.

- **Recommendations**

- Instead of a test of two weeks duration in which three batches of young should be born per female, a test of three or four weeks may be preferred in order to obtain a more thorough judgement of the influence of the test substance on mortality and reproduction: in this period about six to nine batches of young should be born per female.
- It is recommended that a statistical test (such as an analysis of variance) be used to determine whether the test replications can be analysed together.

**"*Daphnia* sp., Acute Immobilisation Test
and Reproduction Test"**

- Standard documents

See references (1) to (6), Section 4, Literature.

2. METHOD

A. INTRODUCTION, PURPOSE, SCOPE, RELEVANCE, APPLICATION AND LIMITS OF TEST

- Definitions

Semi-static test is a test without flow of solution, but with occasional batchwise renewal of test solutions after prolonged periods (e.g. 24 hours).

Flow-through test is a test in which water is renewed continuously in the test chambers, the test substance being transported with the water used to renew the test medium.

EC 50 is the concentration of the test substance in water estimated to result in a 50 per cent reduction in reproduction of the *Daphnia* within a particular period of exposure (which must be stated).

- Reference substances

No reference substances are recommended for the reproduction test. Nevertheless, if a reference substance has been tested in the acute test, the results should be given in the test report.

- Principle of the test method

In the reproduction test, effects on the mortality and the reproductive capacity and other signs of intoxication in *Daphnia* are determined to be used as indications of the toxicity of a substance dissolved in water. For this purpose, the test organisms are exposed to solutions containing the test substance in various concentrations for a period of not less than two weeks, and long enough for the development of at least three broods.

**"*Daphnia* sp., Acute Immobilisation Test
and Reproduction Test"**

The mortality, the time of the first production of young, the number of young born and the signs of intoxication observed are compared with the corresponding parameters in the controls.

• C o n d i t i o n s f o r t h e v a l i d i t y o f t h e t e s t

- The mortality in the controls should not exceed 20 per cent at the end of the test.
- The dissolved oxygen concentration (throughout the test) must have been ≥ 60 per cent of the air saturation value.
- The pH of the controls, and of at least the most concentrated solutions, must be known throughout the test; the deviation from the initial value should be ≤ 0.3 units.
- There must be evidence that the concentration of the substance being tested has been satisfactorily maintained (e.g. at least 80 per cent of the nominal concentration) over the test period. The results should be based on measured concentrations if the deviation from the nominal concentration is greater than 20 per cent.
- The first young should have been born in the controls after a maximum of nine days.
- The average cumulative number of young per female in the controls after three broods, should be ≥ 20 at a temperature of $20 \pm 1.0^\circ\text{C}$.
- If, at the highest concentration tested, no effect on reproduction is detected, this should be reported.

B. D E S C R I P T I O N O F T H E T E S T P R O C E D U R E

• P r e p a r a t i o n s

"*Daphnia* sp., Acute Immobilisation Test and Reproduction Test"

Equipment and material

Normal laboratory apparatus and equipment should be used. Equipment which will come into contact with the test solutions should preferably be all-glass; this glassware should be cleaned with solvents known to remove previously tested chemicals.

Dilution water

Any water, either natural or reconstituted water, can be used, provided that it will sustain good reproduction in *Daphnia* (see Conditions for the validity of the test, above). In addition, the dilution water should meet the criteria given in reference (7). Examples of reconstituted water are given in references (1), (2), (4) and (8).

• Experimental animals

Selection of species

Daphnia magna less than 24 hours old at the beginning of the test, laboratory bred, apparently healthy and with a known history (breeding method, pretreatment) are used in this test. Other *Daphnia* species may be used provided that the relevant reproduction parameters are comparable to those of *Daphnia magna*.

Feeding

In the 14-day reproduction test food (of any kind and in any quantity) that results in meeting the criteria of reproduction given in Conditions for the validity of the test, above, is acceptable. Overloading of the test solutions with food should be avoided in order to minimise sorption of the test substance. Log-phase unicellular green algae are generally suitable.

• Performance of the test

- This reproduction test should not be carried out in a static test system: either a semi-static or flow-through system should be used. The renewal period should be guided by the chemical analysis and (if applicable) the oxygen level in the test solution; the solutions

**"*Daphnia* sp., Acute Immobilisation Test
and Reproduction Test"**

should be renewed at least once every 48 hours (if desired, on Monday, Wednesday and Friday). If the renewal scheme is used, the glassware must be emptied and food residues removed at renewal. It is recommended that the glassware be rinsed with distilled water and kept in a coded series for the following renewal. Each test unit therefore consists of two vessels which are used alternately. If flow-through systems are used, these should be cleaned at intervals of at most twice a week.

- At least 40 animals, preferably divided into four groups of ten animals each, should be used at each test concentration and for the controls.
- The *Daphnia* should be fed at least daily.
- Loading: at least 40 ml of test solution should be provided for each animal.
- The test temperature should be between 18 and 22°C, and for each single test it should be constant within $\pm 1.0^\circ\text{C}$.
- A light-dark cycle is necessary: 8 hours darkness and 16 hours light are recommended.
- The concentrations tested are made up in a geometric series. Substances should not be tested at concentrations exceeding 1 g/l.
- Samples of the test solutions should be taken at the beginning and during the test: the actual concentration must not drop below 80 per cent of the nominal concentration. Aeration of the test solutions is permissible, unless this would cause the actual concentration of the test substance to drop below 80 per cent of the nominal concentration.
- Solutions are preferably made up without the use of a solubilising agent like organic solvent, emulsifiers and dispersants.
- Solutions of test substances of low aqueous solubility may be prepared by mechanical dispersion or, if necessary, by use of vehicles such as organic solvents, emulsifiers or dispersants of low toxicity to *Daphnia*. When such vehicles are used, one control should be exposed to the concentration of the vehicle used with the highest concentration of the test substance.

**"*Daphnia* sp., Acute Immobilisation Test
and Reproduction Test"**

- The concentration of organic solvents, emulsifiers or dispersants should not exceed 100 mg/l. In evaluating the data it must be borne in mind that the results may be due to the combined effects of the substance itself and of the vehicle, which cannot normally be distinguished experimentally.
- The test solutions should be aerated prior to the introduction of the test substance and *Daphnia*.
- When more than 20 per cent of the test substance would be lost through volatility, the test should be carried out either in an almost-closed flow-through system or in an enclosed container of sufficient size to ensure that the oxygen level does not fall below 60 per cent of the saturation value.
- The dissolved oxygen concentration in all test solutions should be checked once every 48 hours (if desired, every Monday, Wednesday and Friday).
- The pH of the controls and of at least the most concentrated solutions is checked before and after each renewal: if necessary the pH of the other solutions should also be checked. The results of these measurements are recorded.
- The live and dead *Daphnia* of the "parental" generation (P) are counted and the dead specimens removed preferably daily, but at least three times a week (Monday, Wednesday and Friday).
- The presence of eggs in the brood pouch, males or winter eggs must be recorded. The condition and size of the P generation should be visually compared with the controls.
- When the parental animals are about seven days old, the first young *Daphnia* emerge from the brood pouch, after which a new batch appears every two to three days. These batches are called "broods" of the F₁ (filial 1) generation.

**"*Daphnia* sp., Acute Immobilisation Test
and Reproduction Test"**

- The newborn young of the F₁ generation should be counted at least three times a week, with an interval of 48-72 hours (e.g. Monday, Wednesday and Friday) and their visually estimated condition recorded. After counting and examination, the young are poured away. The presence of eggs on the bottom of the test vessel from which no young have emerged is checked and recorded.
- Test duration: the minimum duration of the test is 14 days, in which period not less than three broods of the F₁ generation must have appeared in the controls. If this is not the case, the test must be continued until the third brood in the control is complete, which is normally achieved within 21 days.

3. DATA AND REPORTING

- Test report

For both parts of the test, report:

- Test substance: chemical identification data
- Test organisms: source of *Daphnia*, any pre-treatment, breeding method (including source, kind and amount of food, feeding frequency)
- Test method: description of or reference to the method used
- Test conditions:
 - vehicles and/or additives used and their concentrations. If it is observed that the stability or homogeneity of the test solution cannot be maintained, then care should be taken in the interpretation of the results and note made that these may not be reproducible
 - dilution water: source and chemical and physical characteristics, including at least hardness, pH, Ca/Mg ratio, Na/K ratio, alkalinity
 - test temperature

**"*Daphnia* sp., Acute Immobilisation Test
and Reproduction Test"**

- light quality, intensity and periodicity
 - all measurements of pH and oxygen level made during the test, preferably in tabular form
 - results and date of test performed with reference substance, if available
 - description of test vessels: volume of solution, number of test organisms per vessel, number of test vessels per concentration, any treatment of the test vessels, the introduction of the test substance in the dilution water
 - in case of renewal, the renewal procedure and scheme; in case of flow-through, the test substance delivery system and the flow-rate
 - if measured, the actual concentrations of the test substance and the dates of measurement
- Number and percentage of *Daphnia* that showed any adverse effect in the controls and in each treatment group at each observation period and a description of the nature of the effects observed (e.g. immobilisation, mortality) in tabular form
 - Description or reference to statistical procedures applied
 - Any other effects differentiating organisms in tests and controls.

For the 24 hours EC 50 acute immobilisation test, also report:

- The 24h EC 50, preferably with 95 per cent confidence limits, determined by a suitable method
- If possible, the slope of the concentration response curve with its 95 per cent confidence limits
- The highest concentration tested producing no immobile *Daphnia*
- The lowest tested concentration producing 100 per cent immobile *Daphnia*

**"*Daphnia* sp., Acute Immobilisation Test
and Reproduction Test"**

For the reproduction test, also report:

- The EC 50 (immobilisation) and EC 50 (reproduction) values as far as possible at 24 hours, 48 hours, 96 hours, 7 days, 14 days and at the end of the test, preferably with 95 per cent confidence limits, obtained either by computation or graphically and the method applied (for the determination a probit method should be used.)
- The length of time for the appearance of the first brood for each concentration
- The number of young alive in each test vessel at any given day at which counts were made (the minimum requirement is for counts thrice weekly)
- The number of dead young on each day of counting
- Source, kind and amount of food, feeding frequency

For each of the above a statistical analysis of the homogeneity of replicate results for each concentration should be made. If homogeneity is found, it should be determined, through an appropriate statistical analysis, whether a significant difference exists between the control and the test concentrations.

Then report:

- The highest concentration tested at which no significant difference is found versus the controls with respect to mortality, reproduction and other observed effects
- The lowest concentration tested with significant difference versus the controls

4. L I T E R A T U R E

- Standard procedures

1. ISO 6341: Water Quality - Determination of the Inhibition of the Mobility of *Daphnia magna* Straus (Cladocera, Crustacea) (15 March 1982).

**"Daphnia sp., Acute Immobilisation Test
and Reproduction Test"**

2. DIN 38 412, Part 1: Allgemeine Hinweise zur Planung, Durchführung und Auswertung biologischer Testverfahren, (June 1982), and DIN 38 412, Part 2: Bestimmung der Wirkung von Wasserinhaltsstoffen auf Kleinkrebse (Daphnien-Kurzzeitest), (L 11), Normenausschuss Wasserwesen (NAW) beim Deutschen Institut für Normung e.V. (DIN), Arbeitsgruppe "Biotests": Testverfahren mit Wasserorganismen (Gruppe L), Berlin (March 1982).
 3. U.S. EPA: Registration of Pesticides in the United States - Proposed Guidelines, *Federal Register*, 43, N° 132 (July 10, 1978).
 4. AFNOR T 90 301: Qualité d'eau - Détermination de l'inhibition de la mobilité de *Daphnia magna* Straus (Cladocera, Crustacea) (French standard - 1974).
 5. NEN 6501 (Dutch standard on acute test).
 6. NEN 6502 (Dutch standard on reproduction test).
- O t h e r
7. U.S. EPA: *Environmental Effects Test Guidelines*, ES-1, EPA 560/6-82-002, Office of Toxic Substances, U.S. Environmental Protection Agency, Washington, D.C (1982).
 8. D.M.M. Adema, in *Degradability, Ecotoxicity and Bioaccumulation: The Determination of the Possible Effects of Chemicals and Wastes on the Aquatic Environment*, Chapter 5, Government Publishing Office, The Hague (1980).
 9. D.M.M. Adema, *Hydrobiologia* 519 (2), 125-134 (1978).
 10. K.E. Biesinger and G.M. Christensen, *J. Fish. Res. Board of Canada* 29 (12), 1691-1700 (1972).
 11. Commission of the European Communities, Study D.8369, Inter-laboratory Test Programme concerning the Study of the Ecotoxicity of a Chemical Substance with respect to *Daphnia* (1979).
 12. J.H. Canton and D.M.M. Adema, *Hydrobiologia* 519 (2), 135-140 (1978).
 13. J. Finney, *Statistical Methods in Biological Assay*, Griffin, Weycombe, U.K. (1978).

**"*Daphnia* sp., Acute Immobilisation Test
and Reproduction Test"**

14. J.T. Litchfield and F. Wilcoxon, *J. Pharmacol, Exper. Ther.* 96, 99-113 (1949).
15. C.E. Stephan, in *Aquatic Toxicology and Hazard Evaluation: Methods for Calculating an LC 50*, (edited by F.L. Mayer and J.L. Hamelink), ASTM STP 634, pp. 65-84, American Society for Testing and Materials (1977).
16. R.W. Wimmer and M.P. Farrel, *J. Fish Res. Bd. of Canada* 33, 1685-1691 (1976).