

OECD Environment Health and Safety Publications

Series on Testing and Assessment

No. 50

on

**DETAILED REVIEW PAPER ON
AQUATIC ARTHROPODS IN LIFE CYCLE
AND TWO-GENERATION TOXICITY TESTS**

**Environment Directorate
ORGANISATION FOR ECONOMIC CO-OPERATION AND DEVELOPMENT**

**Paris
January, 2005**

Also published in the Series on Testing and Assessment:

No. 1, *Guidance Document for the Development of OECD Guidelines for Testing of Chemicals (1993; reformatted 1995)*

No. 2, *Detailed Review Paper on Biodegradability Testing (1995)*

No. 3, *Guidance Document for Aquatic Effects Assessment (1995)*

No. 4, *Report of the OECD Workshop on Environmental Hazard/Risk Assessment (1995)*

No. 5, *Report of the SETAC/OECD Workshop on Avian Toxicity Testing (1996)*

No. 6, *Report of the Final Ring-test of the Daphnia magna Reproduction Test (1997)*

No. 7, *Guidance Document on Direct Phototransformation of Chemicals in Water (1997)*

No. 8, *Report of the OECD Workshop on Sharing Information about New Industrial Chemicals Assessment (1997)*

No. 9, *Guidance Document for the Conduct of Studies of Occupational Exposure to Pesticides during Agricultural Application (1997)*

No. 10, *Report of the OECD Workshop on Statistical Analysis of Aquatic Toxicity Data (1998)*

No. 11, *Detailed Review Paper on Aquatic Testing Methods for Pesticides and industrial Chemicals (1998)*

No. 12, *Detailed Review Document on Classification Systems for Germ Cell Mutagenicity in OECD Member Countries (1998)*

No. 13, *Detailed Review Document on Classification Systems for Sensitising Substances in OECD Member Countries (1998)*

No. 14, *Detailed Review Document on Classification Systems for Eye Irritation/Corrosion in OECD Member Countries (1998)*

No. 15, *Detailed Review Document on Classification Systems for Reproductive Toxicity in OECD Member Countries (1998)*

No. 16, *Detailed Review Document on Classification Systems for Skin Irritation/Corrosion in OECD Member Countries (1998)*

- No. 17, *Environmental Exposure Assessment Strategies for Existing Industrial Chemicals in OECD Member Countries (1999)*
- No. 18, *Report of the OECD Workshop on Improving the Use of Monitoring Data in the Exposure Assessment of Industrial Chemicals (2000)*
- No. 19, *Guidance Document on the Recognition, Assessment and Use of Clinical Signs as Humane Endpoints for Experimental Animals used in Safety Evaluation (1999)*
- No. 20, *Revised Draft Guidance Document for Neurotoxicity Testing (in approval)*
- No. 21, *Detailed Review Paper: Appraisal of Test Methods for Sex Hormone Disrupting Chemicals (2000)*
- No. 22, *Guidance Document for the Performance of Out-door Monolith Lysimeter Studies (2000)*
- No. 23, *Guidance Document on Aquatic Toxicity Testing of Difficult Substances and Mixtures (2000)*
- No. 24, *Guidance Document on Acute Oral Toxicity Testing (2001)*
- No. 25, *Detailed Review Document on Hazard Classification Systems for Specifics Target Organ Systemic Toxicity Repeated Exposure in OECD Member Countries (2001)*
- No. 26, *Revised Analysis of Responses Received from Member Countries to the Questionnaire on Regulatory Acute Toxicity Data Needs (2001)*
- No. 27, *Guidance Document on the Use of the Harmonised System for the Classification of Chemicals Which are Hazardous for the Aquatic Environment (2001)*
- No. 28, *Guidance Document for the Conduct of Skin Absorption Studies (2004)*
- No. 29, *Guidance Document on Transformation/Dissolution of Metals and Metal Compounds in Aqueous Media (2001)*
- No. 30, *Detailed Review Document on Hazard Classification Systems for Mixtures (2001)*
- No. 31, *Detailed Review Paper on Non-Genotoxic Carcinogens Detection: The Performance of In-Vitro Cell Transformation Assays (draft)*

- No. 32, *Guidance Notes for Analysis and Evaluation of Repeat-Dose Toxicity Studies* (2000)
- No. 33, *Harmonised Integrated Classification System for Human Health and Environmental Hazards of Chemical Substances and Mixtures* (2001)
- No. 34, *Guidance Document on the Development, Validation and Regulatory Acceptance of New and Updated Internationally Acceptable Test Methods in Hazard Assessment* (in preparation)
- No. 35, *Guidance notes for analysis and evaluation of chronic toxicity and carcinogenicity studies* (2002)
- No. 36, *Report of the OECD/UNEP Workshop on the use of Multimedia Models for estimating overall Environmental Persistence and long range Transport in the context of PBTS/POPS Assessment* (2002)
- No. 37, *Detailed Review Document on Classification Systems for Substances Which Pose an Aspiration Hazard* (2002)
- No. 38, *Detailed Background Review of the Uterotrophic Assay Summary of the Available Literature in Support of the Project of the OECD Task Force on Endocrine Disrupters Testing and Assessment (EDTA) to Standardise and Validate the Uterotrophic Assay* (2003)
- No. 39, *Guidance Document on Acute Inhalation Toxicity Testing* (in preparation)
- No. 40, *Detailed Review Document on Classification in OECD Member Countries of Substances and Mixtures Which Cause Respiratory Tract Irritation and Corrosion* (2003)
- No. 41, *Detailed Review Document on Classification in OECD Member Countries of Substances and Mixtures which in Contact with Water Release Toxic Gases* (2003)
- No. 42, *Guidance Document on Reporting Summary Information on Environmental, Occupational and Consumer Exposure* (2003)
- No. 43, *Draft Guidance Document on Reproductive Toxicity Testing and Assessment* (in preparation)
- No. 44, *Description of Selected Key Generic Terms Used in Chemical Hazard/Risk Assessment* (2003)
- No. 45, *Guidance Document on the Use of Multimedia Models for Estimating Overall Environmental Persistence and Long-range Transport* (2004)

No. 46, *Detailed Review Paper on Amphibian Metamorphosis Assay for the Detection of Thyroid Active Substances (2004)*

No. 47, *Detailed Review Paper on Fish Screening Assays for the Detection of Endocrine Active Substances (2004)*

No. 48, *New Chemical Assessment Comparisons and Implications for Work sharing (2004)*

No. 49, *Report from the Expert Group on (Q)SARs on the Principles for the Validation of (Q)SARs (2004)*

No. 50, *Detailed Review Paper on Aquatic Arthropods in Life-Cycle and Two-Generation Toxicity Tests)*

© OECD 2003

Applications for permission to reproduce or translate all or part of this material should be made to: Head of Publications Service, OECD, 2 rue André-Pascal, 75775 Paris Cedex 16, France

ABOUT THE OECD

The Organisation for Economic Co-operation and Development (OECD) is an intergovernmental organisation in which representatives of 30 industrialised countries in North America, Europe and the Asia and Pacific region, as well as the European Commission, meet to co-ordinate and harmonise policies, discuss issues of mutual concern, and work together to respond to international problems. Most of the OECD's work is carried out by more than 200 specialised committees and working groups composed of member country delegates. Observers from several countries with special status at the OECD, and from interested international organisations, attend many of the OECD's workshops and other meetings. Committees and working groups are served by the OECD Secretariat, located in Paris, France, which is organised into directorates and divisions.

The Environment, Health and Safety Division publishes free-of-charge documents in nine different series: Testing and Assessment; Good Laboratory Practice and Compliance Monitoring; Pesticides and Biocides; Risk Management; Harmonisation of Regulatory Oversight in Biotechnology; Safety of Novel Foods and Feeds; Chemical Accidents; Pollutant Release and Transfer Registers; and Emission Scenario Documents. More information about the Environment, Health and Safety Programme and EHS publications is available on the OECD's World Wide Web site (<http://www.oecd.org/ehs/>).

This publication was produced within the framework of the Inter-Organisation Programme for the Sound Management of Chemicals (IOMC).

The Inter-Organisation Programme for the Sound Management of Chemicals (IOMC) was established in 1995 following recommendations made by the 1992 UN Conference on Environment and Development to strengthen co-operation and increase international co-ordination in the field of chemical safety. The participating organisations are FAO, ILO, OECD, UNEP, UNIDO, UNITAR and WHO. The World Bank and UNDP are observers. The purpose of the IOMC is to promote co-ordination of the policies and activities pursued by the Participating Organisations, jointly or separately, to achieve the sound management of chemicals in relation to human health and the environment.

This publication is available electronically, at no charge.

**For this and many other Environment,
Health and Safety publications, consult the OECD's
World Wide Web site (www.oecd.org/ehs/)**

or contact:

**OECD Environment Directorate,
Environment, Health and Safety Division**

**2 rue André-Pascal
75775 Paris Cedex 16
France**

Fax: (33-1) 45 24 16 75

E-mail: ehscont@oecd.org

TABLE OF CONTENTS

PREAMBLE.....	11
ACKNOWLEDGMENTS	13
1.0 EXECUTIVE SUMMARY.....	14
2.0 INTRODUCTION	15
2.1 PURPOSE OF AQUATIC ARTHROPODS LIFE-CYCLE AND TWO-GENERATION TOXICITY TESTS.....	15
2.2 ACRONYMS AND ABBREVIATIONS.....	16
3.0 AQUATIC INVERTEBRATE ENDOCRINOLOGY AND ENDOCRINE DISRUPTION.....	17
3.1 OVERVIEW OF AQUATIC INVERTEBRATE ENDOCRINOLOGY	17
3.1.1 Porifera.....	19
3.1.2 Cnidaria.....	19
3.1.3 Annelida.....	20
3.1.4 Mollusca.....	21
3.1.5 Insecta	21
3.1.6 Crustacea.....	22
3.1.7 Echinoderms	24
3.1.8 Summary	24
3.2 OVERVIEW OF ENDOCRINE DISRUPTION IN AQUATIC INVERTEBRATES	24
3.2.1. Endocrine Disruption in Freshwater Species	25
3.2.2. Endocrine Disruption in Estuarine and Marine Species	28
3.2.3 Summary	32
3.3 EXTRAPOLATION ISSUES	32
3.3.1 Extrapolation From Taxon to Taxon.....	32
3.3.2 Extrapolation from Individuals to Populations	33
4.0 AQUATIC ARTHROPODS IN THE EVALUATION OF POSSIBLE ENDOCRINE DISRUPTION.....	35
4.1 FRESHWATER ARTHROPOD SPECIES	36
4.1.1 Midge Larvae (Chironomus tentans and C. riparius).....	36
4.1.2 Amphipods (Hyalella, Gammarus)	37
4.1.3 Daphnids (Ceriodaphnia dubia, Daphnia magna, D. pulex)	39
4.2 ESTUARINE AND MARINE ARTHROPOD SPECIES.....	40
4.2.1 Copepods (Acartia, Tisbe, Nitocra, Tigriopus).....	40
4.2.2 Amphipods.....	41
4.2.3 Decapods (shrimp, crabs, crayfish, lobsters)	42
4.3 BRACKISH, FRESHWATER AND MARINE ARTHROPOD SPECIES.....	42
4.3.1 Americamysis bahia (Molenock, 1969).....	43
4.3.2 Holmesimysis costata (Holmes, 1900).....	44
4.3.4 Mysidopsis intii holmquist, 1957.....	44
4.3.5 Neomysis integer (Leach, 1814).....	45

4.3.6	Other mysid species	46
4.4	OTHER SPECIES	48
5.0	SELECTION OF AN APPROPRIATE TEST SPECIES	50
5.1	CRUSTACEANS AS REPRESENTATIVE AQUATIC INVERTEBRATES	50
5.2	MYSIDS AS REPRESENTATIVE CRUSTACEANS.....	51
6.0	CONSIDERATION OF APPROPRIATE TEST SYSTEMS FOR REPRODUCTIVE AND DEVELOPMENTAL TOXICITY TESTS	52
6.1	ROUTE OF EXPOSURE.....	52
6.1.1	Water.....	52
6.1.2	Sediment	53
6.2	CONCENTRATION SERIES.....	53
6.3	STATISTICAL CONSIDERATIONS	55
6.3.1	Hypothesis-Testing or Regression Analysis	57
6.3.2	Statistical Versus Biological Significance	60
6.4	DESCRIPTION OF ASSAY ENDPOINTS REFLECTIVE OF REPRODUCTIVE AND DEVELOPMENTAL IMPAIRMENT	62
6.4.1	Growth, morphological and behavioral alterations.....	62
6.4.2	Measures of reproductive performance.....	66
6.4.3	Biochemical measures	68
7.0	RESPONSE TO ECDYSTEROID AGONISTS AND ANTAGONISTS	73
8.0	ANDROGENIC AND ESTROGENIC RESPONSES.....	76
8.1	ANDROGENIC RESPONSES	76
8.1.1	Endpoint Sensitivity.....	76
8.1.2	Gender Differences	77
8.2	ESTROGENIC RESPONSES.....	78
9.0	RESPONSE TO OTHER HORMONAL DISTURBANCES.....	80
10.0	CANDIDATE PROTOCOLS FOR AQUATIC ARTHROPODS LIFE CYCLE AND TWO-GENERATION TOXICITY TESTS	82
10.1	PROTOCOLS USING DAPHNID SPECIES.....	82
10.2	PROTOCOLS USING CHIRONOMID SPECIES AND OTHER SEDIMENT-DWELLING ORGANISMS	83
10.3	PROTOCOLS USING COPEPOD SPECIES	83
10.4	PROTOCOLS USING MYSID SPECIES.....	94
10.5	CANDIDATE PROTOCOL FOR EDC-DETERMINATION STUDIES AND ADDITIONAL DATA NEEDS	97
10.5.1	Preferred species	97
10.5.2	Description of the method.....	97
10.5.3	Endpoints: appropriateness and preferred method for quantification.....	102
10.5.4	Exposure protocol.....	104
10.5.5	Results and reporting	108
11.0	DATA GAPS, RESEARCH NEEDS AND OTHER CONSIDERATIONS.....	111
11.1	SIGNIFICANT DATA GAPS	111
11.2	RESEARCH NEEDS.....	112
11.3	IMPLEMENTATION CONSIDERATIONS	113
11.4	ANIMAL WELFARE.....	113

11.5	RECOMMENDED EQUIPMENT AND CAPABILITY	114
11.6	TESTING WITH NON-NATIVE SPECIES	114
12.0	REFERENCES	114

Tables

Table 1.	Acronyms and abbreviations.....	16
Table 2.	Examples of hormone reported in Invertebrate Taxa. Modified from Summary table from Oehlmann and Schulte-Oehlmann (2003), unless indicated otherwise	19
Table 3.	Types of water delivery systems.....	55
Table 4.	Control data means and coefficients of variation (CV)	58
Table 5.	Example US EPA protocols for evaluating chronic toxicity in Aquatic Arthropods.....	85
Table 6.	Example OECD protocols for evaluating chronic toxicity in aquatic arthropods.....	89
Table 7.	Recommended test holding conditions for <i>Holmesimysis costata</i> and <i>Mysidopsis intii</i>	94
Table 8.	Recommended Mysid Life Cycle Toxicity Test Conditions	100
Table 9.	Mysid two-generation toxicity test conditions recommended for conducting tests on potential endocrine disrupting chemicals	105
Table 10.	Measurement of effects of three classes of hormones	112

Figures

Figure 1.	Diluter	54
Figure 2.	Power of a one-sided independent-samples t-test as a function of the percentage difference (δ) detected between the test and control means, with 5 replicates per treatment ($\alpha = 0.05$).....	58
Figure 3.	Coefficient of variation (%CV) of control mysid survival as a function of time. The solid line is the fitted regression; the dashed lines are the 95% confidence intervals.....	58

PREAMBLE

In 1998, a Task Force on Endocrine Disrupter Testing and Assessment (EDTA) was established at the request of OECD member countries. The EDTA Task Force is a Special Activity of the Test Guidelines Programme and its main objectives are to:

- identify the needs and prioritize the development of new and enhanced guidelines for the detection and characterization of endocrine disrupting chemicals;
- develop a harmonized testing strategy for the screening and testing of endocrine disrupters;
- manage validation work for newly developed and enhanced Test Guidelines as appropriate; and,
- provide practical tools for sharing testing results and assessments.

The need for new and updated test methods to detect and characterise endocrine disrupting chemicals has been expressed by the Task Force for the assessment of human health effects and environmental effects. At early meetings of the EDTA Task Force, it appeared that existing OECD Test Guidelines would insufficiently cover for endocrine-related effects, especially for the environment. Member countries decided to list test methods which could potentially cover effects of chemicals on the reproductive system (estrogen agonists/antagonists and androgen agonists/antagonists) and on the development (thyroid system), and proposed enhancements where needed. Invertebrate life-cycle tests, encompassing development and reproduction, are proposed as *in vivo* assays including endpoints that indicate mechanisms of adverse effects (endocrine and other mechanisms) and potential population damage.

This Detailed Review Paper (DRP) is intended to provide the current state-of-the-knowledge in the area of aquatic invertebrate endocrinology, with a particular focus on the evaluation of possible endocrine disruption in aquatic arthropods through life cycle and two-generation tests.

The content of this review is intended to describe test methods, already existing or in development, that could be used in testing chemicals for their potential to interact with the endocrine system of aquatic arthropods and likely to create population damage. However, the methods described do not focus on endocrine disruption in terms of endpoints measured, but rather on reproductive and developmental performance of individuals exposed. It is planned that test methods under development or enhanced, will be validated to establish their relevance (i.e. whether the test is meaningful and useful for the intended purpose) and their reliability (i.e. reproducibility of test results over time within and among laboratories).

General principles for the conduct of validation studies have been defined following the OECD Stockholm Conference on Validation and Regulatory Acceptance of New and Updated Test Methods in Hazard Assessment. The OECD draft Guidance Document No. 34 describes these guiding principles and addresses the important steps and aspects that must be considered prior to and during the validation process. They include: (i) the definition of the test method and related issues (e.g., purpose, predictions, endpoints, limitations), (ii) the design and conduct of the initial phase of the validation process leading to the optimisation of the test method (often referred to as the pre-validation), (iii) the design and conduct of

the broader, multi-laboratory follow-up validation work, based on the outcomes of the initial phases and aiming at accumulation of data on the relevance and reliability of the test method, and (iv) the overall data evaluation and subsequent validation study conclusion, keeping in mind the requirements of regulatory authorities for submission of information relating to new or modified test procedures. It also discusses the need for and the extent of an independent evaluation, or peer review, of validated test methods.

The planning and conduct of a validation study should be undertaken on a case-by-case basis since there may be several ways of assessing the validity of the method. As described in the draft Guidance Document No. 34, the validation process is sufficiently flexible so that it can be applied equally well to a wide variety of tests and procedures. The flexibility also applies regardless of whether tests are for health or environmental effects. Flexibility is also encouraged on issues such as the amounts of information required at each phase, the number of chemicals tested, when and to what extent to use blind testing, and the number of laboratories participating.

A Validation Management Group for Ecotoxicity Testing (VMG-eco) has been established at the OECD level to supervise the planning and conduct of experimental work in fish, birds, amphibians and invertebrates. This VMG-eco reports back to the Task Force on Endocrine Disrupters Testing and Assessment (EDTA). To discuss the technical details of the test methods, an *Ad hoc* Expert group on Invertebrate Testing was created in 2002 and met for the first time in October 2003 to discuss the various tests proposed and to prepare initial validation work.

The U.S. Environmental Protection Agency took the lead in preparing the initial and revised versions of this Detailed Review Paper for their national programme on endocrine disrupters. A draft of this document was circulated for comments in October 2002 to OECD member countries. Comments were received from several internationally recognised experts (see acknowledgement section). A revised draft taking into account comments received served as the basis for the present OECD Detailed Review Paper.

ACKNOWLEDGMENTS

The OECD Secretariat would like to acknowledge the contributions of primary authors and national experts who provided extensive comments on the draft version of this detailed review paper:

- Roy K. Kropp, Margaret R. Pinza, and Michael L. Blanton, as primary authors; Battelle (United States)
- Environmental Health Science Bureau, Health Canada (Canada)
- Chemicals Division, Danish Environment Protection Agency (Denmark)
- UFZ Center for Environmental Research, Chemical Ecotoxicology, Leipzig/Halle (Germany)
- ECT Ecotoxicologie, Flörsheim (Germany),
- Rheinisch-Westfälische Technische Hochschule, research Group Aquatic Ecotoxicology and Mathematical Modelling, Aachen (Germany)
- DIN working group on marine biotests and collaborators (Germany)
- Federal Environment Agency, Berlin (Germany),
- Water Research Centre and the Health and Safety Executive, United Kingdom
- ABC Laboratories, Inc.; Chemical Development Group; Columbia (MO, United States)
- Rohm & Haas Company; Toxicology Department; Spring House (PA, United States)
- U.S. Environmental Protection Agency Federal Advisory Committee; Endocrine Disruptor Methods Validation Subcommittee; Office of Science Coordination & Policy; Washington DC
- U.S. Environmental Protection Agency; Office of Research & Development; NHEERL Atlantic Ecology Division; Narragansett (RI, United States)

1.0 EXECUTIVE SUMMARY

i) Invertebrates (especially arthropods such as insects and crustaceans) constitute the vast majority of animal species on earth. Although many invertebrate toxicity test protocols are routinely used in regulatory toxicity testing, few have been designed with endocrine-specific endpoints in mind. Although the growth, reproduction, development, and other aspects of invertebrate physiology and life cycle are known to be regulated by endocrine control, the endocrine systems and the hormones produced and used in the invertebrate body are not directly analogous to those of vertebrates. For example, ecdysone is a steroid hormone that regulates growth and molting in arthropods, and exhibits some functional and structural similarities to estrogen. Therefore, methods for testing effects of potential endocrine disrupters on aquatic arthropods are relevant to assess the adverse consequences of such chemicals on development and reproduction, through life-cycle or two-generation tests.

ii) This paper reviews the endocrinology of aquatic invertebrates in general (annelids, mollusks, insects, crustaceans and echinoderms, mainly) and provides examples of endocrine disruption. Issues associated with extrapolation are also explained in [Section 3](#).

iii) The remaining sections of the paper focus on aquatic arthropods. [Section 4](#) lists those species that have been studied most and where test methods have sometimes been developed to assess reproductive and developmental impairments following exposure to chemical substances. This is followed by an evaluation of possible endocrine disruption in aquatic arthropods.

iv) Criteria for the selection of appropriate species are proposed in [Section 5](#). This is followed in [Section 6](#) by considerations of an appropriate test system, including the route of exposure and concentration series and statistical procedures. A description of assay endpoints, such as survival, growth rate, and specific reproductive output (time to first brood, viability of offspring, clutch size, and sex ratio) is provided in the same section. These endpoints should provide adequate quantitative information on the adverse consequences of a putative endocrine disrupting chemical to the invertebrate species considered.

v) Responses observed following exposure to ecdysteroid (ant-)agonists, androgenic and estrogenic compounds and other hormonal disturbances are presented in [Section 7](#), [Section 8](#) and [Section 9](#) respectively.

vi) A review of candidate protocols using various species of aquatic arthropods is provided in [Section 10](#); together with recommendations on preferred options for endpoint measurement and test conduct.

vii) Data gaps, research needs and implementation considerations are described in [Section 11](#); followed by references in [Section 12](#).

2.0 INTRODUCTION

2.1 PURPOSE OF AQUATIC ARTHROPODS LIFE-CYCLE AND TWO-GENERATION TOXICITY TESTS

1. Test methods proposed at Level 5 of the EDTA Conceptual Framework are meant *i)* to provide data on effects from endocrine and other mechanisms of action, and *ii)* to be used for assessment of adverse effects and potential population damage. To fulfill this purpose, tests are often longer-term studies designed to encompass critical life states and processes, a broad range of doses, and administration by relevant route of exposure. In addition, the effects associated with potential endocrine disrupters can be latent and not manifested until later in life or may not be apparent until reproductive processes occur in an organism's life history. Thus, tests for endocrine disruption often encompass two generations to address effects on fertility and mating, embryonic development, sensitive neonatal growth and development, and transformation from the juvenile life state to sexual maturity.

2. The results from such long-term tests should be conclusive in documenting a discernable cause-and-effect relationship of chemical exposure to measurable manifestation in the test organisms. The tests are generally expected to:

- Establish exposure/concentrations/timing and effects relationships;
- Be sensitive;
- Assess relevant endpoints, i.e. indicative of developmental and reproductive impairment;
- Include a dose range for full characterization of effects;
- Adhere to good laboratory practices; and
- Be suitable for validation.

3. Invertebrates (especially arthropods such as insects and crustaceans) constitute the vast majority of animal species on earth. Although many invertebrate toxicity test protocols are routinely used in regulatory toxicity testing, few have been designed with endocrine-specific endpoints in mind. Although the growth, reproduction, development, and other aspects of invertebrate physiology and life cycle are known to be regulated by endocrine control, the endocrine systems and the hormones produced and used in the invertebrate body are not directly analogous to those of vertebrates. For example, ecdysone is a steroid hormone that regulates growth and molting in arthropods, and exhibits some functional and structural similarities to estrogen. It has been reported that the vertebrate androgen testosterone acts as an ecdysteroid antagonist in a crustacean (Mu and LeBlanc 2002a). Also, the aromatase inhibitor fenarimol, which prevents the conversion of testosterone to the vertebrate estrogen, has been demonstrated to inhibit ecdysteroid synthesis and interfere with normal molting processes in a crustacean (Mu and LeBlanc 2002b). Therefore, a method for testing crustaceans for effects of potential endocrine disrupters is relevant to assess the adverse consequences of chemicals indicated to be endocrine active in screening and short-term assays and tests.

2.2 ACRONYMS AND ABBREVIATIONS

4. Table 1 lists the acronyms and abbreviations used in the DRP, with the exception of commonly used units, such as h for hour or L for liter. Each of the acronyms and abbreviations is also introduced at first use in the text.

Table 1. Acronyms and abbreviations

20-E	20-hydroxyecdysone
4NP	4-nonylphenol
AFDW	ash-free dry weight
ASTM	American Society for Testing and Materials
BpA	Bisphenol A
BPDH	black pigment-dispersing hormone
CHH	crustacean hyperglycemic hormone
CYP	cytochrome P450 enzyme
DAH	dark adapting hormone
DDT	dichlorodiphenyl trichloroethane
DES	diethylstilbestrol
DRP	detailed review paper
EC	effects concentration
EC₅₀	median effective concentration
EDC	endocrine-disrupting chemical
EPA	United States Environmental Protection Agency
GIH	gonad-inhibiting hormone
GSH	gonad-stimulating hormone
HPV-inerts	high production volume inert compounds
IC	inhibition concentration
IGR	insect growth regulator
JH	juvenile hormone
LAH	light-adapting hormone
LC	lethal concentration
LC₅₀	median lethal concentration
LOEC	lowest observed effects concentration
MAR	metabolic androgenization ratio
MIH	molt-inhibiting hormone
MSD	minimum significant difference
NIEHS	National Institute of Environmental Health Sciences
NOEC	no observed effect concentration
NPPE	nonylphenol polyethoxylate
OECD	Organization for Economic Cooperation and Development
OPPTS	EPA Office of Prevention, Pesticides and Toxic Substances
PCB	polychlorinated biphenyl
PoA	ponasterone A
SAB	Scientific Advisory Board
SAP	Scientific Advisory Panel
SDWA	Safe Drinking Water Act
TBT	tributyltin

3.0 AQUATIC INVERTEBRATE ENDOCRINOLOGY AND ENDOCRINE DISRUPTION

5. Invertebrates comprise 95% of the world's animal species (Wilson 1988), and certainly a larger percentage of the Earth's total animal abundance. Certainly, it is understood that hormones are important in controlling physiological processes in invertebrates (Lafont 2000, Oehlmann & Schulte-Oehlmann 2003). However, until recently invertebrates received little attention regarding potential endocrine disruption, the main exception being studies of imposex in gastropods. Invertebrate endocrine systems, except perhaps those of arthropods, are generally poorly studied compared to those of vertebrates and this limits ability to evaluate potential EDCs (Lafont 2000, Oehlmann and Schulte-Oehlmann 2003). Because many invertebrate endocrine systems are not very well understood, some changes to endocrine systems after exposure to a chemical may not be detected or may not be measurable (Oehlmann and Schulte-Oehlmann 2003). However, the situation is improving with many studies being done within the last few years on such groups as sponges, corals, polychaetes, and echinoderms (Section 3.1), adding to what has been learned about arthropod endocrinology. Nonetheless, to gain a full understanding of the complex endocrine disruption issue, invertebrates must be included in a tiered testing approach (Vandenbergh et al. 2003).

3.1 OVERVIEW OF AQUATIC INVERTEBRATE ENDOCRINOLOGY

6. There are several recent general reviews that provide good summaries of invertebrate endocrinology (LeBlanc et al. 1999, Lafont 2000a, Feix and Hoch 2002). Additionally, there are some reviews that pertain to specific groups of invertebrates including Cnidaria (Leitz 2001), Annelida (Hardege 1999, Andries 2001, Salzet 2001), Insecta (e.g., Lafont 2000b), and Crustacea (Fingerman 1987, 1997, Huberman 2000, Subramoniam 2000). The intent of this section is to provide a brief overview of invertebrate endocrinology. More details can be found in the review articles.

7. As animal body plans increased in complexity from simple cell-based to organ system-based structures, the need for a means to coordinate internal processes became more important. Neural networks are developed in the Cnidaria, one of the more primitive invertebrate groups (Lafont 2000a). The complexity of the coordination systems increased through the invertebrate evolutionary line, progressing to the presence of endocrine cells in the Annelida, endocrine glands in the Mollusca, and perhaps culminating in complexity in the Arthropoda, although Lafont (2000a) offered that this conclusion might reflect the general lack of study of invertebrate endocrine systems. The complexity of coordinating systems is reflected in the pathways by which stimulation of the central nervous system (CNS) generates a response in a target organ (Lafont 2000a). The stimulus can induce the CNS to produce a neurotransmitter or a neurohormone. Both messengers can then act directly on the target organ or on an endocrine gland. The endocrine gland can then act on the target organ or on another endocrine gland, which then acts on the target organ.

8. There are two main types of chemical messengers (Lafont 2000a):

- Neurotransmitters may be fast- or slow-acting depending on whether the link to ion channels is direct or indirect (acetylcholine, glutamate); these are low molecular weight

molecules and have low diversity throughout invertebrates, i.e., they are evolutionarily conservative.

- Neurohormones/hormones include peptide/protein and lipid molecules that can be grouped by similarity of structure (e.g., steroids, peptides, terpenoids); these messengers show important evolutionarily diversification and either are related to vertebrate messengers or are specific to invertebrates.

9. Lafont made two key generalizations about invertebrate endocrine systems that are important to consider when evaluating endocrine disruption in invertebrates: (1) there are two sets of hormones in invertebrates—those that are similar to those found in vertebrates (i.e., they likely share a common ancestry) and those specific to inverts (e.g., ecdysteroids); (2) structurally related molecules may have different functions. This last generalization is very important and has been shown to be applicable to some studies of EDCs (Section 3.2)

10. LeBlanc et al. (1999) gave an introductory overview of animal lineages as they relate to endocrinology. Briefly, animals diverged from the ancestral “stem” line into two primary lineages, the protostomes and the deuterostomes. Traditionally, embryological features have been used to separate the two lineages. Major stem groups that will be discussed here are the sponges (Porifera) and jellyfish, hydroids, and corals (all Cnidaria). Major Protostome groups include worms (Annelida), snails and clams (Mollusca), midges (Insecta), and amphipods, daphnids, copepods, mysids, and decapods (Crustacea). The primary deuterostomes mentioned here are sea stars and feather stars (Echinodermata) and fish (Vertebrata). This divergence into two main lineages corresponds to important endocrinological differences. Protostomes primarily rely on neuropeptides to regulate physiological processes, although more advanced groups (insects, crustaceans) have increased reliance on ecdysteroids and terpenoids. Invertebrate deuterostomes (echinoderms) rely more on vertebrate-like steroids (estrogens, androgens, progesterone) and terpenoids, but do not have ecdysteroid hormones.

11. LeBlanc et al. (1999) also discussed the endocrinology of the major invertebrate groups. Many studies of individual invertebrate groups have identified single hormones or groups of hormones. Examples of these studies, selected primarily to build on the LeBlanc et al. (1999) review, are discussed in the following sections. The groups are presented in approximate phylogenetic sequence. Oehlmann and Schulte-Oehlmann (2003) prepared a table of the hormones that have been identified for the major invertebrate taxa, and the processes they control. Table 2 is a modification of their table that has been updated to include some additional taxa and hormones that have been identified recently.

Table 2. Examples of hormone reported in Invertebrate Taxa. Modified from Summary table from Oehlmann and Schulte-Oehlmann (2003), unless indicated otherwise

Taxon	Hormone Type	Example	Controlled Process
Porifera	Unknown	Unknown	Unknown
Cnidaria	Neuropeptides Thyroids Retinoids Steroids	GLWamides Thyroxine 9- <i>cis</i> -retinoic acid 17 β -estradiol ¹	Metamorphosis Strobilation Strobilation Reproduction
Nematoda	Ecdysteroids Terpenoids Neuropeptides	Unknown Juvenile hormone-like FMRFamide	Unknown Growth Unknown
Annelida	Ecdysteroids Neuropeptides Terpenoids	Ecdysone FMRFamide Gonadotropin ² Eicosatrienoic acid ³ Aracidonic acid ³	Unknown Neuromodulation Vitellogenesis Metamorphosis
Mollusca	Ecdysteroids Steroids Terpenoids Neuropeptides	Unknown Testosterone, 17 β -estradiol, progesterone JH reported APGWamide, dorsal body hormone, FMRFamide, egg-laying hormone, molluscan insulin-like peptides	Unknown Sexual differentiation, prosobranch reproduction Questionable Sexual differentiation, gonad maturation, spawning, neuromodulation, growth, development, energy metabolism
Crustacea	Ecdysteroids Steroids Terpenoids Neuropeptides	Ecdysone 17 β -estradiol, testosterone, progesterone Methyl farnesoate Androgenic hormone Crustacean Hyperglycemic Hormone Molt-inhibiting hormone Vitellogenesis-inhibiting hormone	Molting, vitellogenesis Uncertain Metamorphosis Sexual differentiation, vitellogenesis inhibition Energy metabolism Ecdysteroid production Vitellogenesis
Echinodermata	Steroids Neuropeptides	Testosterone, 17 β -estradiol, estrone Gonad-stimulating substance Maturation-promoting factor	Vitellogenesis, gametogenesis, spawning Spawning Fertilization
Tunicata	Steroids Neuropeptides Thyroids	Testosterone, 17 β -estradiol GRH analog Thyroxine	Gametogenesis, spawning Gonad development Tunic formation (?)

¹ Tarrant et al. (1999), Pernet and Ancill (2002). ² Gaudron and Bentley (2002). ³ Laufer and Biggers (2001).

3.1.1 *Porifera*

12. The current literature review did not identify any studies that have demonstrated the occurrence of hormones or endocrine functions in the sponges. However, one study (Hill et al. 2002) did examine the effects of three chemicals known to disrupt endocrine systems in some animals on freshwater sponges (see Section 3.2.1).

3.1.2 *Cnidaria*

13. The cnidarians are of particular interest because of the group's position as one of the stem invertebrate phyla that existed before the divergence into the protostome and deuterostome lineages. Cnidarians may, therefore, show aspects of endocrinology conserved in both groups (LeBlanc et al. 1999).

Also, some taxa within the phylum are of particular importance in tropical waters, e.g., coral reefs comprise a major ecosystem. Cnidarian neurosecretory cells may have been precursors to the evolution of the neurohormonal systems of higher animals (Feix and Hoch 2002).

14. Leitz (2001), in a thorough review of cnidarian endocrinology, reported that cnidarians do not possess defined endocrine glands, but that neurones are the major source of signaling compounds in cnidarians, although the target cells rarely have been identified. Leitz identified several major groups of regulatory compounds. Non-peptide regulatory compounds include catecholamines and their precursors (dopamine, adrenaline, norepinephrine), serotonin (5-hydroxytryptamine), taurine, and gamma-aminobutyric acid (GABA). The functions of most have not been identified clearly. Serotonin may be involved in metamorphosis, taurine may function in osmoregulation, and GABA may affect feeding.

15. Most of the peptide regulatory compounds are neuropeptides and inhibit or stimulate muscle contraction in hydrozoans and anthozoans (Leitz 2001). Gonadotropin-releasing hormones (GnRHs) comprise a peptide family that is conserved in length and amino-acid sequence composition (Ancil 2000). There is some evidence that they may also be active in other invertebrates (e.g., mollusks and tunicates). Ancil (2000) found evidence for GnRHs in the sea pansy *Renilla koellikeri* that are physiologically active and function in modulation of peristalsis (GnRH inhibits peristalsis), which is strongly enhanced during spawning. They were also found in starlet anemone *Nematostella vectensis*.

16. Retinoids occur in cnidarians. Retinoic acid X receptors (RXR) are nuclear hormone receptors found in vertebrates, echinoderms, arthropods, nematodes (Kostrouch et al. 1998). Kostrouch et al. found jellyfish RXR (jRXR), which is a close homolog of vertebrate RXR, in the jellyfish (*Tripedalia cystophora*). This hormone targets genes that encode soluble crystallins in lens of eye (as in vertebrates), which suggests that cnidarians are ancestral to other phyla, not an independent offshoot.

17. Sex steroids were identified in cnidarians very recently. Pernet and Ancil (2002) studied the sea pansy, which is dioecious and forms separate male and female colonies. Pernet and Ancil discovered the vertebrate estrogen 17- β estradiol (E_2) in all colonies and found that the levels varied through the reproductive cycle with a strong peak in March (at the onset of maturation) and June (start of the spawning period). These observations argue that 17- β estradiol has a role in the reproductive biology of the sea pansy. Tarrant et al. (1999) identified 17- β estradiol in a scleractinian coral (*Montipora capitata*) and thought that it functioned in reproduction. Corals contain a variety of steroids (Tarrant et al. 2003 among others), but there have been few studies on steroid metabolism. Water-born estrogens can be taken up by corals (Tarrant et al. 2001), metabolized (Tarrant et al. 2003), and can affect coral physiology (Tarrant et al. 2004). The presence of a vertebrate estrogen (17- β estradiol) in cnidarians is important for eventual understanding of evolution of hormonal systems.

3.1.3 *Annelida*

18. All hormones that have been identified in annelids to date are neuropeptides secreted by neurosecretory cells located primarily in the head (LeBlanc et al. 1999; Salzet 2001). A cardioactive peptide, FMRFamide (first found in mollusks), has been found in polychaetes, but not oligochaetes; ecdysteroids occur in some annelids, but no function for them has been determined; juvenile hormone affects larval settling when studied in the laboratory. Gaudron and Bentley (2002) discovered that the prostomium produces a gonadotrophic hormone that controls vitellogenesis in the oocytes. The hormone production is under environmental control. A second hormone induces oocyte maturation.

19. Andries (2001) reviewed the endocrine regulation of reproduction in polychaete worms. A single brain hormone controls reproductive development in worms that breed once and then die (semelparous reproduction) by inhibiting gonad maturation and promoting somatic growth. Hormone levels remain

consistent throughout the extended period of gamete development and the population of oocytes, which includes various stages of development during the one- to three-year period, becomes homogeneous only when the hormone levels decrease. The hormone apparently also inhibits spermatogenesis. Pheromones are involved in the timing of broadcast spawning (Hardege 1999, Andries 2001). Reproduction in worms that breed annually or continuously (iteroparous reproduction) is regulated by a combination of environmental factors and hormones (e.g., supraesophageal hormone stimulates ovarian-protein synthesis).

20. Laufer and Biggers (2001) reviewed the role of methyl farnesoate and juvenile hormone-active fatty acids in annelid metamorphosis and reproduction. They reported the *Capitella* trochophores respond very quickly to exposure to methyl farnesoate and eicosatrienoic acid (sperm maturation factor). Eicosatrienoic acid is present in adults of *Arenicola* and was proposed as a hormone that functions in metamorphosis and also induces spawning.

3.1.4 *Mollusca*

21. Neurosecretory centers, which produce neuropeptides, occur in the cerebral, pleural, pedal, abdominal ganglia of the central nervous system and comprise the molluscan endocrine system (LeBlanc et al. 1999). Typical invertebrate steroids (e.g., ecdysone) have been reported only rarely to occur in mollusks. Vertebrate-type steroids (testosterone, progesterone) can be synthesized from precursors in the ovotestes, which underscores the hermaphroditic character of many mollusks.

22. Ecdysteroids and juvenoids occur in some mollusks, but their particular functions have not been determined. The neuropeptide FMRFamide, a cardioacceleratory peptide that was first identified in mollusks, is one of the best known and most widespread neuroendocrine hormones. It regulates several physiological processes. Other peptides are involved in gonad maturation and egg production (egg-laying hormone), the development of female accessory sex organs, gonad maturation and ovulation (dorsal body hormones), and in growth, development, and metabolism (molluscan insulin-like peptides).

3.1.5 *Insecta*

23. Insect endocrinology, which is probably better understood than that of any other invertebrate group, has been thoroughly reviewed (e.g., LeBlanc et al. 1999, Lafont 2000b). Three types of structures comprise the endocrine system—neurosecretory cells, endocrine glands (epitracheal glands, corpora allata, prothoracic glands), and reproductive organs (LeBlanc et al. 1999). The neurosecretory cells produce neuropeptides that permit insects to respond to environmental factors that include food availability, temperature, and photoperiod, among others. Many hormones, most of which are unique to arthropods, have been identified in insects. Some insect hormones are similar to vertebrate hormones, but their functions have not been identified yet.

24. Two of the key insect hormones that are very relevant to an EDC evaluation program are ecdysteroids and juvenile hormones. Ecdysteroids, which are compounds structurally related to ecdysone (Goodwin et al. 1978), comprise one of the more important groups of insect hormones because they are involved in growth and development (molting) and reproduction (Lafont 2000b). Ecdysteroids are secreted by the prothoracic gland and by endocrine cells in the gonads. Insects cannot synthesize cholesterol; therefore, ecdysteroids are prepared from ingested cholesterol and plant steroids (LeBlanc et al. 1999). The synthetic pathway leading to the production of ecdysteroids is not completely known, partly because unstable compounds are included in some of its early stages (Lafont 2000b). Lafont mentioned two of the main questions concerning ecdysteroids that have yet to be answered: (a) How many active ecdysteroids and ecdysteroid receptors are there? and (b) How do varying concentrations of the hormones influence their action?

25. Juvenile hormones (JH) are terpenoids produced by the corpora allata that primarily regulate metamorphosis and reproduction. The presence or absence of JH determines the type of molt that occurs in the insect. The presence of JH during the initial rise of ecdysteroids in a larval stage will induce a molt to another larval stage (LeBlanc et al. 1999). When ecdysone levels start to rise during the final larval stage, the absence of JH will result in metamorphosis. In hemimetabolous species, this molt leads to the adult stage. In holometabolous species, JH again is present during this molt and the larva transforms into a pupa. The pupa molts to the adult in the absence of JH. JH biosynthetic pathways are known and are reviewed by Lafont (2000b). There are three main types of JH, called JHI, JHII, and JHIII, and several related compounds, including methyl farnesoate (MF) (Lafont 2000b). Lafont described a possible evolutionary scenario for JH with primitive insects (cockroaches) using MF and JHI and the most advanced insects (flies) using a unique form of JHIII (JHIII bisepoxide).

26. Many of the insecticides used to control outbreaks of agricultural pests have been formulated to interact with either of these two hormones. Many of these compounds, especially JH analogs, are not species-specific in action; thus they have the potential to impact nontarget animals severely. Therefore, these compounds are often the focus of endocrine disruption studies.

3.1.6 *Crustacea*

27. For crustaceans, biological processes are regulated by a complex endocrine system (Cuzin-Roudy and Saleuddin 1989). Study of this regulation began about 1921 with the publication of the results of experiments by R. Courrier that showed that a hormone produced by an endocrine gland, not the testes, determined male secondary sex characteristics (Fingerman 1997). Fingerman's (1997) historical review traces the passage of crustacean endocrinology from its early development by emphasizing key discoveries such as the nature and role of the sinus gland and other glands, and the roles of specific hormones in crustacean biology. Many of the descriptions are enhanced by the author's personal experiences gained through 40 years of active research on crustacean endocrine systems. Crustacean endocrinology has been reviewed about every decade since the early studies of the 1920s and 1930s. Among the relatively recent reviews, Quackenbush (1986) reviewed studies of the four types of compounds that help regulate crustacean biology—peptides, steroids, terpenoids, and biogenic amines. Later reviews have approached crustacean endocrinology by examining physiological processes, particularly growth and reproduction (Charmantier et al. 1997, Chang 1997, Subramoniam 2000). Chang (1993) compared endocrine control of molting and reproduction in crustaceans to that in insects.

28. Basically, inputs from the environment are integrated by a central nervous system. Neurotransmitters and neuromodulators govern the release of neuropeptides, which govern the production of hormones by the endocrine glands (Cuzin-Roudy and Saleuddin 1989). Molting, for example, is controlled by the release of molting hormones, which are ecdysteroids, and by neurosecretions for the central nervous system, which are accumulated and released by the sinus gland. The main endocrine centers for crustaceans described to date include the Y-organ, mandibular organ, androgenic gland, X-organ, and sinus gland.

29. Peptide hormones in crustaceans include compounds, such as red pigment concentrating hormone (RPCH) and pigment concentrating hormone, that affect the chromatophores and retinal pigments that comprise the complex color control systems found in crustaceans (Quackenbush 1986). Peptides also affect crustacean molting and reproduction. Molt-inhibiting hormone (MIH) is one factor that regulates molting. Peptides function in the control of vitellogenesis in crustaceans. Crustacean hyperglycemic hormone (CHH) is a peptide that regulates blood sugar, particularly glucose, in crustaceans (Quackenbush 1986). The hormone is unique among crustaceans in that it is taxon-specific. For example, CHH produced by crayfish does not affect crabs.

30. Steroid hormones include the ecdysteroids, which are the molting hormones in crustaceans (Charmantier et al. 1997, Chang 1997). These growth-regulating hormones also function in the control of reproduction and embryogenesis (Subramoniam 2000). Ecdysteroids are synthesized by the ecdysial glands or Y-organs. The Y-organ secretes ecdysone, which is converted to 20-hydroxyecdysone, the active ecdysteroid in most crustaceans. Several studies have shown that the Y-organ in some brachyuran crabs also secretes 3-dehydroxyecdysone and 25-deoxyecdysone (summarized in Subramoniam 2000). 25-deoxyecdysone is the precursor to ponasterone A (PoA), the primary circulating ecdysteroid in the premolt stage of the crabs (Subramoniam 2000). Other sources for ecdysteroids are the ovary, epidermis, and the oenocytes (Delbecque et al. 1990). In many crustaceans, molting, and hence somatic growth, continue after maturity, with the result that the Y-organ is active in adults. For most crustaceans, growth and reproduction can be grouped into three functional categories. In the first category, represented by crab and lobster, reproduction occurs after a long intermolt period. The second category includes isopods and amphipods, the growth and reproduction of which are concurrent. The last category relates to the rapidly molting cirripedes, for which reproduction requires several molt cycles. Molting and limb regeneration are intertwined (Fingerman et al. 1998). When limb regeneration occurs, first a limb bud develops within a layer of cuticle, and then becomes free and unfolds when ecdysis occurs as part of the molting process. Synthesis and secretion of ecdysteroid by the Y-organs is inhibited by molt-inhibiting hormone, a peptide that is released from the sinus gland.

31. Among the mysids, there is synchronization between reproduction and molting. Accumulation of ovary ecdysteroid takes place during the premolt stage, when the hemolymph ecdysteroid levels rise sharply. It is presumed that the hemolymph ecdysteroids are transported to the ovary along with the yolk precursor material. This trend is seen in other species as well, and shows that the Y-organ is active during premolt and that it produces ecdysteroids that are transported to the ovaries. This observation was confirmed by Subramoniam (2000). A Y-organ ablation was performed on the shrimp, *Lysmata seticaudata*, which caused a subsequent depression of vitellogenin synthesis and retardation in ovarian growth. Further findings on the same shrimp revealed a failure of folliculogenesis, which is a necessary prerequisite for vitellogenin-uptake by oocytes during secondary vitellogenesis.

32. Terpenoids, which are unique to arthropods (Quackenbush 1986), include methyl farnesoate (MF). MF is secreted by a mandibular organ and there is evidence that this compound is involved with the control of ecdysteroid synthesis. When a mandibular organ was experimentally implanted into the shrimp, *Penaeus setiferus*, there was a subsequent shortening of the molt cycle (Subramoniam 2000). Secretion by the Y-organ is controlled by methyl farnesoate, whereas inhibition is exercised by MIH from the X-organ sinus gland. The mandibular organ has also been implicated in the control of reproduction in crustaceans. Mandibular organ implants stimulated ovarian growth in the juvenile spider crab females and methyl farnesoate levels increased in the hemolymph and the mandibular organ during vitellogenesis in the crab, suggesting that this compound has a gonadotropic role similar to that of JH in insects. However, other studies showed no methyl farnesoate-level effects within the vitellogenic period in the lobster, for example (Subramoniam 2000).

33. Serotonin (5-hydroxytryptamine, 5-HT), a biogenic amine, is one of the most important biologically active substances in animal kingdom as it regulates many physiological and behavioral functions (Moreau et al. 2002). 5-HT has been found in many invertebrate groups, including crustaceans. Moreau et al. (2002) documented its presence in mysids, although they did not study its specific function. Studies reviewed by Quackenbush (1986) suggested that 5-HT induces the release of molt-inhibiting hormone and crustacean hyperglycemic hormone from the eyestalks of decapods.

34. Rather than providing a detailed review of crustacean endocrinology, the few paragraphs presented here summarize the four main types of compounds involved in the regulation of crustacean biology and show the interplay among them in regulating major physiological processes. In summarizing

about 75 years of crustacean endocrinological studies, Fingerman (1997) concluded that despite the many significant advances, work in the field “has really just begun.” This is especially true considering the tasks ahead in examining the potential for the disruption of crustacean endocrine systems by anthropogenic compounds.

3.1.7 Echinoderms

35. Echinoderms (e.g., sea stars, feather stars, sea cucumbers) are deuterostomes and are relatively closely related to vertebrates. Therefore, their endocrine systems may have some similarities to those of vertebrates and may share similar targets/susceptibilities to chemicals known to have endocrine-disruptive effects on vertebrates (LeBlanc et al. 1999). Vertebrate sex steroids may play a role in echinoderm reproduction (Oberdörster and Cheek 2001).

36. No ecdysteroids or juvenoids are known in echinoderms (echinoderms don't molt). Processes under hormonal, neurohormonal, or local growth factor control include gametogenesis, spawning, growth, and regeneration (reviewed in LeBlanc et al. 1999). Reproduction influenced by steroids and neuropeptides (summarized in LeBlanc et al.).

37. Sea stars synthesize two steroids, progesterone and testosterone, and steroidogenic pathway enzymes (3β -hydroxysteroid hydrogenase, cytochrome P450) occur in sea stars. Estrogen and estradiol synthesis has not been demonstrated, but a receptor for estradiol has been identified. High levels of progesterone in males have been demonstrated at the beginning of spermatogenesis. Therefore, it is very likely that steroids function in sea star reproduction. There is very little information about endocrine functions in other echinoderm groups.

38. Gonad-stimulating substance (GSS), which has been found in radial nerve extracts and stimulates spawning, may be a neuropeptide. GSS indirectly stimulates release of a maturation-promoting factor that readies oocytes for fertilization. Other neuropeptides have been discovered in echinoderms and may regulate feeding.

39. Regeneration is a form of asexual reproduction in some taxa, but also serves to replace lost body parts. Loss of body parts, which may be a defense from predation, most frequently involves arms (e.g., sea stars), but also often may involve sections of epidermis (e.g., sea cucumbers; Kropp 1982). Arm regeneration in feather stars (Crinoidea) occurs by the proliferation of migratory undifferentiated cells and is under nervous system control, which provides the primary regulatory factors (Candia Carnevali et al. 2001b).

3.1.8 Summary

40. The endocrine system of an invertebrate differs from that of a vertebrate organism in the type of endocrine glands present and in the chemical structure (and consequently in the function) of specific hormones that are produced. Invertebrates produce some hormones that vertebrates do not. For example, crustaceans and most other invertebrates produce di- and tri-iodothyronine, but have no thyroid gland, and the function of the thyronines is unknown. Invertebrates use hormones that are not found in vertebrates. Crustecdysone is in some ways analogous to a vertebrate's estrogen hormone, but it is structurally, functionally, and metabolically different from the vertebrate hormone (J.M. Neff, personal communication, January 15, 2002).

3.2 OVERVIEW OF ENDOCRINE DISRUPTION IN AQUATIC INVERTEBRATES

41. There are two basic types of compounds with the potential to disrupt endocrine systems: synthetic chemicals (xenobiotics) and natural plant chemicals (phytoestrogens) (Crisp et al. 1998). Some

examples of xenobiotics are compounds used in plastics (nonylphenol, bisphenol-A), PCBs, and some pesticides. Phytoestrogens include hormone-mimicking substances contained in some agricultural plants and in paper mill effluent. These can be estrogenic or antiestrogenic.

42. Lafont (2000) described and gives examples of four levels at which EDCs can disturb endocrine systems. At the first level, EDCs block the availability of the precursors required for the synthesis of hormones. The interruption of hormone biosynthesis occurs at the second level. For example, some chemicals inhibit cytochrome P450s, which catalyzed biosynthesis, thus breaking the synthetic pathway leading to hormone production (Lafont 2000). The third level at which an EDC can act is on hormone catabolic processes. In this case, an EDC could act to increase the rate at which a hormone is catabolized, resulting in lower levels in an animal. At the fourth level, an EDC directly interferes with the actions of hormones. EDCs can act as agonists by binding to a hormone receptor and acting as that hormone would to regulate gene transcription (LeBlanc et al. 1999). EDC also may act antagonistically by binding to a hormone receptor without inducing its activity (LeBlanc et al. 1999).

43. The following sections present a brief overview of endocrine disruption in invertebrates that builds on recent reviews by LeBlanc et al. (1999), Oehlmann and Schulte-Oehlmann (2003), and Segner et al. (2003) among others. The overview is not meant to be exhaustive and is organized by major taxon within habitat type. Some information about endocrine disruption in aquatic arthropods, especially crustaceans, is included here, but a more detailed discussion, structured in the context of hormonal effects, is presented in Sections 7, 8 and 9.

3.2.1. *Endocrine Disruption in Freshwater Species*

44. Porifera: Hill et al. (2002) tested the effects of three chemicals known to disrupt endocrine systems in some animals (ethylbenzene, nonylphenol, bisphenol-A) on two species of freshwater sponges (*Heteromyenia* sp., *Eunapius fragilis*). The study was initiated with gemmules and tested the effects of these chemicals on growth (morphology and rate). Hill et al. found that higher doses caused a morphological abnormality that was similar across chemicals for both species. Germination rates for gemmules in the control treatments were relatively low (70% and <50% for each species) and germination could be inhibited at highest concentrations. Hill et al. could not determine if the observed effects were attributable to a disruption of an endocrine pathway or a more general toxic effect, yet argued that the former was likely (despite the general lack of information about sponge endocrinology).

45. Hydrozoa: Pascoe et al. (2002) examined the effects of 17 α -ethinylestradiol and bisphenol A on the hydrozoan *Hydra vulgaris* and found no physical or physiological damage to the polyps at environmentally relevant concentrations (ng/L) of either chemical. However, they determined that toxicity occurred at relatively high chemical concentrations and concluded that signaling processes necessary for normal development, regeneration, and sexual reproduction were not affected by these estrogenic pollutants at low, environmentally relevant concentrations.

46. Cladocera: Baldwin et al. (2001) documented the effects of several EDCs on the *Daphnia magna*. One experiment focused on exposure to 20-hydroxyecdysone (20-E), the crustacean molting hormone, and to ponasterone A (PoA), an endogenous compound that has 20 times higher affinity for the ecdysone receptor. The 21-day exposure had little effect on reproduction for either compound, except at the highest concentrations. However, adults suffered high mortality rates and either did not produce broods or produced smaller broods. Second-generation effects were not observed as a result of 20-E exposure, but there was a slightly significant effect on reproduction from PoA exposure. The effect on reproduction could be attributed to the structure of PoA: it has fewer hydroxyl groups, and could be less easily metabolized as is 20-E. The resulting longer exposure could allow second-generation effects. It is also possible that the higher affinity for PoA to ecdysone receptor sites caused a limited effect on

secondary vitellogenesis in developing daphnids, which manifested itself as reduced reproduction because of smaller brood size (Baldwin et al. 2001).

47. Recently, many studies conducted by LeBlanc and coworkers have elucidated many aspects of daphnid endocrinology and have subsequently documented the impacts of EDCs. For example, Olmstead and LeBlanc (2002) determined that methyl farnesoate was the likely endocrine factor regulating the development of males in *D. magna*. They later predicted that a juvenile hormone analog, pyriproxyfen, would stimulate the production of males and found that this indeed occurred (Olmstead and LeBlanc (2003). Tatarazako et al. (2003) also found that methyl farnesoate and pyriproxyfen stimulated male production in *D. magna*, even under uncrowded, high-food (i.e., low stress) conditions. They also found that three additional juvenoids—fenoxycarb, methoprene, and JHIII—stimulated male production. The five juvenoids showed similar dose-dependant responses (reduction in fecundity, male production), although fenoxycarb and pyriproxyfen caused the effects at substantially lower concentrations than the other three compounds. Companion studies by Mu and LeBlanc (2002a, 2002b) showed that ecdysteroids are important in regulating daphnid development and that ecdysteroid antagonists, testosterone and fenarimol, interfered with normal development. Testosterone affects early and late developmental stages (acting as a hormone receptor antagonist), whereas fenarimol affected late development (acting as a hormone synthesis inhibitor). Mu and LeBlanc (2004), in a study with more significance to EDC evaluations, examined the potential synergistic effects of a fenarimol-testosterone mixture on daphnid development. Mu and LeBlanc focused on the different actions of each compound, predicting that fenarimol would lower ecdysone levels in daphnids, which would increase the testosterone binding to the ecdysone receptor with the net overall effect of increasing toxicity beyond that of either individual chemical. Their study confirmed this prediction and underscored the importance of considering the potential synergistic of mixtures of chemicals in the environment.

48. Kashian and Dodson (2004) studied the effects of several vertebrate hormones on sex determination and development in *D. magna*. Among the hormones they tested, only long-term (26 days) exposure to progesterone affected sex determination. The second clutch of young produced by *D. magna* exposed to 100 $\mu\text{g/L}$ progesterone contained significantly more males than controls. The effect disappeared with succeeding clutches. Daphnids exposed to 100 $\mu\text{g/L}$ testosterone had significantly reduced fecundity compared to control animals. Two hormones affected daphnid growth. Exposure to diethylstilbestrol (100 $\mu\text{g/L}$) significantly reduced growth, whereas exposure to gonadotropin (30 $\mu\text{g/L}$) significantly increased growth compared to controls. Resting egg production and molting were not affected by any of the vertebrate hormones tested.

49. Other studies of endocrine disruption in daphnids showed that styrene dimers and trimers reduced fertility in *Ceriodaphnia dubia* (Tatarazako et al. 2002) and the 17 α -ethinylestradiol decreased the number of offspring in *Daphnia magna* (Goto and Hiromi 2003). Goto and Hiromi also found that another contraceptive ingredient, norethindrone, did not affect offspring production or sex ratio in *D. magna*.

50. Amphipoda: Vandenbergh et al. (2003) used a multigenerational assay to examine the effects of sublethal doses of 17 α -ethinylestradiol, a synthetic estrogen, on sexual development in *Hyalella azteca*. They compared the second gnathopod size and reproductive tract histology among treated and untreated males and found that F1 males exposed to the lowest 17 α -ethinylestradiol doses, had significantly smaller second gnathopods than controls, whereas the second gnathopods of those exposed to higher doses did not. This suggested a U-shaped response where effects were observed at a low concentration, but were masked at high concentrations. Vandenbergh et al. also observed cellular abnormalities in males exposed to all concentrations. Among these were larger and more spherical germ cells, a hollow cell structure, and less dense cytoplasm, features that are analogous to female gonad morphology. Oocyte-like structures were detected at some concentrations and smaller spermatids, fewer vas deferens, and irregular spermatogonia were found at highest concentration. The observed histological abnormalities indicated some degree of

hermaphroditism in the exposed amphipods. The authors concluded that sublethal exposures of 17 α -ethinylestradiol may affect sexual development in *H. azteca*, but they also noted that test concentrations were higher than those reported to affect vertebrates and higher than those observed in natural environments.

51. Watts et al. (2001b) looked at the toxicity of 17 α -ethinylestradiol and bisphenol A and effects of the chemicals on the precopulatory guarding behavior of *Gammarus pulex*. They found that 17 α -ethinylestradiol was more toxic than bisphenol A, but that there was no disruption of precopula or the propensity to reestablish it except at the highest concentrations, which were close to doses that were determined to be acutely toxic. Therefore, these two xenoestrogens had no effect on any endocrine systems that facilitate the behavior. In a follow-on study, Watts et al. (2002) designed a longer study specifically to determine if the lack of response to 17 α -ethinylestradiol in the acute test predicted a similar lack of response at the population level in *G. pulex*. The results showed that all of the 17 α -ethinylestradiol concentrations that were tested showed population increases greater than those in control treatments (the control treatments did increase over the starting population values). The difference was attributed to the recruitment of neonates and juveniles. The dosed treatments also had more females than the control treatments and had a M:F ratio of ~0.4–0.5:1 versus the ratio of 1:1 attained in the control treatments. The authors concluded that exposure to 17 α -ethinylestradiol resulted in significant increase in population size, despite prediction of no effect on reproductive behavior that was derived from the acute assay. The increase in population size was attributed to an accelerated female maturation rate, such that more young were produced earlier, and an increase in number of females. These two studies underscore the need for a chronic assay to develop a more complete understanding of potential EDC effects on populations.

52. Insecta (*Chironomus*): Most of the studies of endocrine disruption in midge larvae has been done on *Chironomus riparius*. In the sole study found during this review that directly concerned endocrine disruption in *C. tentans*, Kahl et al. (1997) conducted an assay to examine the effects on 4-nonylphenol on the life cycle of the midge. The only significant result was on 20-d survival at highest concentration. There were no effects on egg production or viability, emergence time, and sex ratio. Kahl et al. (1997) noted that some egg cases were deformed, particularly at the high nonylphenol concentration.

53. Studies on potential endocrine disruption in *C. riparius* have considered two basic types of endpoints, mouthpart deformities and life cycle reproductive parameters. Deformities of the mouthparts (mentum, mandibles, epipharyngeal pecten) have been observed in association with some contaminated sediments. Physiological disturbance during molting are thought to cause the deformities. Meregalli et al. (2001) tested the hypothesis that hormonal disruptions during molting contributed to the development of the deformities by subjecting the midge larvae to the known endocrine disrupter 4-nonylphenol. Mentum deformities were significantly more common in larvae exposed to sublethal concentrations of nonylphenol than in the control larvae. The mandibles and epipharyngeal pecten showed few deformities. In a more broadly scoped study, Watts et al. (2003) studied the effects of sublethal doses of the estrogenic endocrine disruptors 17 α -ethinylestradiol and bisphenol A on molting and mouthpart structures in *C. riparius*. The highest concentration (1 mg/L) of both chemicals significantly reduced larval weight and delayed molting. Watts et al. noted that the high concentration was environmentally unrealistic and much greater than the concentration known to elicit a response in fish. The remaining doses of either chemical did not affect either parameter. The lowest concentration (10 ng/L) of each chemical caused deformities of the mentum, but not the other two primary mouthparts. 17 α -ethinylestradiol had a slightly stronger effect than bisphenol A on the mentum. No mouthpart deformities occurred at the highest concentration of each chemical. The mouthpart studies also highlights that classical dose-response curves often do not apply to EDCs, but that inverted U-shaped curves (no/low effects at the concentration extremes, high effects at middle concentrations) more accurately depict the response (Watts et al. 2003).

54. Watts et al. (2001a) used a two-generational study of *C. riparius* to evaluate the effects of 17 α -ethinylestradiol and bisphenol A on the midge life cycle. They found that the main effect of the two chemicals was delay of emergence at higher test concentrations, especially for the second generation. The delay did not affect the typical protandrous emergence pattern in chironomids (males emerge before females). There was no effect on egg production or egg viability. One interesting effect was the alteration of the second generation adult sex ratio where males outnumbered females 2:1 at all but the highest 17 α -ethinylestradiol concentration. Watts et al. thought that this effect was aligned with that expected for an estrogenic compound and suggested that this could mean the chemical did not act as an estrogen in this particular case. Bisphenol A did not affect sex ratio.

55. Hahn et al. (2001, 2002) tested the effects of tebufenozide, a pesticide that acts as a molting hormone agonist, on *C. riparius* development and vitellogenin/vitellin (Vg/Vn) levels. The molting hormone agonist was anticipated to accelerate development. Tebufenozide affected development, but that effect did not occur until the final molt from pupa to adult. Hahn et al. pointed out that this effect was counter to that desired for the pesticide's intended butterfly targets, stimulation of an early molt that leads to death. The effects levels determined during this chronic study (e.g., LC50 = 21.14 μ g/L) were much lower than those determined for acute exposures. The 2002 study also examined the effects of bisphenol A and 4-nonylphenol on Vg/Vn immunoreactivity. Males exposed to most contaminant concentrations showed reduced Vg/Vn immunoreactivity. Females were not affected by the contaminants, except the highest bisphenol A concentration reduced Vg/Vn levels. These results were not expected because bisphenol A and nonylphenol are known to stimulate vitellogenesis in vertebrates. The observed responses were not dose-dependent as all concentrations reduced Vg/Vn somewhat equally.

56. Hahn and Schulz (2002) found that relatively short-term exposure to tributyltin reduced ecdysteroid synthesis in female *C. riparius* larvae at test concentrations as low as 50 ng/L, whereas males showed increased biosynthesis at a concentration of 500 ng/L. Imaginal discs developed more slowly in females, but faster in males at all test concentrations than in the respective control treatments. The authors thought that these gender-specific differences showed that either the ecdysteroid pathways were impacted differently, or that different reactions occurred to the same impact.

3.2.2. Endocrine Disruption in Estuarine and Marine Species

57. Cnidaria: Tarrant et al. (2004) evaluated the effects of exogenous estrogens (17 β -estradiol, estrone) on scleractinian coral reproduction. They determined that 29% fewer egg bundles were released by *Montipora capitata* colonies in estradiol-treated colonies than in control colonies. Thus, estrogens seem to be natural bioregulators in corals, which suggests that exogenous estrogens can reduce coral fecundity. However, the number of eggs per bundle did not differ in treated corals than from corals in control treatments. Growth in *Porites compressa* was significantly reduced in treated corals at estrone levels only slightly greater than ambient levels. However, the mode of action of estrogen in corals is unknown. Tarrant et al. suggested that these results (and others for invertebrates) imply that the potential disruptive effects of estrogens are not limited to interruption of mammalian reproduction.

58. Annelida: The review prepared by LeBlanc et al. (1999) mentioned that there was only one known case of endocrine disruption in annelid worms; juvenile hormone and JH analogs stimulate larval settling and metamorphosis in *Capitella*. In a 78-d study, Hansen et al. (1999) found that the lowest treatment of sediment-bound 4-n-nonylphenol stimulated asymptotic body volume growth and increased mean brood size, but that these effects did not translate to changes in population growth rates. The highest concentration of nonylphenol significantly reduced several reproductive metrics, including brood size, volume-specific fecundity, time to first reproduction, and the growth rate of individuals. The authors did not link the observed effects to possible endocrine disruption, although they noted that the stimulation at the low dose was another case of hormesis, which has been documented for other organisms.

59. Mollusca: The most frequently cited examples of endocrine disruption in mollusks concern imposex, which is an irreversible condition in gastropod snails in which females develop secondary male sex organs (Matthiessen and Gibbs 1998). Imposex, which is most frequently a response to exposure to tributyltin contained in anti-fouling paints, may result from the inhibition of aromatase or other steroid precursors, or from direct impacts to neurohormones (Rotchell and Ostrander 2003). Since the topic has been the subject of very many studies and has been considerably reviewed, it will not be discussed in detail here. Reviews and other significant discussions of imposex are included in Matthiessen and Gibbs (1998), Evans and Nicholson (2000), the series of studies by Oehlmann and others (Oehlmann et al. 2000, Schulte-Oehlmann et al. 2000, Tillman et al. 2001), and Axiak et al. (2003).

60. Nice et al. (2003) studied the effects of single short-term (48 h) nonylphenol exposure of 7-d old larvae of oyster *Crassostrea gigas* on long-term physiological processes. Exposure to environmentally relevant levels of nonylphenol resulted in a sex ratio biased towards females and an increased incidence of hermaphroditism. The exposure also affected gamete viability, severely impacting embryonic and larval development in second generation.

61. Jobling et al. (2003) tested effects of 17 α -Ethinylestradiol, 4 *tert*-octylphenol, bisphenol A, and sewage effluent on egg and embryo production in a common freshwater European snail *Potamopyrgus antipodarum*. All estrogen and xenoestrogen treatments stimulated embryo production, except at the highest concentration, which had an inhibitory effect. These reproductive effects were generally similar to those the authors observed for three fish species (fathead minnow, rainbow trout, carp) in a companion study. Jobling et al. suggested that this snail is a sensitive species whose testing against estrogens may be relevant to estrogenic activity in vertebrates.

62. Crustacea: Marine crustaceans are important organisms to include in the evaluation of the adverse consequences of EDCs, and the selection of suitable species is the one focus of the present review. Early studies of the effects of EDCs on estuarine crustaceans, most of which have occurred within the last few years, have focused on three primary groups, barnacles, copepods, and decapods (Ingersoll et al. 1999, Hutchinson 2002). The results of these studies showed that some crustacean groups may be affected by exposure to EDCs, but others may not and, therefore, extrapolation of the results from testing one crustacean group to another is problematic.

63. *Barnacles*.—Billingham et al. (1998, 2000) examined the effects of two estrogens, 4-*n*-nonylphenol and 17 β -estradiol on larval settlement and the production of a larval storage protein (cyprid major protein, CMP) in *Balanus amphitrite*. Cyprids use CMP during settlement and the early post-settlement development. Because CMP is structurally related to vitellin, which is analogous to vitellogenin, it can be used as a biomarker of estrogen exposure in lower vertebrates. The expectation in these studies was that cyprid settlement might be affected by the stimulation of CMP synthesis after larval exposure to environmental estrogens. The results of the 1998 study, however, showed reduced settlement after exposure to both estrogens, but that the cause was not related to endocrine disruption. The second study (Billingham et al. 2000) measured levels of CMP and found that they were elevated after exposure of nauplii to low levels of the estrogens.

64. *Copepods*.—Hutchinson et al. (1999a, 1999b) found that exposure to several steroids had no effect on the survival and development of harpacticoid copepod (*Tisbe battagliai*) nauplii and cautioned against extending the reported effects of steroid exposure in some species of crustaceans to the group as a whole. At about the same time, Bechmann (1999) showed that high levels (>62 μ g/L) of nonylphenol were acutely toxic to *T. battagliai*, but that exposure to a low level (31 μ g/L) did not affect any of the measured life-table parameters (survival, sex ratio, fecundity) measured. Breitholtz and Bengtsson (2001) did not find evidence of endocrine disruption in the harpacticoid copepod *Nitocra spinipes* after exposure to the estrogens 17 β -estradiol, 17 α -ethinylestradiol, and diethylstilbestrol. Chandler and coworkers studied

potential endocrine disruption in a third harpacticoid species, *Amphiascus tenuiremis*. Bejarano and Chandler (2003) used chronic multi-generation exposures to evaluate the effects of the herbicide atrazine on reproduction and development in *A. tenuiremis*. While atrazine did not have significant effects on several parameters (e.g., time to maturity, time to egg extrusion, time to hatching), some concentration-related effects occurred. Reproductive failure (mating pairs unable to produce living offspring or females unable to extrude more than one brood) increased with atrazine dose. Nauplii production by F₀ females was reduced at the highest concentration and that by F₁ females was reduced at all concentrations. Both factors combined to reduce total population growth of the F₁ generation at doses lower than those considered “safe” for chronic exposures (26 µg/L). Chandler et al. (2004) found that sublethal, environmentally relevant concentrations of the insecticide fipronil delayed the maturation of *A. tenuiremis* copepodites to adults and reduced or virtually eliminated the production of young. These effects were modeled and predicted a 62% decline in the population size of the copepod at the lowest concentration tested (0.16 µg/L).

65. Copepods probably have been tested against more potential EDCs than any other marine group. For example, Anderson et al. (2001) tested the effects of 14 compounds, including natural vertebrate hormones (17β-estradiol, estrone, testosterone, progesterone), natural invertebrate hormones (20-hydroxyecdysone, juvenile hormone-III), hormone antagonists (flutamide, tamoxifen, hydroxyflutamide), xenoestrogens (17α-ethinylestradiol, 4-octylphenol, bisphenol A), and environmentally relevant compounds (nonylphenol ethoxylate, diethyl phthalate) on larval metamorphosis (nauplius to copepodite) in the calanoid copepod *Acartia tonsa*. The important points of this study were that chemicals differed in their relative effects on survival and larval development. Some of the chemicals tested affected development at concentrations well below those determined to be toxic. Others delayed development only at concentrations close to those that were toxic, so that main effect of the chemical was likely toxicity. Chemicals having very similar effects on toxicity, and having similar octanol/water partition coefficients (K_{ow}) could have very different effects on larval development. For example, flutamide and testosterone have the same K_{ow} and had similar effects on survival of the copepods, yet flutamide was a much stronger inhibitor of development than testosterone.

66. *Amphipods*.—There have been few studies of endocrine disruption in marine amphipods. Brown et al. (1999) found that exposure to 4-nonylphenol reduced growth in the gammaridean amphipod *Corophium volutator*, but that this was probably a general response to the exposure rather than an interaction with molting hormones. They also reported that males exposed to 4-nonylphenol had larger second antennae than those in control treatments and suggested that the compound may have acted on the androgenic gland. Field studies suggested that intersexuality observed in an estuarine amphipod, *Echinogammarus marinus*, might be indicative of endocrine disruption, although a causative link was not established (Ford et al. 2004a, 2004b).

67. *Mysids*.—Mysid crustaceans have been used in regulatory (and other) toxicity testing for more than 20 years. Standard testing protocols have been developed for some species. Despite that little is known about general endocrine functions in mysids (Ingersoll et al. 1999) and there have been few direct links between potential EDCs, beyond certain IGRs, and endocrine disruption in mysids, they have been suggested as providing a useful model of the hormonal control of crustacean molting (Cuzin-Roudy and Saleuddin 1989). McKenney and Celestial (1996) studied the effects of methoprene, which is a JH analog used to control mosquitoes, on *Americamysis bahia* and found that mysids grown at sublethal concentrations were smaller, had a longer time to the production of the first brood, and produced fewer young per female than control animals. They suggested that the effects shown were likely from the interruption of endocrine function by the methoprene.

68. Advances in biological control agents to control insect pests have inspired the synthesis of insect growth regulators (IGR), which find their way into the estuarine environment by either direct or indirect

application. Crustaceans, which along with insects are in the phylum Arthropoda, could also be sensitive to these compounds. The mysid group has been shown to be among the most sensitive members of the estuarine community (McKenney 1982, 1985, 1986, 1996; Nimmo and Hamaker 1982; Nimmo et al. 1981). One study focused on exposure of *Americamysis bahia* to methoprene, a JH analog (McKenney and Celestial 1996). The goal was to determine whether typical application rates shown to control mosquito larvae also cause problems for nontarget organisms. The results showed a significant effect during the mysid life cycle test. Total lethality occurred at 125 µg/L in a 4-day test. Similar concentrations caused significant mortality in the larvae of an estuarine crab and shrimp. Other sublethal endpoints, such as reduced growth (weight), longer time to first brood, and a significant reduction in brood size were also observed. These results suggest that methoprene could interfere with the endogenous endocrine system, which uses hormones that act like JH. Retarded growth rates were also accompanied by bioenergetic disruption, resulting in lower net growth efficiency values. This suggested that increased metabolic demands reduced the amount of assimilated energy available for new tissue production (McKenney 1982, 1985). The delays in mysid first brood production could be the result of slowing sexual maturity and/or embryogenesis. Diminished reproductive success could be the result of inhibited vitellogenesis, modifications in ovarian development, or disruption of successful embryogenesis. In either case, further work is required with the mysid to determine a more conclusive cause-and-effect relationship between potential EDCs and their effects, as observed by test measurement endpoints.

69. Verslycke et al. (2002, 2003a, 2003b, 2004b) studied testosterone metabolism by mysids (*Neomysis integer*) and examined the changes in energy and testosterone metabolism after exposure to potential EDCs. Because these studies focus on potential endpoints, they are presented in more detail in Section 6.4.3.

70. *Decapods*.—Several studies have investigated the effects of EDCs on crustacean life cycles using decapod larvae as the test organisms (e.g., Lee and Oshima 1998; McKenzie et al. 1998; Nates and McKenzie 2000). Several studies (reviewed by McKenzie 1999) have reported effects of the exposure of decapod larvae to JH analogs suggestive of the interruption of endocrine processes, but direct links were not established. More recently, Nates and McKenzie (2000) found that exposure to the pesticide fenoxycarb disrupted lipid metabolism in mud crab larvae and suggested that the compound could be interfering with the endocrine regulation of lipid metabolism. Exposure to fenoxycarb delayed maturation of xanthid crab (*Rhithropanopeus harrisi*) larvae by about 25% compared to controls (Cripe et al. 2003). Some evidence of endocrine disruption in decapods was provided by Snyder and Mulder (2001) who found that exposure of lobster (*Homarus americanus*) larvae to the pesticide heptachlor altered ecdysteroid hormone levels that were linked to delays in molting. Metals such as mercury, cadmium, and zinc have been reported to affect molting and limb regeneration in crabs. Organic compounds such as Aroclor 1242 and sodium pentachlorophenate reportedly had a similar effect—inhibition of limb regeneration—in the grass shrimp, *Palaemonetes pugio*, but had no effect on the molting cycle. This suggests that these chemicals act directly on limb development, but not on the hormonally controlled molting cycle (Fingerman et al. 1998).

71. Echinodermata: Candia Carnevali et al. (2001a, 2001b) studied the feather star *Antedon mediterranea* (Crinoidea) arm regeneration in response to PCB exposure as endocrine disruptor effect. They found that the early phases of regeneration were not different in feather stars exposed to PCBs than in those from control treatments, but noticeable effects occurred later (first at ~7 d, more extensive effects at ~14 d). The primary effect observed in exposed treatments was increased growth rate. Effects, including rearrangement and/or dedifferentiation of some tissues in the regenerating stump, also occurred at cellular level. No abnormal histological effects on feather star arms were observed. The authors concluded that the observed growth and tissue/cellular effects from the exposure PCBs were consistent with pseudoendocrine activities and steroid dysfunction.

3.2.3 Summary

72. Rather than summarize the various impacts, or lack of impacts, of the EDCs that have been studied to date, this section will highlight some of the general findings that are important to consider.

- Chronic assays often reveal impacts by a chemical at doses that are much lower than those eliciting effects during acute exposures. This emphasizes the importance of chronic assays in an EDC evaluation program (e.g., Watts et al. 2001b, 2002).
- Potential EDCs may have effects in invertebrates other than those anticipated by knowledge of their actions in vertebrates (e.g., Hahn et al. 2002, Watts et al. 2001b).
- Pesticides that target certain aspects of insect physiological process may affect nontarget organisms in ways not predicted by the pesticide's desired action (e.g., Hahn et al. 2001, 2002).
- Males and females may react differently to chemicals that are not thought to cause gender-specific responses (e.g., Hahn and Schulz 2002).
- Chemicals that have very similar effects on toxicity, and have similar octanol/water partition coefficients (K_{ow}), could have very different effects on endocrine-controlled processes (e.g., Anderson et al. 2001).
- Some of the endpoint response measured in a study of a potential EDC may also be caused by non-EDC stressors.

73. Finally, and perhaps most importantly, chemical mixtures may show synergistic impacts that are not evident in the actions of the individual components (e.g., Mu and LeBlanc, 2004). In nature, most animals will most certainly be exposed to chemical mixtures rather than to single compounds, thus considering the potential effects of mixtures is important to an EDC evaluation program. However, to be able to predict the possible outcomes of mixtures, at a minimum the individual impacts and mechanisms of action of the component chemicals need to be understood.

3.3 EXTRAPOLATION ISSUES

74. The traditional practice in toxicology of testing the effects of a stressor on individuals from one or a few species has raised two major issues that are of concern to an EDC evaluation program. Simply put, can data collected from tests involving individuals of a species be used to predict potential impacts to other species, or to populations? These two extrapolation issues are discussed in this section.

3.3.1 Extrapolation From Taxon to Taxon

75. It is not practical, or even possible, to conduct toxicity tests on every species in a particular ecosystem (e.g., Hutchinson 2002), therefore, the reasonableness of extrapolating the effects of a stressor (e.g., an EDC) observed for one species to another is an important concern (Segner et al. 2003). There are several approaches to establishing the transferability of the results from testing one species to another. One method is to compare the structural and functional similarity of endocrine systems and/or hormones among taxa. For example, Oberdörster and Cheek (2001) reviewed studies on the ecdysteroid systems of various arthropods, mentioning that the structure of ecdysone in crustaceans is identical to that in insects and that the primary function of the hormone in both groups is to regulate molting. They concluded that information on this system in one arthropod species should be applicable to another. More support for this type of argument is that there is some evidence that peptides are conserved through various evolutionary

lineages from invertebrates to vertebrates (Anctil 2000). This could be interpreted to mean that effects on particular ecdysteroid or peptide systems in one species are likely to be the same as in another species that has the same or very similar system. There are several issues that render this reasoning inadvisable. Lafont (2000) stressed that one of the key general observations concerning invertebrate endocrine systems was that molecules may be structurally related and yet have very different functions in different taxa. Chang et al. (2001) reviewed many studies and showed that the same hormone may have functions that differ among the different life stages of an individual. Finally, even if a hormone can be shown to have the same function in a taxon, the sensitivity of that hormone to disruption by a chemical could vary among taxa. For example, Watts and Pascoe (2000) who clearly demonstrated that *Chironomus tentans* and *C. riparius* differed significantly in their responses to the same chemical stressor. These observations argue for caution in the use of data from a few species to craft generalizations about many (Lafont 2000).

76. Despite this difficulty in extrapolating from one taxon to another, a program to evaluate potential EDCs can be successful if it selects a sensitive taxon that is ecologically relevant (Nimmo and Hamaker 1982). This approach may not offer protection to all species, but should at least offer some degree of protection to critical parts of the ecosystem that have similar exposure probabilities. The program should also develop robust test protocols (Hutchinson 2002) that can be used efficiently by many laboratories and eventually should include representatives from several taxonomic groups (Hutchinson 2002, Oehlmann and Schulte-Oehlmann 2003).

3.3.2 *Extrapolation from Individuals to Populations*

77. The toxicological effects of chemicals, including potential EDCs, as measured during tests of exposure to individuals may not have the same potential to predict possible impacts to populations because the population response may not occur at the same exposure levels as the individual response or may be masked by another stress response. Therefore, the challenge posed to programs charged with assessing the potential risks of EDCs is to establish likelihood that a chemical, or suite of chemicals, will have adverse effects on populations (Crisp et al. 1998, Gleason and Nacci 2001). Thus, the critical need is to be able to detect specific cause-and-effect relationships between a chemical and the responses observed in the field when several factors may cause similar responses (Crisp et al. 1998).

78. Two general approaches to extrapolating from EDC effects on individuals to potential impacts to populations have been identified. One approach uses data gathered during bioassays to attempt direct extrapolation to populations. The other approach uses population modeling techniques to connect effects observed on individuals to potential population impacts.

79. As an example of the first approach, Watts et al. (2002) used survival and sublethal endpoints (growth, development, reproduction) to attempt to predict effects at higher levels of organization. They used a life table or demographic study where they recorded age-specific mortality and fecundity of individuals during life span of a single cohort and used those data to estimate the intrinsic rate of population increase. Watts et al. (2001b) had done an earlier acute study that showed no effect of 17 α -ethinylestradiol on the freshwater amphipod *Gammarus pulex*. A subsequent chronic exposure resulted in changes in maturation rate and sex ratio that translated to measurable effects at the population level (at least in the laboratory) (Watts et al. 2002). Watts et al. concluded that chronic tests provide a more integrated approach to predicting population responses.

80. McTavish et al. (1998) presented a general modeling approach to the evaluation of potential EDCs. McTavish et al. mentioned that many EDCs may persist in natural environments for some time because they are lipophilic and have fairly slow decay rates. These persistent EDCs may change mortality, reproductive, and life-stage transition rates, might become noticeable only after some delay, could affect different life stages present, and may affect offspring of exposed individuals. This introduces a level of

complexity into EDC evaluations that may interfere with efforts to use typical toxicity test data to predict population-level effects. Mathematical models are useful tools to investigate these complex dose-response relationships (McTavish et al. 1998, Gleason and Nacci 2001). McTavish et al. built a general model that enabled mortality, reproductive, and life-stage transition rates to be evaluated alone or in combination. The model also allowed delayed responses and transgenerational impacts to be analyzed. The model evaluated the effects of two dosing scenarios; a single pulse of a chemical into a system and its subsequent decay, and continuous dose of a chemical. McTavish et al. argued that the effective dose response should be based on chemical concentrations in the test animals rather than those in the water because traditional dose-response assessments based on water concentrations may not detect all potential risk to natural systems. This approach probably is not very practical for many invertebrates, especially small arthropods because the amount of tissue required for the analyses would significantly increase the number of animals required for the test. The authors concluded that models are very useful for examining the changes from chemical exposure to the effective dose, and from that dose to populations. Models can synthesize laboratory toxicity test data and use them to extrapolate to a composite picture under a variety of dosing regimes and also can be used to assess the relative importance of various bioassay endpoints. Combinations of stressors can act synergistically to cause significant effects on individuals or populations even though the individual compounds did not when tested separately (Arnold et al. 1996, Mu and LeBlanc 2004). These synergistic effects can be evaluated by the model that McTavish et al. used.

81. Kuhn et al. (2000, 2001) applied the general principals proposed by McTavish et al. in a general evaluation of the ecological relevance of mysid toxicity tests using *Americamysis bahia*. Kuhn et al. (2000) used a model to predict the concentration of a contaminant that would result in no population growth and compared that concentration to standard toxicity test data. They also evaluated which test endpoint, survival or reproduction, was better at predicting population responses. Later, Kuhn et al. (2001) evaluated the ability of their age-classified projection matrix model to predict the population response of *A. bahia* over more than three generations maintained in the laboratory. Data incorporated into the model were gathered from daily records of survival and reproduction, which begins at about Day 17, during toxicity testing with several types of chemicals (e.g., metals, organic compounds). The analyses showed that the population growth rates were dependent on concentration for most of the chemicals tested. Kuhn et al.'s comparison of endpoints showed that the LC_{50} estimated from a 96-h exposure and the chronic life-cycle test endpoints were highly correlated to population changes ($r = 0.96$ and 0.93 , respectively). However, linear regression analysis showed that the life-cycle test was better than the LC_{50} at predicting population-level effects. Within the chronic assay, reproduction was more strongly correlated than survival to population growth rate ($r = 0.98$ and 0.79 , respectively). The model was verified (Kuhn et al. 2001) by using the results of a 28-d life-cycle test conducted with *para*-nonylphenol as the toxicant, to generate data that allowed the model predictions of mysid abundances at the end of the multigenerational (55-d) assay. The predicted abundances then were compared to actual abundances from the assay. The results showed that the model could predict population effects "reasonably well" (Kuhn et al. 2001) in the laboratory. The model-predicted concentration ($16 \mu\text{g/L}$) at which no population growth would occur did not differ significantly from that derived from the multigenerational test data ($19 \mu\text{g/L}$). The data from the 28-d test also were used to calculate a chronic exposure value ($12 \mu\text{g/L}$), which was compared to the 55-d "no-growth" concentration. The lower value was selected as the conservative concentration at which the population should be protected from exposure to *para*-nonylphenol.

82. While either or both approaches may be appropriate, the important consideration is that if population-level impacts are ignored, the possibility that the direct effects of EDCs on ecosystems will be underestimated increases (McTavish et al. 1998).

4.0 AQUATIC ARTHROPODS IN THE EVALUATION OF POSSIBLE ENDOCRINE DISRUPTION

83. Many anthropogenic pollutants eventually end up in the world's oceans, carried there through riverine and estuarine pipelines (Nimmo and Hamaker 1982). Since the mid-1990s, there is an increased awareness that many sewage constituents or chemicals associated with industrial production that enter the environment can disrupt endocrine systems, and that these compounds will likely affect marine organisms (Depledge and Billingham 1999; Oberdörster and Cheek 2001). Although early concern over EDCs focused on vertebrates, attention recently has broadened to include invertebrates because they are ecologically important and have distinctive endocrine systems that differ from those of vertebrates. Estuaries, which are intrinsically and commercially important ecosystems, are among the earliest recipients of EDCs. Among the many estuarine organisms that could be adversely affected by these compounds, crustaceans are good candidates for study of potential impacts. Some of these species are discussed in this section. Attention here is focused on aquatic arthropods (insects and crustaceans) because they are often among the most abundant organisms in freshwater and estuarine systems, form vital links in estuarine food webs, and are known to be susceptible to the effects of EDCs.

84. Hutchinson et al. (2000) offered a counter argument to the inclusion by the EDSTAC and EDSP of an arthropod, specifically a daphnid or mysid, in the endocrine screening program at this time. The principal objection applicable to both taxa was the general data gap in the basic endocrinology of mysids and daphnids. They argued that the assertion by EDSTAC that vertebrate estrogen or androgen disruptors could interfere with ecdysteroid activity was only speculation and that any effects on reproduction or development by known vertebrate EDCs have not been directly linked to ecdysteroid or juvenile hormone activity. Hutchinson et al. also expressed concern that the gender identification of mysids in the laboratory was technically difficult and that the test results might be confounded by cannibalism of offspring by parents. These three concerns can be addressed. The understanding of invertebrate endocrine systems is improving rapidly (Section 3.0) and the links between hormones and processes in crustaceans are being established. For example, in companion studies, Mu and LeBlanc (2002a, 2002b) showed that ecdysteroids regulate the embryonic development of daphnid and that this development could be interrupted by testosterone and fenarimol, a fungicide used in agriculture. Testosterone did not display ecdysteroidal activity, but interfered with ecdysteroids by outcompeting them to occupy the ecdysteroid receptor (Mu and LeBlanc 2002a). Fenarimol acted by reducing body levels of ecdysone in individual daphnids (Mu and LeBlanc 2002b). Vandenberg et al. (2003) found that the synthetic estrogen 17 α -ethinylestradiol altered sexual development in individuals of the amphipod *Hyaella azteca*. Males did not develop secondary sex features, reproductive system morphology was altered, and sex ratios in exposed groups favored females. Also, recent studies of crustaceans have identified that methyl farnesoate has a role in regulating larval development in lobsters (summarized by Chang et al. 2001) and stimulates gonad development in a freshwater prawn (Nagaraju et al. 2003). As Verslycke et al. (2003a) stated, the discussion over whether or not an observed effect is directly related to endocrine disruption or not may be appropriate, but neither that discussion, nor the cause of the effect, is of concern to the affected organism. Determining sexual maturity may be difficult, although Khan et al. (1992) used maturity, based on the development of gonad tissue, as a sensitive endpoint and thought that such determinations could be done by unsupervised laboratory staff. Females and males have anatomical differences that can be used to indicate maturity. In females, the appearance of the oostegites that form the marsupium are one indication of maturity. Maturity is more difficult to recognize in males, but one pair of abdominal pleopods is

modified to assist in copulation. Molt staging could be used to alert the laboratory staff of the potential appearance of mature individuals because *A. bahia* reaches maturity at the fourth molt (Touart 1982). The susceptibility of young to being eaten by adult mysids is also well-known (e.g., Johnston and Ritz 2001, Quirt and Lasenby 2002). Premolt newborns are particularly susceptible. Testing protocols call for newly released young to be removed from the tanks as soon as possible and placed into separate rearing chambers (or new test chambers if they are to be exposed to a potential toxicant. Mysids are also typically fed liberal quantities of food (*Artemia* nauplii) to lessen the likelihood that adults will eat the young. Clearly, considerable evidence exists that counter the concerns raised by Hutchinson et al. (2000).

4.1 FRESHWATER ARTHROPOD SPECIES

85. Although many freshwater invertebrate species might be chosen for use in EDC evaluations, the focus in this section is on three groups of arthropods. The groups selected are ecologically important, have had well-developed testing protocols prepared and evaluated, and have been widely used in general toxicological testing in addition to endocrine disruptor screenings.

4.1.1 Midge Larvae (*Chironomus tentans* and *C. riparius*)

86. Natural History: *Chironomus* spp. are true midges (Order Diptera) that have a widespread worldwide distribution (Pennak 1989). Adult midges are small, nonbiting, and swarm near bodies of water at night. Larval stages are aquatic, inhabiting fine tubes in the upper layers of the sediment. Midge larvae feed on algae, higher plants, and organic detritus (Pennak 1978). Chironomid larvae are ecologically relevant for toxicity testing because of their widespread distribution, numerical abundance, and importance as prey for juvenile and adult fish. Two species of *Chironomus* are typically used in toxicity testing; *C. tentans* in North America and *C. riparius* in Europe. *Chironomus tentans* eggs hatch about 2 d to 6 d after laying and begin the first of four aquatic instar larval stages (summarized in Benoit et al. 1997). The larval development lasts about 23 d and is followed by a short (~1–2 d) pupal stage. Males emerge about five days before females (protandry). Adults are short-lived, about 7 d. The aquatic development period for *C. riparius* is shorter as adults emerge within about 15–17 d after beginning the first instar stage (Watts and Pascoe 1996, Watts et al. 2001a) and live about 4 d (e.g., Ristola et al. 2001). This was confirmed in a direct comparison in natural and artificial control sediments between the two species (Watts and Pascoe 2000).

87. Availability, Culture, and Handling: Both species of *Chironomus* are readily available from commercial sources and do not need to be collected from the field. They are easily maintained in laboratory culture. Detailed culture requirements for *C. tentans* have been described by EPA (2000). Larval are held at about 23°C in glass aquaria (e.g., ~19 L volume containing ~8 L water) with finely shredded paper toweling or silica sand used as substrate. About 600 eggs per 8 L should be stocked into each aquarium. High densities of larvae increase development time, delaying adult emergence. Larval are fed a slurry prepared of commercial fish food flakes at a concentration of ~0.04 mg dry food/mL culture water. Adult emergence commences in about three weeks after egg hatching (23°C). Chronic tests must be initiated with <24-h old larvae, therefore, laboratories must maintain cultures to ensure larvae of the correct age are available for testing.

88. McCahon and Pascoe (1988) presented information on culturing *C. riparius*. Conditions for culture are somewhat similar to those for *C. tentans*. The substrate is based on homogenized cellulose paper (filter paper). Cultures are maintained at 18°C to 20°C and are fed a ration of commercial fish food flakes daily.

89. Strengths and Weaknesses: The principal strengths of using *Chironomus* in toxicological testing are the high ecological relevance, ready availability, and ease of culturing of the two primary species.

Formal test procedures for both species are well-developed and the species have been used in a variety of laboratory (e.g., Benoit et al. 1997, Environment Canada 1997a, EPA 2000) and *in situ* (Castro et al. 2003) testing situations. At least one other species, *C. prasinus*, has been used in some EDC testing (Sánchez and Tarazona 2002). The life cycle of chironomids is relatively short and amenable to multigenerational testing. The life cycle of *C. tentans* is about 33–38 days (Benoit et al. 1997); that of *C. riparius* is likely shorter (Watts & Pascoe 1996, Watts et al. 2001a). Recent studies have included several evaluations of potential endocrine disrupting chemicals on *C. riparius* (Hahn et al. 2001, Watts et al. 2001a, Hahn et al. 2002, Hahn and Schulz 2002,) and *C. tentans* (Kahl et al. 1997). One possible weakness is the apparent lack of response, or conflicting responses, of life cycle endpoints in *C. riparius* to exposure to two known xenoestrogens, bisphenol-A and 17 α -ethinylestradiol, or in *C. tentans* exposed to the surfactant 4-nonylphenol (Kahl et al. 1997). Chironomid larvae are sediment dwellers, therefore the testing protocol must provide the test animals with a suitable substrate. Watts and Pascoe (2000) showed the type of sediment (artificial or natural) was an important contributor to the differences in response that they observed. Concentrations of contaminants were greater in the porewaters of the artificial sediment than in those of the natural sediment.

90. Another important issue that must be considered when using either species of *Chironomus* is that the two were shown to have different sensitivities to the same toxicant. Watts and Pascoe (2000) found that *C. tentans* was more sensitive than *C. riparius* to the same toxicants evaluated under the same test conditions. They also determined the *C. tentans* was less physically robust than *C. riparius*, which led to increased variability in the data for *C. tentans*, especially emergence data (there was poor emergence in control animals). The response criteria were affected by type of sediment, the choice of species, the particular toxicant, and the duration of the experiment.

4.1.2 Amphipods (*Hyalella*, *Gammarus*)

Hyalella azteca Saussure 1857

91. Natural History: Historically, *Hyalella azteca* has been considered a very common inhabitant of permanent fresh water systems throughout North, Central, and northern South America (Bousfield 1973, Pennak 1989, Gonzalez and Watling 2002). As discussed in the “Strengths and Weaknesses” section below, the species’ actual distribution is probably much more restricted. *Hyalella* inhabits a variety of unpolluted springs, streams, brooks, pools, ponds, and lakes (Pennak 1989), but also may occur in large coastal rivers and into tidal fresh waters (Bousfield 1973). These epibenthic burrowing amphipods may occur in large numbers; they are detritivorous, feeding primarily on algae and bacteria. *Hyalella* produces about 18 eggs per brood although the number can vary with female age and size (Pennak 1989). At warmer temperatures (24°C–28°C), eggs hatch in about 5–10 d and females can produce about 15 broods every 150 days (Pennak 1989, EPA 2000).

92. Availability, Culture, and Handling: *Hyalella azteca* has been used in toxicity testing for many years and is widely available from many commercial suppliers. EPA (2000) provides a detailed description of culture requirements. The amphipods can be cultured in containers of various sizes (e.g., 2L–80L) at a water temperature of about 23°C and a light:dark photoperiod of 16:8 h. The water should be renewed periodically and monitored for various parameters such as hardness, alkalinity, pH, and ammonia. Amphipods can be fed commercial fish food flakes, or a combination of yeast-Cerophyl®-trout chow (YCT, prepared according to the recipe in EPA 2000 or purchased commercially) and the alga *Selenastrum capricornutum*.

93. Strengths and Weaknesses: *Hyalella azteca* is an ecologically relevant testing organism. It has been used in many types of toxicity testing and several protocols, including life cycle testing, to guide testing have been written for it (e.g., Environment Canada 1997b, EPA 2000). This amphipod is readily

available and relatively easy to culture. However, endocrine disruption evaluations, and any other toxicity evaluations, with this species are hampered by recent revelations that the widespread “species” *Hyalella azteca* is actually a mixture of many species. Several studies, beginning in the late 1990s, have investigated various populations of *H. azteca* from around North America. Duan et al. (1997) studied six lab “populations” of *H. azteca*; one from Burlington, Ontario, one from Conesus Lake NY; and four originally derived from the Corvallis, Oregon population [the Nebeker strain that was originally collected in 1982 (Duan et al. 2000b)], and another species in the genus, *H. montezuma*. They found that the four populations derived from Oregon were relatively closely related and distinct from the other two and from *H. montezuma*. Duan et al. concluded that the populations represented four species (including *H. montezuma*). McPeck and Wellborn (1998) found allelic and genotypic differences between large-bodied and small bodied ecotypes of *H. azteca* in southeastern Michigan. They determined that the two ecotypes did not interbreed and that it was likely that they represented different species. Duan et al. 2000a studied 12 populations, all presumably *H. azteca*, from geographically isolated parts of N. America. They used electrophoresis to identify nine groups. The large genetic differences that were identified led them to conclude that all nine probably represent different species. Witt and Hebert (2000) studied populations from wide geographical area in NA, most of which were from around the Great Lakes (24 lakes and ponds in Ontario and Wisconsin), but also included the Yukon and New Brunswick. They found low levels of gene flow, reduction in genetic variability, low heterozygosity, unique alleles, and strong genetic differentiation and divergence among populations, concluding that there were at least seven mitochondrial lineages. Gonzalez and Watling (2002) conducted a morphological study of type specimen of *H. azteca* and compared it to specimens from Maine, Texas, Mississippi, Michigan, Oklahoma, and Hawaii concluding that *H. azteca* is a species complex. Thus, the taxon currently know as “*Hyalella azteca*” is probably comprised of at least a dozen species. Gonzalez and Watling (2002) recommend that any further toxicological studies confirm identification of species. If animals are obtained from a commercial supplier, the original source of that supplier’s culture should be identified.

94. The potential ramifications of using different genetic species on toxicity testing was underscored by Duan et al. (2000b, 2000c) who tested the effects of various metals, pH, and fluoranthene on different genotypes within a population of *H. azteca* derived from the Nebeker strain. They discovered that some genotypes were more susceptible to the stressors than others, and that the susceptibilities differed across genotypes. This differential susceptibility implies that a population that has a higher frequency of a particular genotype that happens to be resistant to a particular stressor will show better survival than a population that has a high frequency of a genotype that is susceptible to the stressor. That is, different genetic populations may provide different responses to the same stressor.

Gammarus pulex

95. Natural History: *Gammarus pulex*, an amphipod that is also known as scud, is widespread throughout Europe, where it is often the most abundant invertebrate in streams. *G. pulex* is a primary prey item of fish and used in biotic indices to describe pollution in rivers (summarized by McCahon and Pascoe 1988). The reproductive biology of *G. pulex* involves precopulatory guarding of the female by a male (e.g., Watts et al. 2001b). The male clasps onto a female and swims with her until she molts to assuring sexual contact during the time at which mating can occur. This particular behavior has been developed into a bioassay procedure (Poulton and Pascoe 1990, Pascoe et al. 1994). It is likely that a chemical signal facilitates this behavior, although evidence for pheromonal control of mating in this species is contradictory (summarized by Watts et al. 2001b). A similar assay was developed for the estuarine gammarid, *Gammarus duebeni* (Lawrence and Poulter 1996).

96. Availability, Culture, and Handling: *G. pulex* is available from at least one commercial supplier, but it is not clear if those animals are cultured by the supplier or field collected. Most studies report using field-collected animals and hold them for a short time before testing. McCahon and Pascoe (1988)

developed culture methods for *G. pulex*. Gravid females and pre-copula pairs are removed from the collection and placed in breeding chambers (1-L plastic jars with an open, mesh-covered bottom to allow juveniles to escape); these chambers are placed into larger (2–8 L) tanks that are provided with flow-through dechlorinated water (the chambers can be maintained static, if water is renewed periodically). Adults are fed a variety of conditioned tree leaves (e.g., horse chestnut, elm, sycamore, or oak) that are collected during fall, air dried, and stored until just before required for feeding. The leaves are prepared by being placed in organically enriched dechlorinated water to promote the bacterial and fungal growth that provide an important food for *G. pulex*. About 200 gravid females in a chamber will provide enough juveniles (500–1000) to stock the rearing tank. Juveniles are fed conditioned leaves, which is supplemented with adult feces from the adult containers. Timing can be arranged to provide different age ranges for testing. Culture temperatures were not provided, but at 13°C about 70% of juveniles reach maturity in 130 days, after completing 10 molts.

97. **Strengths and Weaknesses:** *Gammarus pulex* provides a test organism that has considerable ecological relevance in European freshwaters. Because organisms are usually field collected, the potential for population-related differences in pollutant sensitivities is high and should be assessed before comparing the results of tests performed on different populations. However, collected animals are relatively easily maintained in culture systems once established and can provide a suitable number of test organisms for some time. *G. pulex* has been used in many types of bioassay procedures, including behavioral (Poulton and Pascoe 1990, Pascoe et al. 1994) and *in situ* (Matthiessen et al. 1995) testing. However, formal chronic test protocols have not been developed (Segner et al. 2003).

4.1.3 *Daphnids (Ceriodaphnia dubia, Daphnia magna, D. pulex)*

98. **Natural History:** These three species of daphnids are widespread, occurring in freshwater ponds or lakes throughout much of the world (Weber 1993). *C. dubia* is a small species commonly found among plants in the littoral zone of lakes and ponds. *D. magna* lives primarily in lakes where the hardness exceeds 150 mg/L CaCO₃, whereas *D. pulex* lives in clean, well-oxygenated ponds (Weber 1993). Much of the year, when environmental conditions are favorable, populations of these species consist primarily of females; males are produced when the environment begins to become unfavorable. Male *C. dubia* usually appear in autumn and those of *D. magna* in spring or autumn. *C. dubia* and *D. magna* reproduce via cyclic parthenogenesis with the male contributing genetic material only during certain times of the year. *D. pulex* may reproduce by either a cyclic or obligatory parthenogenesis. Daphnids produce ephippia, encased embryos that are resistant to harsh environmental conditions, during periods of sexual reproduction.

99. The life span of these species is highly variable, depending principally on environmental conditions. At 25°C, *C. dubia* and *D. magna* live about 30 days and 40 days, respectively. *D. magna* lives about 56 days at 20°C, whereas *C. dubia* and *D. pulex* live about 50 days at that temperature (Weber 1993). These daphnids produce about 10 eggs per clutch, which hatch into juveniles after about 38 h; the maximum egg production occurs at temperatures from 18°C to 25°C.

100. **Availability, Culture, and Handling:** All three species are readily available from commercial sources and are easily maintained in culture. Only 20–30 individuals are required to start a culture within a testing laboratory. Cultures should be maintained for at least two generations under the same conditions and feeding regimes that will be used during testing. Cultures can be maintained in natural waters, but Weber (1993) recommends “synthetic” water because it produces reliable results. Culture temperature is optimal at about 20°C. *D. magna* requires relatively hard water (160–180 mg/L CaCO₃), whereas *D. pulex* requires softer water (80–90 mg/L CaCO₃). Dissolved oxygen concentrations should be held above 5 mg/L. Cultures should be maintained carefully. The culture medium should be replaced three times per week and the cultures should be thinned when the populations exceed 200 individuals per 3 mL of culture water. Daphnids should be fed a YCT mixture and the alga *Selenastrum capricornutum*.

101. **Strengths and Weaknesses:** Daphnids offer many advantages to an EDC evaluation testing program. They are ecologically relevant, widely distributed, and are readily available via commercial suppliers. Cultures are easy to start and maintain. Daphnids are usually tested at intermediate temperatures (20°C). Acute and chronic protocols for testing daphnids are available and daphnids are actively used in many testing scenarios. Daphnids have been actively used in the evaluations of potential endocrine disrupting chemicals (Section 3.2). The primary weakness with using daphnids in endocrine disruption evaluations is that the asexually reproducing stage is typically the one used in testing because the presence of sexual reproduction is taken as an indication that culture conditions are less than ideal or that testing methods are deficient (Olmstead and LeBlanc 2000). Parthenogenic reproduction results in lower genetic variability within the test population than would occur with a sexually reproducing taxon (Lagadic and Caquet 1998). This reduced variability would reduce the potential variation in response to stressors. However, the emphasis on only testing the parthenogenic phase for endocrine disruptive effects ignores the sexual reproduction phase, which is vital to daphnid population viability (Lagadic and Caquet 1998, Olmstead and LeBlanc 2000). Therefore, EDC impacts to daphnid populations may be severely underestimated. The species discussed above are temperate species and may not be feasible for use in tropical areas. Buratini et al. (2004) evaluated a tropical species, *Daphnia similis*, and found it to be similar in sensitivity to the two temperate *Daphnia* species. *D. similis* is easily cultured and may be more appropriate for use in evaluating potential EDCs in tropical waters that have low hardness.

4.2 ESTUARINE AND MARINE ARTHROPOD SPECIES

102. Although insects, crustaceans, and pycnogonids (sea spiders) are the primary arthropod taxa that occur in estuarine and marine communities, crustaceans are certainly the more abundant and, therefore, the more ecologically important group. Many insect species occur in these waters, including the larvae of chironomid midges, but they are relatively poorly studied. Little is known about pycnogonid biology and endocrinology. Thus, the focus in this section is directed towards the estuarine and marine crustacean species most likely to be amenable to EDC evaluations.

4.2.1 Copepods (*Acartia*, *Tisbe*, *Nitocra*, *Tigriopus*)

103. **Natural History:** Copepods comprise one of the most important groups of marine invertebrates in estuarine and marine systems whether planktonic or benthic. They form a vital link in water column food webs, feeding on smaller plankton and being consumed by larger predators, such as fish. In benthic systems they are often important grazers on microalgae, yet provide forage for young life stages of many fish. The life history of copepods involves morphologically distinct stages (nauplius and copepodite stages). Therefore, it is possible to test the effects of chemicals, including those that are potential endocrine disruptors, on the metamorphosis from nauplius to copepodite stages that might detect interference with processes regulated by ecdysteroids (Anderson et al. 2001).

104. **Availability, Culture, and Handling:** Some species of copepods, *Acartia tonsa* for example, may be available commercially. However, the species typically used in toxicity testing are easy to maintain in cultures so that sufficient animals for testing are always available. Some species have been maintained for at least 20 years (Anderson et al. 2001). Culture temperatures depend on the species involved. *A. tonsa* and *Tisbe battagliai* are cultured at 20°C (Anderson et al. 2001, Hutchinson et al. 1999a, 1999b), *Nitocra spinipes* at 22°C (Breitholtz et al. 2003), and *Tigriopus japonicus* at 25°C (Marcial et al. 2003). Copepods are fed the unicellular algae, such as *Rhodomonas*, *Nanochloropsis*, and *Isochrysis*.

105. **Strengths and Weaknesses:** Copepods are ecologically relevant animals for use in the evaluation of EDCs. They are readily available and easy to maintain and handle. The test conducted by Anderson et al. was an acute test with a very simple and easy to measure endpoint, the proportion of larvae that metamorphose from nauplius to copepodite. *A. tonsa* was more sensitive than *Daphnia magna* to

interruption of growth and molting. Chronic (full life-cycle) tests may be completed within a relatively short time, from 21 days to 25 days. Although formal chronic protocols have not been developed many copepod species have been used in toxicity testing, including the evaluation of potential EDCs (Section 3.2) and the procedures used for those experiments could be adapted for standardized testing worldwide.

4.2.2 Amphipods

106. Several species of benthic estuarine and marine amphipods are used regularly in regulatory toxicity testing, including *Ampelisca abdita*, *Eohaustorius estuaries*, *Rhepoxynius abronius*, and *Leptocheirus plumulosus*. Test guidelines have been established for these species (e.g., EPA/USACE 1998, EPA 2001). Several other species have been used in studies of the toxic effects of various contaminants. Among these are *Corophium volutator*, *Gammarus* spp., *Microdeutopus gryllotalpa*, *Echinogammarus marinus*, and *Grandidierella japonica*. Formal protocols for these latter species have not been developed. Among these amphipods, *L. plumulosus* probably is the best suited for adaptation to an EDC testing program and, therefore, is discussed in more detail below.

Leptocheirus plumulosus:

107. Natural History: *Leptocheirus plumulosus* is a common infaunal amphipod occurring in estuaries along the east coast of the U.S. from northern Florida to Massachusetts (Bousfield 1973). *L. plumulosus* lives in simple burrows it digs into the upper layers of the sediment. It is a detrital feeder, extracting organic material that is suspended in the water and from ingested sediment. *L. plumulosus* tolerates a wide range of salinities, from 1‰ to 35‰ (EPA 2001). Its generation time is short, requiring about 24 days at 23°C (EPA 2001). Another species in the genus, *L. pinguis*, is known to use parental care, sheltering juveniles in its burrows (Thiel 1997). *L. plumulosus* has become increasingly selected as the test species in contaminant sediment evaluations. A chronic survival, growth, and reproduction testing protocol for the species was recently developed (EPA 2001).

108. Availability, Culture, and Handling: *L. plumulosus* is readily available from several commercial suppliers who maintain large cultures of the species. Laboratories can obtain amphipods having a wide variety of ages. Laboratories conducting the chronic testing protocol need to culture the amphipods in-house because the protocol requires that the test be initiated with neonates (<48-h old). Cultures are easy to start and are maintained at temperatures of 20°C to 25°C and at the salinity that will be used in testing, which is usually 20‰ (EPA 2001). Cultures are fed finely milled fish food flakes and the culture water is renewed three times per week. Cultures may have to be thinned because these amphipods are very prolific. Cultures require about 6 weeks to mature sufficiently to provide enough neonates for testing.

109. Strengths/Weaknesses: The principal strengths of *L. plumulosus* for EDC evaluations are its ready availability, ease of culture, and sensitivity to contaminants. The species is becoming widely used in toxicity testing programs. A 28-day chronic testing protocol, which could be adapted to EDC testing programs, has been developed for the species (Emery et al. 1997, EPA 2001). Some aspects of the species' reproductive endocrinology have been studied (Volz et al. 2002, Block et al. 2003) and the impacts of some chemicals known to affect endocrine systems have been tested on the species (Lussier et al. 2000, Zulkosky et al. 2002). Spencer and McGee (2001) used data collected from field populations of *L. plumulosus* to develop a stage-structured population model that allows extrapolation from laboratory toxicity tests to population impacts. McGee and Spencer (2001) tested the model by using laboratory testing data to project the effects of sediment toxicity on the population growth rate. They found that the population projections were very similar to amphipod abundances observed at the site from which the sediment was collected. *L. plumulosus* abundance in the area from which the sediment was collected. *L. plumulosus* occurs only along the east coast of the U.S. and, therefore, serves only as a surrogate species for estimating the impacts of EDCs on faunas from other regions.

4.2.3 Decapods (shrimp, crabs, crayfish, lobsters)

110. Decapods probably have been the subjects of more endocrine function and possible EDC effects-related testing than any other crustacean group (Fingerman et al. 1998, Ingersoll et al. 1999). Decapods are particularly appealing to EDC studies because much is known about their endocrine systems and their free-swimming larvae are likely to be susceptible to JH analogs and other Insect Growth Regulators (IGRs) produced to control insects (McKenney 1999).

111. Two species, the mud crab (*Rhithropanopeus harrisi*) and the grass shrimp (*Palaemonetes pugio*), have been the primary subjects for many of the endocrine studies and have been advocated as being potentially useful in studies of EDCs (McKenney 1999). Several studies have investigated the effects of EDCs on crustacean life cycles using decapod larvae as the test organisms (Section 3.2). Most of the recent work on the impacts of various EDCs on decapods has centered on the grass shrimp *Palaemonetes pugio* and the mud crab *Rhithropanopeus harrisi*, which, therefore, are the focus of this section.

112. Natural History: The grass shrimp, *Palaemonetes pugio*, is widespread in estuaries from Nova Scotia to Texas (Williams 1984) and has been used considerably in bioassays. Grass shrimp feed on bacteria-laden benthic detritus and are an extremely important component of energetic flux pathways in estuaries (studies summarized in Williams 1984). Gravid females are available from late spring to late summer. Grass shrimp usually mature about six months after hatching, although maturation may occur quicker where water temperatures are warm (Williams 1984, Volz et al. 2002). The mud crab, *Rhithropanopeus harrisi*, occurs in estuaries from the southwestern Gulf of St. Lawrence, Canada to Veracruz, Mexico and has been introduced into parts of Europe and the U.S. west coast (Williams 1984). Mud crabs tolerate a wide range of salinities and temperatures. Mud crabs are omnivorous, feeding on a variety of aquatic plants, detritus, and other estuarine animals. Gravid females typically can be collected during the summer months. Mud crab larvae, rather adults, are typically used in toxicity studies.

113. Availability, Culture, and Handling: Grass shrimp are available from commercial suppliers and can be maintained in culture by testing laboratories. Mud crabs must be collected from the field before testing. Grass shrimp can be held at a wide range of conditions with temperatures ranging from 18°C to 25°C and salinities ranging from 20‰ to 28‰. Oviparous females are held in the laboratory until the eggs hatch. Larvae, which are used in toxicity testing can be cultured at the above conditions in the laboratory. Many studies involving the mud crab use the larval stages rather than adults. Oviparous females can be collected and held in cultures until the larvae, which are relatively easy to raise, are released. Larvae can be cultured at a water temperature of about 25°C and a salinity of about 20‰. Larvae of both species are fed brine shrimp (*Artemia*) nauplii daily.

114. Strengths/Weaknesses: The principal strength that both species offer an EDC evaluation program is that the knowledge of decapod endocrine systems is relatively well-developed. Both are common animals in estuaries of the U.S. Gulf and East Coasts. One species, the grass shrimp, is commercially available. Both are amenable to individual, focused studies of the specific impacts of EDCs on various endocrine system components and, thus, are important species to study. However, neither is advantageous for use in an overall EDC screening program, primarily because the long generation time precludes relatively rapid multigenerational studies. Additionally, the mud crab must be collected from the field and would be difficult for many testing laboratories to obtain.

4.3 BRAKISH, FRESHWATER AND MARINE ARTHROPOD SPECIES

115. Several mysid species are considered below for their potential utility in EDC testing. For each, a discussion of natural history, availability, and culture and handling is offered, along with a summary paragraph on strengths and weaknesses of the species as test organism. Ingersoll et al. (1999) included

several additional characteristics that are important considerations in the selection of test species. Some of the features, for example mode of reproduction and knowledge of endocrinology, do not allow for discrimination among candidate mysid species. [For several of the species in the section below, the name of the researcher who described them is cited. If the species was originally described in a genus other than the one listed, the researcher's name is enclosed within parentheses.

4.3.1 *Americamysis bahia* (Molenock, 1969)

116. *Americamysis bahia* is a small mysid crustacean that occurs in coastal estuaries and embayments ranging from the Gulf of Mexico to Narragansett, Rhode Island (Price et al. 1994).

117. Natural History: *A. bahia* typically occurs in areas where the salinity is >15‰, but is more abundant in higher-salinity habitats. Molenock (1969) originally described the species as *Mysidopsis bahia*, but Price et al. (1994) transferred it to a new genus, *Americamysis*, during their taxonomic revision. The natural history of the species is well known; the following description is based on Weber (1993). Adults of *A. bahia* may reach almost 10 mm in total length, and females attain a larger size than males. Individuals become sexually mature in about 12 to 20 days, and the genders can be distinguished when the animals reach about 4 mm in total length, at which time the brood pouch typically has started development. At about 12 days, the female's ovaries begin to contain eggs, and the brood pouch is fully formed at about 15 days. The developing young are carried in the brood pouch for an additional 2 to 5 days, resulting in a life cycle of about 17 to 20 days. New broods may be produced about 4 to 7 days. Females produce an average of about 11 eggs per brood, and the number is directly related to female body length.

118. Availability, Culture, and Handling: *A. bahia* is cultured commercially by many laboratories located throughout the United States and therefore is readily available to testing laboratories. With only a few days' notice, commercial suppliers can ship <24-h-old mysids via overnight service, allowing testing laboratories time to acclimate the animals to test conditions. However, testing laboratories can also easily culture the species. The mysids can be raised in 80-L to 200-L aquaria provided with continuous flow-through or recirculating systems. The water temperature and salinity within the culture tanks are typically 24°C to 26°C and 20‰ to 30‰, respectively. The cultures are fed newly hatched brine shrimp (*Artemia* sp.). Several sources provide information on culturing this species (e.g., Lussier et al. 1988). Although a small species, individuals of *A. bahia* are relatively easily handled.

119. Strengths and Weaknesses: As a candidate test species, *A. bahia* has many strengths and few weaknesses. Its primary advantages include its widespread availability and ease of culture. Because animals can be obtained from commercial cultures, the likelihood of misidentifying the species is very low. Its relatively short generation time makes it desirable in life cycle testing. The species is widely used in toxicological testing, and appropriate test conditions are well known. Standardized life cycle test protocols have been developed (EPA 1996; ASTM 1997) and applied (e.g., McKenney 1982, 1985, 1986, 1994; McKenney et al. 1991; McKenney and Celestial 1996) or evaluated (Lussier et al. 1999). Included among the standardized test protocols are many of those necessary to measure EDC-related endpoints; others appear in peer-reviewed publications (Ingersoll et al. 1999). McKenney (1998) synthesized the results of several of his earlier studies on the effects of chronic exposures to various pesticides on *A. bahia* and concluded that life-cycle endpoints (slowed juvenile growth and reduced production of young) were more sensitive than survival. More recently, studies showed that the results from toxicity tests, which included standard and multigenerational tests using *A. bahia*, could be used to extrapolate from laboratory to population effects (Kuhn et al. 2000; 2001). Verslycke et al. (2004a), in their review of the use of mysids for endocrine disruptor evaluations, favored *A. bahia* as a reasonable test species. One criticism of the widespread use of *A. bahia* in toxicological testing is that because it is a warm-temperate or subtropical species, it may not be ecologically relevant to colder-water materials testing.

4.3.2 *Holmesimysis costata* (Holmes, 1900)

120. *Holmesimysis costata*, previously referred to as *Acanthomysis sculpta*, is an ecologically important species that ranges from southern California to British Columbia (Hunt et al. 1997).

121. Natural History: *H. costata* is a dominant member of the plankton community living within the surface canopy of the giant kelp, *Macrocystis pyrifera* (Chapman et al. 1995) and is an important pelagic prey of the California gray whale, *Eschrichtius robustus* (Dunham and Duffus 2002). Adults may reach lengths of about 7 mm to 13 mm (Daly and Holmquist 1986), and females attain a larger size than males (Turpen et al. 1994). Sexual maturity occurs at about 42 days, at which time males, which are recognized by an extended fourth pleopod, can be distinguished from females, recognized by the developing brood pouch (Turpen et al. 1994). Young are released when the females are about 65 to 73 days old. Brood size among laboratory-cultured females averaged about 16 released juveniles per female; in contrast, field-collected females released substantially larger broods that averaged 27 young (Turpen et al. 1994). Brood size was directly related to female size. EPA (2002a) reported that females may produce multiple broods during their 120-day life span. However, Turpen et al. (1994) also reported that all laboratory-reared females died before releasing a second brood of young.

122. Availability, Culture, and Handling: Field-collected animals are available from a few suppliers and are likely to be available year round, but the species is not cultured commercially (Turpen et al. 1994). Brood stock can be collected by sweeping a small-mesh net through the canopy of the giant kelp (EPA 2002a). Field-collected mysids can be cultured in the laboratory, and guidelines for doing so have been established (Chapman et al. 1995, EPA 2002a). However, broodstocks should be rejuvenated periodically by the addition of field-collected animals (Turpen et al. 1994). Culture tanks can range in volume from 4 L to 1000 L and should be provided with aeration and fronds from the giant kelp, *M. pyrifera* (Chapman et al. 1995, EPA 2002a). *H. costata* is typically maintained and tested at temperatures that range from about 13°C to 15°C for animals collected north or south of Point Conception, CA, respectively (EPA 2002a). Adults are fed newly hatched *Artemia*, whereas juveniles are fed *Artemia* supplemented with a small amount of ground fish-food flakes (e.g., Tetramin). The animals are easily handled by using a combination of small-mesh dip nets and pipettes to transfer them from culture tanks to test chambers.

123. Strengths and Weaknesses: The primary strengths of this species are its ecological relevance to northeast Pacific regional testing conditions, its relatively large brood sizes, and its ease of handling and maintenance. *H. costata* occurs in relatively cold waters and may serve as a coldwater alternative to *A. bahia*. The species has been used in several toxicological tests (e.g., Singer et al. 1998) and standardized test protocols for the species have been developed (Chapman et al. 1995, EPA 2002a) and evaluated (Martin et al. 1989, Hunt et al. 1997). Martin et al. (1989) determined that *H. costata* had sufficient sensitivity to evaluate effluent toxicity and that the results were repeatable by different testing laboratories. The principal disadvantages inherent in using *H. costata* are its long generation time (~70 days) and the difficulty in raising multiple broods in the laboratory. The tests required to measure many EDC-related endpoints must be developed (Ingersoll et al. 1999) or are impractical because of the species long generation time and the difficulty in raising multiple broods. Because the original animals are field-collected, they must be identified carefully prior to their use in testing. Also, potential population-related differences in pollutant sensitivities should be assessed before comparing the results of tests performed with mysids from different populations.

4.3.4 *Mysidopsis intii holmquist*, 1957

124. *Mysidopsis intii* is an epibenthic species that occurs in the eastern Pacific from South America to the southern California coast of the United States (Price et al. 1994; Langdon et al. 1996).

125. Natural History: *M. intii* has only recently been reported from the United States (off Los Angeles), but it could be more widespread (Langdon et al. 1996). *M. intii* is a relatively small species that attains body length of about 6 mm to 7 mm. The genders can be distinguished at about 9 to 10 days after hatching. Eggs enter the brood pouch at about 13 days, and juveniles are released at about Day 20 (Langdon et al. 1996).

126. Availability, Culture, and Handling: Animals to be used in testing must be obtained from field collections, because there is no commercial culture of the species. Individuals of *M. intii* are collected by using an epibenthic sled, and the wild-caught animals are then used to establish breeding stocks in the laboratory. The species is easily cultured in the laboratory in 40-L to 90-L tanks continuously supplied with flowing, filtered seawater. Langdon et al. (1996) determined that the optimal temperature for high juvenile production is 20°C. To ensure high reproductive output, adults should be fed recently hatched *Artemia* and adult copepods, *Tigriopus californicus* (Kreeger et al. 1991; Langdon et al. 1996). The *Artemia* diet can be enriched with fatty acid supplements, and the mysid cultures provided with fatty acid boosters (UCSC 1998). Separation of juvenile mysids from the adult cultures is easily accomplished by using light to attract them through a 1-mm-mesh divider into an isolation chamber (Langdon et al. 1996).

127. Strengths and Weaknesses: *M. intii* represents an indigenous, ecologically-relevant species for testing contaminants that could negatively affect northeast Pacific coast ecosystems. The life cycle of *M. intii* is much shorter (~20 days) than that of *H. costata* (~70 days), the other Pacific coast species commonly used in toxicity testing. The EPA sponsored the development of a 7-day toxicity test protocol (Langdon et al. 1996) that has been applied (UCSC 1998) and evaluated (Harmon and Langdon 1996). *M. intii* was recently used in a series of tests examining acute and chronic effects of nickel on three species of marine organisms (Hunt et al. 2002). The chronic test was a 28-d full life-cycle test that included survival, growth (as the change in length and weight), and two reproductive endpoints (the percentage of females carrying eggs or juveniles in the brood sac, and the number of live juveniles produced). However, during this test no juveniles were released and the number of gravid females was low in the test controls. Hunt et al. did not discuss possible explanations for the reproductive failures. The primary disadvantages associated with using *M. intii* is the lack of available commercial culture and that testing protocols to measure EDC-related endpoints need to be developed. Regardless of source of test animals, *M. intii* individuals used in testing must be identified carefully prior to their use in tests. Further, the requirement to supplement an *Artemia* diet with copepods (*T. californicus*), which initially must be field collected, could be an impediment to the use of *M. intii* by some laboratories. Finally, potential population-related differences in pollutant sensitivities should be assessed before comparing the results of tests performed on different populations.

4.3.5 *Neomysis integer* (Leach, 1814)

128. *Neomysis integer* is found throughout northern Europe (Mees et al. 1994) and has been suggested as an appropriate species for use in European toxicity testing programs (Roast et al. 1998, 2000b).

129. Natural History: *N. integer* is a relatively large, hyperbenthic species that occurs in relatively low-salinity portions of estuaries (Roast et al. 2001). Females may attain a standard length of about 18 mm (measured from the base of the eyestalk to the end of the last abdominal segment); males are smaller (Mees et al. 1994). Brood size is strongly correlated with the size of the female: the number of larvae per brood extends to about 80 individuals for females of 16 mm or more in length (Mees et al. 1994). Winkler & Greve (2002) determined that individuals matured in about 45 d after hatching at 15°C, but at about 3.5 months at 10°C; the total generation time was about 69 d at 15°C.

130. Availability, Culture, and Handling: Because it is the dominant mysid inhabiting northern European estuaries (Mees et al. 1994), it is readily available, but animals to be used in testing must be field

collected. It is not cultured commercially, but wild-collected animals are easily maintained in the laboratory.

131. **Strengths and Weaknesses:** The principal strength of this species for use in toxicity testing is that it is a common and ecologically important component of European estuaries. It is also a very well-studied species, ecologically, physiologically, and toxicologically. However, it is not commercially cultured and all animals to be used in tests must be collected from estuaries and raised in the testing laboratories. *N. integer* can be tested at relatively cool temperatures of about 15°C, although it is probably not a useful species for testing at colder temperatures (~10°C) because of the much longer generation time at cold temperatures. Testing to measure potential EDC-related endpoints and studies addressing some aspects of the endocrinology and of this species, and its potential utility in EDC testing, has recently been completed. Verslycke and Janssen (2002) developed an indicator, the CEA, that could be used to detect changes to energy metabolism of *N. integer* in response to environmental stressors and tested the effects of tributyltin chloride on the CEA (Verslycke et al. 2002, 2003a). Verslycke et al. (2002) found that *N. integer* produced 11 monohydroxy testosterone metabolites and two nonpolar metabolites (androstenedione and dihydrotestosterone). They also found the anabolic steroid β -boldenone, which had not been previously reported in invertebrates. The function of the steroids in mysids is not clear. Verslycke et al. (2002) did not detect the vertebrate estrogen 17 β -estradiol in *N. integer*. Biotransformation experiments conducted by Verslycke et al. (2002) revealed that mysids have a complex steroid hydroxylase system comprised of several P450 isozymes. Alteration of P450 activity in mysids could be used as an endpoint in EDC testing. Verslycke et al. (2003b) tested the effects of TBTCI on testosterone metabolism in *N. integer* and found nonpolar and polar metabolite induction at the lowest TBTCI concentration (10 ng/L) tested, but there were no significant differences from controls at higher concentrations. TBTCI had no effect on the elimination of testosterone by glucose conjugation, in part because of high within-treatment variability. Testosterone elimination by sulfate conjugation was significantly lower at high TBTCI concentrations. [Verslycke et al. calculated a metabolic androgenation ratio (the ratio of the oxido-reduced products to the hydroxylated plus conjugated products), which has been used to interpret the total effect of a chemical on testosterone metabolism, and found no significant treatment-related differences from the control.] Despite this, they contended that the ratio could be used to summarize the effect of TBTCI, which was an increase in the ratio at the two lowest concentrations. What was also clear in the data, but not discussed, was the very high variability in the ratio within treatments, especially within the controls. Control variability, as indicated by coefficient of variation values of almost 100%, was very high, which calls into question the general utility of the metric in evaluating endocrine disruption.]

132. Gorokhova (2002) stated that to understand growth of crustaceans (which is a potential EDC endpoint) it is imperative to understand the molt staging of the organism. Gorokhova determined that the duration of the molt cycle of *N. integer* was about 9 days and that the effects of salinity and temperature on molt staging differed according to the food supply.

133. These animals must be identified carefully prior to testing. Also, potential population-related differences in pollutant sensitivities should be assessed before comparing the results of tests performed on different populations.

4.3.6 *Other mysid species*

134. A few other species of *Americamysis*, including *A. almyra* (Bowman 1964) and *A. bigelowi* (W. Tattersall 1926), have been used in toxicity testing or related studies. *A. almyra* is closely related to *A. bahia*, and the two species have similar geographic distributions; however, *A. almyra* inhabits less saline waters (Price et al. 1994). The reproductive biology including brood size and generation time of the species is very similar to that of *A. bahia* (reviewed in Reitsema and Neff 1980). The species is amenable to laboratory culture. Recirculating (Reitsema and Neff 1980) and static (Domingues et al. 1998, 1999)

culture systems have been developed. *A. almyra* can be maintained on a diet of *Artemia* nauplii (Domingues et al. 2001a, 2001b). *A. bigelowi* is also biologically similar to *A. bahia* and to *A. almyra*. It occurs along the east coast of the United States from Massachusetts to Florida (Price et al. 1994). Although Gentile et al. (1982) found it to be suitable for use in toxicology testing, *A. bigelowi* has not received widespread attention as a test species.

135. *Neomysis mercedis* Holmes 1897, an ecologically important Pacific coast species, was advocated as an acute toxicity test organism appropriate for estuarine waters having low salinities, ranging from 1‰ to 3‰ (Brandt et al. 1993). The suggested temperature range for testing with this species is 16°C to 19°C (Brandt et al. 1993); thus it could serve as a cool water alternative to *A. bahia*. It occurs in freshwater and brackish waters from California to southern Alaska (Daly and Holmquist 1986) and recently has been used in toxicity testing (Farrell et al. 1998a, 1998b; Hunt et al. 1999, 2002). *N. mercedis* is an important predator on *Daphnia* in a freshwater lake (Lake Washington, Washington), where its consumption is enough to control *Daphnia* populations (Chigbu 2004). Brandt et al. (1993) found that *N. mercedis* was similar to *A. bahia* in sensitivity to several environmental contaminants. A standardized acute toxicity testing protocol has been developed for *N. mercedis* (ASTM 1997). *N. mercedis* is not cultured commercially, but can be reared in the laboratory (Brandt et al. 1993). *N. mercedis* has a relatively long generation time of about three to four months (Brandt et al. 1993). The tests required to measure many EDC-related endpoints must be developed (Ingersoll et al. 1999) or are impractical because of the species' long generation time.

136. *Neomysis americana* (S.I. Smith, 1874) is a western Atlantic species that occurs on sandy bottoms at depths of 0 to -240 m from Florida to Newfoundland, and also South America (Anderson et al. 2004). *N. americana* plays a dual role in marine food webs, being an important item in the diets of several species of demersal fish (Steimle et al. 2000) and a significant predator on zooplankton (Winkler et al. 2003). The species was used in some toxicity tests in the late 1970s, early 1980s, but has been used very little. The U.S. EPA lists *N. americana* as an alternative species for an acute effluent testing protocol (EPA 2002a), but not for chronic effluent testing (EPA 2002b). Typical test temperatures are 20°C and 25°C and the typical salinity range is from 10‰ to 32‰ (EPA 2002a).

137. *Praunus flexuosus* (Müller 1776) is one of the two predominant mysid species in northern European waters, particularly in shallow water in the outer Elbe Estuary, where it is predominant in summer (*N. integer* is other abundant species) (Winkler and Greve 2002). *P. flexuosus* is tolerant of a wide range of salinities and temperatures. It is not commercially cultured and must be collected from coastal waters before testing. Two studies by Garnacho and coworkers have highlighted one of the important caveats that must be recognized when using wild-collected animals for EDC evaluations. Garnacho et al. (2000, 2001) examined the effects of copper on survival and the metabolism of *P. flexuosus*. The mysids were fed <48-h old *Artemia* nauplii during the holding period and daily during testing. Testing was conducted with water at a salinity of 33‰ and temperatures of 10°C and 20°C. Water and test solutions were renewed every 48 h. Copper was found to be toxic to the mysids collected and tested during the summer, whereas it was not toxic to mysids collected and tested during the winter (Garnacho et al. 2000). Garnacho et al. found that the O:N ratio varied seasonally, being 2.5 times greater in winter than in spring. Metabolism of species is protein based all year. Garnacho et al. also found that the O:N ratio decreased quickly (24 h) at the highest copper dose and decreased in all doses by 10 days. The response in summer was faster than that in winter. The shift in O:N ratio reflects a greater metabolic reliance on protein. The faster response in summer occurs when metabolism is already more dependent on protein. The observed response to copper differs from other mysid responses to stress. Garnacho et al. concluded that O:N ratio changes appears to be a sensitive early indicator of stress in mysids. However, one significant finding from the two studies was that the effects of a particular toxicant or stressor on animals that are wild-collected may vary importantly depending on the season during which the animals were collected. Winkler and Greve (2002) determined that *P. flexuosus* did not mature at a test temperature of 10°C, but

did grow to large size (20 mm). At 15°C, individuals matured at about 3.5 months after hatching and reached a mean size of 16 to 18 mm. The mean incubation time within the marsupium was about 23 d at 15°C, and the total generation time (from egg to mature individual) was about 133 d. Despite its ecological importance, *P. flexuosus* would not be a useful colder water test animal, and its long generation time makes it undesirable for EDC evaluations.

138. *Tenagomysis novaezealandiae* Thomson 1900 is found on New Zealand's North and South Islands (Nipper and Williams 1997). The species is abundant and widely occurring. Although it is not available through commercial culture, field-collected animals can be maintained in the laboratory. The species' life cycle lasts about four weeks and laboratory populations can be reproductively active year-round (Nipper and Williams 1997). Mysids can be collected by using hand dip nets and are cultured in the laboratory at a temperature of 20°C, a salinity of 34 ‰, and a light:dark photoperiod of 16:8 hours. Mysids are held at a density of 10–20 individuals/L and are fed newly hatched brine shrimp nauplii daily. Nipper and Williams (1997) conducted several experiments designed to determine the appropriate physical conditions and food regimen for holding and testing the species. One of the more interesting findings from these experiments was that survival was higher when tests were conducted in complete darkness compared to the 16:8-h light:dark cycle. Nipper and Williams attributed this to the typical vertical migration habits of mysids, which involves migration up in the water column at night and resting in dark areas near the bottom during the day. Nipper and Williams (1997) concluded that *T. novaezealandiae* is suitable for use in toxicity testing and can be cultured for year-round testing.

4.4 OTHER SPECIES

139. Estuarine and marine communities are comprised of taxa from many phyla other than the Arthropoda. Some taxa, especially polychaete worms, are of very high ecological importance and are probably very susceptible to the EDCs. Many frequently are used for various types of regulatory toxicity testing. However, the endocrine systems of most have not been studied sufficiently, nor have adequate protocols been developed, for these taxa to be included in an EDC evaluation program. Nonetheless, three of these groups of taxa are briefly reviewed in this section.

140. Polychaetes: Polychaete worms comprise an important component of most estuarine and marine ecosystems. Primarily infaunal sediment dwellers, polychaetes contribute significantly to sediment bioirrigation and occupy a key part of marine food webs. They represent an ecologically significant group on which to evaluate the effects of potential EDCs. Several species of polychaetes are widely used in sediment toxicity testing. Polychaete (e.g., *Nereis*) bioaccumulation assays are one of the primary tools used in evaluating the suitability of dredged material for disposal into U.S. coastal waters (EPA/USACE 1998 ITM). A general toxicity testing protocol has been developed for several species of nereid polychaete worms, including a chronic survival and growth assay for *Neanthes arenaceodentata* (ASTM 2000) and a fertilization/embryo-larval development assay for *Platynereis dumerilii* (Hutchinson et al. 1995). *P. dumerilii* has been widely used in Europe, especially in genotoxicity studies (e.g., Hagggar et al. 2002) and some aspects of its reproductive endocrinology have been studied (review in Andries 2001). Several age categories of *N. arenaceodentata* are available from a commercial supplier. Several breeding cultures of *P. dumerilii* have been established at European laboratories (e.g., the European Molecular Biology Laboratory in Heidelberg, Germany). The species can be bred continuously in the laboratory (Fischer and Dorresteijn 2004) and may eventually be a good candidate for EDC evaluations. The prime weakness in using polychaetes for EDC evaluations probably is the lack of understanding of functional roles for hormones in polychaetes (Andries 2001).

141. Mollusks: Mollusks may seem an obvious choice for use in EC evaluations because imposex is frequently offered as the best example of endocrine disruption in invertebrates (Section 3.2). Bivalve (e.g., *Macoma*) bioaccumulation studies are important in contaminated-sediment evaluations (EPA/USACE

1998), general chronic assays have not been developed. Endocrine disruption in marine gastropods has been studied considerably, almost exclusively with respect to the induction and incidence of imposex. However, the effects, including reproductive endpoints, of some endocrine disrupting chemicals on the hermaphroditic freshwater snail *Lymnaea stagnalis* have been studied (Czech et al. 2001, Coeurdassier et al. 2004). *Lymnaea peregra* was included in a multi-species multi-generation, whole life-cycle testing protocol developed by Sánchez and Tarazona (2002). However, at this point suitable chronic protocols for estuarine and marine mollusks have not been developed that could be adapted for a chronic reproductive assay.

142. Echinoderms: Echinoderms are often included in some regulatory testing protocols, particularly those involving effluent discharges or the effects of the suspended particulate phase of sediments, but these focus on acute effects on embryonic and larval development (e.g., EPA 2002b, 2002c). Some echinoderms have been used in studies of specific EDCs (Section 3.2), but these do not involve chronic procedures or reproductive endpoints. Echinoderms would be of interest to an EDC evaluation program because they are known to have steroids that also occur in mammals (LeBlanc et al. 1999). Currently, echinoderms do not seem well-suitable to multi-generational EDC evaluations.

5.0 SELECTION OF AN APPROPRIATE TEST SPECIES

143. Several characteristics should be considered in determining the most appropriate species for use in EDC-related testing. Nimmo and Hamaker (1982) and Roast et al. (1998) suggested that the test organism should be ecologically relevant, and sensitive to contaminants. Because it must be available as needed for testing, it must be either abundant and easily collected, or amenable to laboratory culture. Its diet should be well understood and easy to provide in the laboratory, therefore allowing it to be well-adapted to laboratory conditions and reducing the need for extensive acclimation periods. The ideal test organism should have a short, relatively simple life cycle that allows for the testing of successive generations.

144. Ingersoll et al. (1999) included several additional characteristics that are important considerations in the selection of test species. These characteristics include the species' mode of reproduction and knowledge of their endocrinology. Species that reproduce parthenogenetically may produce a test population within which genetic variability is relatively low compared to species that reproduce sexually. High genetic variability may yield highly variable responses to a stressor within a test population (Lagadic and Caquet 1998), which may make it difficult to detect impacts other than those that are substantial. However, using parthenogenetically reproducing animals means that important processes related to sexual reproduction (e.g., gametogenesis) are not being evaluated (Lagadic and Caquet 1998, Olmstead and LeBlanc 2000). Knowledge of the endocrinology of a candidate species is important to the selection of appropriate test endpoints and the interpretation of the test results (Segner et al. 2003). Also important to selecting a candidate species for use in the multigenerational EDC testing program are the ability to culture the species in the laboratory (with a strong probability that transgenerational testing is possible); relatively short generation time, allowing for full life-cycle testing; size (larger animals provide more tissue for measuring hormone titers, but usually have longer generation times); and the availability of standard (consensus-based) testing methods, including whether or not new methods must be developed to measure EDC-relevant endpoints.

145. Recent studies of the genetic variability within a common and widespread species emphasizes the need to document the test species identity and the source from which the animals originated. Several studies have shown that the freshwater amphipod *Hyalella azteca* is a species complex comprised of at least a dozen species and that different genotypes may vary in their response to the same stressors (discussed in detail in section 4.1.2). The identity of test organisms should be verified and documented through a voucher collection, as described by Huber (1998), that is maintained by the testing laboratory.

5.1 CRUSTACEANS AS REPRESENTATIVE AQUATIC INVERTEBRATES

146. Many insecticides are considered to be EDCs, because they are specifically formulated to attack insect endocrine systems, affecting in particular the systems that are involved in molting and larval metamorphosis (Oberdörster and Cheek 2001). Most of these insecticides are JH analogs (Oberdörster and Cheek 2001) or ecdysteroids. Although insects and crustaceans represent two classes within Phylum Arthropoda and consequently exhibit many similarities as well as differences, several recent studies showed that insecticides formulated as JH analogs adversely affect crustacean larvae by disrupting molting and metamorphosis (e.g., see McKenney and Celestial 1996, which focuses on mysids). Crustaceans probably do not synthesize JH (LeBlanc et al. 1999); however, they do produce methyl farnesoate in the

mandibular gland, and it is likely a natural JH analog. It is known to be involved in crustacean reproduction, but its specific role is uncertain (LeBlanc et al. 1999).

5.2 MYSIDS AS REPRESENTATIVE CRUSTACEANS

147. Mysid crustaceans are distributed from 80 °N to 80 °S and occur in most aquatic environments, including brackish, freshwater, and marine (e.g., Mauchline 1980). In some habitats, particularly coastal temperate waters, mysids are very abundant and are very important in freshwater, estuarine, and marine food webs. Mysids diets include consumption of detritus, phytoplankton, and zooplankton. They may be abundant enough to control population densities of some prey (Chigbu 2004). Stomach-content analysis revealed that mysids are a staple food for striped bass, the tidewater silverside, and several species of flounder (Gentile et al. 1983).

148. Mysids, despite their superficial resemblance to decapod shrimp, have been considered more closely related to amphipods and isopods (Brusca and Brusca 1990). However, recent studies of 28S rDNA sequences among several malacostracan orders indicated that mysids (Suborder Mysida) are more closely related to krill (Euphausiacea) than to amphipods or isopods (Jarman et al. 2000). This proposed phylogeny has not been accepted universally (Richter and Scholtz 2001). Regardless of phylogenetic relationships, mysids, amphipods, and isopods are characterized in part by the retention of developing young in a marsupial brood pouch. All three taxa would be good candidates for toxicological testing, and amphipods and mysids are routinely used. However, for EDC testing, especially for life cycle tests, mysids offer clear advantages over amphipods. Most marine amphipods used in toxicological testing must be collected from their natural habitats just prior to use in tests. Although they can be held for a few weeks prior to testing, they generally are not cultured for tests. Currently, only one marine amphipod, *Leptocheirus plumulosus*, has been cultured successfully and used for growth and reproduction tests (EPA 2001). Conversely, several mysid species have been cultured in the laboratory and used in such life cycle tests.

149. Nimmo and Hamaker (1982) advocated the use of mysids in toxicity testing, including acute and chronic life-cycle testing. Nimmo and Hamaker compared acute and chronic toxicity data for a variety of taxa and concluded that mysids were more sensitive than most of the taxa, including grass shrimp, an estuarine fish, and 17 freshwater species. Nimmo and Hamaker foresaw a role for mysids in the evaluation of the effects of pesticides and metals on successive generations, in part because the life cycle of mysids is short. This role is being fulfilled by the inclusion of mysids in an EDC evaluation program. Verslycke et al. (2004a) evaluated the potential for using mysids in an endocrine disruption evaluation program. They reviewed aspects of the biology and ecology of mysids in the context of this potential use and summarized the literature on toxicity studies done using mysids. Their review also included a discussion of many endpoints that may be useful when using mysids to test for endocrine disruption. Verslycke et al. (2004a) concluded that mysids have the ecological relevance and sensitivity to stressors required of a taxon that would be suitable for evaluation of endocrine disruption in marine and estuarine invertebrates and could serve as a surrogate for other crustacean species.

6.0 CONSIDERATION OF APPROPRIATE TEST SYSTEMS FOR REPRODUCTIVE AND DEVELOPMENTAL TOXICITY TESTS

150. Mysids are typically pelagic, and consequently have contact with EDCs through the water-column. The main route of exposure is through their swimming and feeding in the water-column (Roast et al. 1998). Many estuarine mysids are also hyperbenthic (Roast et al. 1998) and make diurnal migrations into the water column (Dauvin et al. 1994). For these reasons, mysids could serve as sensitive indicator species to monitor the effects of EDCs through exposure to either the water column or sediment or through dietary uptake.

6.1 ROUTE OF EXPOSURE

151. In a testing program, the routes of administration could be through spiking a potential EDC of interest into the water-column or into clean sediment, or by mixing the compound with the mysid food source prior to feeding. Because of the particular diet of the mysids proposed for use, the two most practical routes are through the water-column or sediment, as described below.

6.1.1 *Water*

152. In water-column exposures, the test compound is mixed with the dilution water. For poorly soluble substances, use of a saturation column, a solvent carrier, or other technology may be necessary (Lussier et al. 1985; Nimmo and Hamaker 1982; OECD 2000). The dilution water should be acceptable to mysids, be of uniform quality, and should not unnecessarily affect results of the test. For detailed discussion of dilution water, see EPA (1996). If a solvent carrier other than dilution water is used, its concentration in the test solution should be kept to a minimum, and should be low enough that it does not affect the survival, growth, or reproduction of the mysids (EPA 1996; OECD 2000). It is important that the highest concentration of the EDC not exceed the single-phase seawater solubility of the compound. It could require direct measurement of the test chemical's solubility in clean, particle- and dissolved-organic-carbon-free seawater of the salinity to be used in the test.¹ Delivery systems are designed to provide either continuous or intermittent flow of the chemical and dilution water mixture (McKenney et al. 1991). The EDC is mixed with dilution water in the mixing chamber, agitated, and then delivered to the replicated test chambers.

153. Continuous-flow systems (Figure 1) are designed to deliver a constant concentration to the test chamber, often by means of the metered pump of a siphon-flush system, which produces a 50% exchange of volume every 4 h, with an incoming flow rate of 30 mL/min and an outgoing flow rate of 100 mL/min (Gentile et al. 1982). The chosen system should be calibrated prior to use to ensure that the appropriate concentration of the test substance is achieved into the test chambers. The general operation of the

¹ Dissolved organic carbon and particulate organic carbon in exposure water can substantially affect the amount of the toxicant that is in true solution and therefore the most bioavailable to the test organisms. Dissolved organic carbon has a strong affinity for binding nonpolar organic compounds and some metals, which decreases their bioavailability. It also thereby could decrease the exposure concentration of the toxicants, and accordingly, influence the results of the bioassay. Therefore, there probably should be a requirement developed as part of the testing protocol for the acceptable concentration of dissolved and particulate organic carbon in exposure water, particularly when natural seawater is used (J.M. Neff, personal communication, January 15, 2002; ASTM 1997).

delivery system should be checked twice daily, with a target 24-h flow rate through the test chamber equal to at least five times the volume of the testing chamber. Also, the flow rates should not vary more than 10% between replicate chambers or over time (EPA 1996). Table 3 presents some examples of the types of systems used in toxicity testing, along with their advantages and disadvantages.

6.1.2 Sediment

154. Mysids are currently used in routine toxicity tests to examine the potential toxicity of marine sediment (Carr et al. 1998; Cripe et al. 2000). It has been documented (Cripe et al. 2000) that the mysids were observed to collect sediment, manipulate it at the mouth region, and drop it. This suggests that mysids could be used to test sediment suspected of containing EDCs.

6.2 CONCENTRATION SERIES

155. The goal of the chronic life cycle test should be to calculate endpoints such as survival, growth, or reproduction that are inhibited at a specific chemical concentration. If nothing is known about the sensitivity of the mysid to the chemical being tested, then a geometrically spaced series of test concentrations is used to establish a dose-response relationship for the key endpoints. If there is some information concerning the concentration-effect curve then several other approaches could be taken:

- Conduct an acute range-finding test in which the organisms are exposed to a control and three or more concentrations of the test material that differ by a factor of 10. The test design for the acute test should use the same testing conditions and same age of organisms as the chronic test. The results of a water-only acute test could be used to calculate a more precise concentration series to target for the chronic test.
- Alternatively, estimation of concentration series could use an acute-to-chronic test ratio for a species of comparable sensitivity: the result of the acute test can be divided by the acute-to-chronic ratio. Generally, acute-to-chronic ratios determined with saltwater mysids are often less than 5.

156. If no other useful information is available, the highest concentration of test material in a life cycle test with mysids is often selected to be equal to the LOEC in a comparable acute test (ASTM 1997). Tests can also be used to generate the median lethal concentration (LC_{50}), median effective concentration (EC_{50}), or median inhibition.

- concentration (IC_{50}), using a toxicant concentration series (0.5 or higher) should be selected that will provide partial mortalities at two or more concentrations of the test chemical (EPA 2000).

Figure 1. Diluter




Table 3. Types of water delivery systems

Type	Description	Advantages	Disadvantages
Static	solution not changes during testing	no maintenance, inexpensive, performed in large numbers	toxicant concentrations may shift due to uptake by organisms, volatilization, decreased dissolved oxygen, or changes in pH.
Static Renewal	test solution changed at regular intervals using either a manual pump and freshly prepared replacement water	improved consistency of toxicant concentration	replacement of solution may stress the test organisms
Flow-through	solution replaced by automated system using a controlled dosing system a proportional diluter or a continuous flow siphon system	improved consistency of toxicant concentration, savings in labor hours	equipment can be expensive large volumes of test solution may require costly disposal

6.3 STATISTICAL CONSIDERATIONS

157. Statistical approaches for conducting and interpreting mysid reproductive toxicity tests and their relative sensitivity have been presented by a number of authors. The most comprehensive document on the statistical analysis and design of genotoxicity tests was a joint project from ISO and OECD (OECD 2003). This document, which was also circulated within ISO as a working draft (ISO TC 147/SC 5 N 18, ISO/WD 1 “Water Quality – Guidance Document on the Statistical Analysis of Ecotoxicity Data”), covers the differences in the general design and statistical analysis between approaches used to estimate an NOEC and a dose-response modeling approach used to estimate an effective concentration (EC_x). Kuhn et al. (2000, 2001) discuss the extrapolation of standard toxicity endpoints to population risk assessment. Lussier et al. (1985), McKenney (1986), McKenney et al. (1991), McKenney and Celestial (1996), Lussier et al. (1999), and Cripe et al. (2000) provide a measure of variability on survival, growth, and reproduction of mysids. Nimmo and Hamaker (1982) compare the sensitivity in toxicity testing of mysids to other marine species, and EDSTAC (1998) considers the relevance of mysid testing to other invertebrate groups.

158. The objective of a mysid two-generation reproductive and developmental toxicity test is to provide the most precise and accurate estimate of toxicity associated with endocrine disruption and reproductive fitness for an identified potential EDC. The results of the Tier 2 testing should be conclusive in documenting a discernible cause-and-effect relationship of chemical exposure to measurable manifestation in the test organisms. The test protocol will be designed to

- • Determine whether effects are a primary or secondary disturbance of endocrine function
- • Establish exposure/concentrations/timing and effects relationships
- • Be sensitive and specific
- • Assess relevant endpoints
- • Include a dose range for full characterization of effects (EDSTAC 1998).

159. Thus, the assay must be biologically sensitive, have minimal variability associated with dose exposure throughout the test duration, and have a statistically powerful inference. Biological sensitivity is a function of the choice of species tested, the relevance of the endpoints measured to species productivity and survival, and the route, duration, and level of the chemical exposure. Design-associated variability in dose exposure is a function of exposure route and duration, chemical stability and purity within the testing environment, and the testing protocol. The power of a statistical inference is a function of the inherent variability in response; design-associated variability; the degrees of freedom and the source of variability for testing; and the estimation process and decision criteria.

160. Ideally, an experimental design incorporates randomness, independence, and replication (Cochran and Cox 1957). Randomness is used to remove noise, independence is used to extend the inferences made, and replication provides a measure of variability for testing (Chapman et al. 1996). Randomization of 1) experimental containers within a testing environment, 2) treatment application to experimental containers, and 3) assignment of organisms to experimental containers allows one to incorporate the variability associated with the environmental conditions, the containers, and the organism equally across all treatments. Thus, when one evaluates the difference between treatment means, the variability associated with experimental environment, experimental containers, and organisms being treated is removed and only the effect of the treatment remains.

161. Independence of treatment application and the creation of the treatment, and thus, the inference associated with the treatments being tested, incorporate the variability associated with more than one individual, in more than one location, making and applying the same treatment. The random sample of organisms from a given population actually limits the inference to that population. However, one can evaluate the stability of the inherent variability of the population over time. An experimental unit is defined as the group of material or individuals to which a treatment is applied independently in a single trial of the experiment (Cochran and Cox 1957). Replication of experimental units for each treatment provides a measure of all the necessary sources of variability needed to extend the inference across time and space. A reduction in the sources of variability that are truly independent constrains the inference (Hurlbert 1984). Thus, if only one mix of each treatment is made and then divided among replicates, the source of variation associated with making the treatment is not included in the variability for testing, and the inference is limited. Some would say that this variability is nuisance noise, too small to be of concern. Therefore, if this source of variability is not included, it should at least be acknowledged. The variability among replicate experimental units could also include noise that was not randomized out due to a poor randomization or variable measurement error. These sources of variability can be reduced without loss to inference.

162. Statistical power is the probability of rejecting the null hypothesis of equal means when the alternative is true—that is, detecting a difference when there is a difference. Statistical power is a function of the variability among replicate experimental units within a treatment, the number of replicate experimental units, the size of the Type I error, and the percentage difference one wishes to detect. One can control the latter three components; however, the variability in response is inherent in the test organism. Thus, the choice of which species to test and the relevant endpoints to measure should include a comparison of inherent variability or coefficients of variation (CVs), defined as the standard deviation \div mean \times 100%. Data that have high CVs are associated with low power for detecting small-scale differences. For example, with five replicates per treatment, a CV of 50% would rarely detect differences less than 70% between the test and control-treatment responses at a Type I error rate of $\alpha = 0.05$ (Figure 2). For a given CV, one can increase power by increasing the number of replicates. Test species and endpoints with the least inherent variability, by default, require the least replication for a given level of power and thus are more cost-effective.

163. Life-cycle studies provide a number of continuous and discrete random variables over the course of the exposure. Examples of continuous data include growth measured as length and dry weight, biochemical markers, and time to brood release. These data are analyzed using ANOVA and pair-wise comparison techniques to determine difference between treatments and controls, and regression or maximum likelihood techniques to estimate an ECx. Williams' parametric test of ordered alternatives (Williams 1971) or Jonckheere-Terpstra nonparametric trend test (Jonckheere 1954, Terpstra 1952) are used to determine the significance of a specified trend in the response associated with the treatments. Life-cycle studies also produce a large number of discrete data points, such as the molt frequency, morphological characteristics, and survival. These data can be analyzed by ANOVA if arcsine square root transformed, and by pair-wise comparison techniques, contingency table techniques to assess association, and regression or maximum likelihood techniques to estimate an ECx. Fisher's Exact Test is an example of a technique for comparing two sets of discrete quantal data.

164. Data collected by Lussier et al. (1985), McKenney (1986), McKenney et al. (1991), McKenney and Celestial (1996), Lussier et al. (1999), and Cripe et al. (2000) can be used to compare the CVs for selected endpoints. Four- to 26-day survival data for control water and sediment (McKinney and Celestial 1996, Lussier et al. 1999, Cripe et al. 2000) showed a significant increase ($p = 0.01$) in the survival CV with time ranging from 2% to 20%. (Figure 3). The control data CVs for dry weight did not vary significantly with time ($p = 0.64$) and averaged 22% in control water and 7% in control sediment (McKinney and Celestial 1996, Lussier et al. 1999). Reproductive endpoints had CVs of 12% to 55% (Table 4). In general, there are few life-cycle data for which a power analysis can be conducted with great confidence. An increase in CV as a function of dose is dependent on the organism being tested, the exposure chemical, and its toxicity to the endocrine system.

6.3.1 Hypothesis-Testing or Regression Analysis

165. There has been much debate over the use of the NOEC in toxicity assessment and the associated risk analysis (Crane and Newman 2000, Chapman et al. 1996). The desire to detect effects implies a comparison of means. ANOVA methods are appropriate for comparing means, asking the question of whether the treatment means are statistically different from the control, such as in a screening test or a validation test. However, ANOVA methods are not appropriate when a precise and accurate estimate of toxicity and the pattern of response are required. There is also a false positive error rate in ANOVA, because the many parameters assessed are not all independent. Regression techniques provide an estimate of the level of effect as a function of exposure (nominal or actual concentration) and the functional relationship between dose and response. Further, by analyzing the different dose-response relationships, one can compare the sensitivity and potential thresholds of effect for different endpoints.

Figure 2. Power of a one-sided independent-samples t-test as a function of the percentage difference (delta) detected between the test and control means, with 5 replicates per treatment ($\alpha = 0.05$)




Figure 3. Coefficient of variation (%CV) of control mysid survival as a function of time. The solid line is the fitted regression; the dashed lines are the 95% confidence intervals.




Table 4. Control data means and coefficients of variation (CV) ¹

Variable	Number of Studies	Mean Response	CV
----------	-------------------	---------------	----

Days to release first brood	4	20.4	12%
Number of young in first brood	2	23.7	20%
Number of young per female (first brood)	3	5.6	55%
Young per female reproductive day	1	0.55	48%

¹ Lussier et al. (1985), McKenney (1986), McKenney et al. (1991), McKenney and Celestial (1996)

166. Although the NOEC is used widely, it should not be relied on as the sole indicator of low toxicity (Crane and Newman 2000, Chapman et al. 1996, OECD 2003). The largest dose for which statistical differences have failed to be detected is a direct function of the power of the test: failure to reject the null hypothesis of no difference does not mean that there was no effect. For example, for certain endpoints with CVs greater than or equal to 40%, it is unlikely that differences less than 60% will be detected with 5 replicates per treatment (Figure 2). It is also conceivable that short-term range-finding experiments will have difficulty in predicting the location of a NOEC in a multi-generational test. It may also prove difficult to achieve effects bracketing the 50% response in the F1 population. However, effect concentration calculations are an appropriate alternative for estimating doses associated with low to medium toxicity. Care must be taken not to estimate an effects concentration that is more sensitive than the data and the experimental design will allow. Precision and accuracy of the effects concentration is a function of the spread between treatment concentrations and the number of concentrations tested (Chapman et al. 1996).

167. The design and analysis requirements for estimating the NOEC differ from those for fitting a dose-response model (Chapman et al. 1996, Stephan and Rogers 1985). ANOVA methods require experimental unit replication and achieve greater power in testing as a function of the number of replicates. As shown in Figure 2 and Table 4, the different endpoints would require different amounts of replication to achieve the same level of power. However, 5 replicates provide greater than 80% power for detecting less than a 30% change at $\alpha = 0.05$ from the control for some of the growth and reproductive endpoints in Table 4, assuming the CVs do not increase with dose much beyond 20%. Transformation of the data to satisfy homogeneity of variance is required for the parametric test and the regression approach. Estimation of the NOEC should be done with the more powerful tests of ordered alternatives such as Williams' parametric test of ordered alternatives (Williams 1971) or Jonckheere-Terpstra nonparametric trend test (Jonckheere 1954, Terpstra 1952).

168. The design of a study intended for dose-response modeling does not require replication of the treatments (Snedecor and Cochran 1980). Each individual response is assumed to be a random response from a normal population of responses for a given dose. The variance is assumed to be equal for each population. Replication of doses provides a test of equal variance and lack-of-fit (Draper and Smith 1981). Further, because of the unpredictable nature of survival and fertility in the two-generation test and the large variability in specific endpoints, it is desirable to have some level of treatment replication to provide a more accurate estimate of the mean population response for a given dose. The number of replicates would depend on the maximum expected variability in response for each dose. The variability in response may be a function of the dose. In this case, either a weighted analysis should be conducted or a data transformation applied that satisfies the assumption of homogeneity of variance.

169. Benefits of the regression approach include 1) estimation of the pattern, or slope, of toxicity as a function of dose; 2) estimation of the distance between effect concentrations and environmental concentrations; 3) estimation of effective doses (ED_x) and their associated confidence intervals for x equal to a low to medium effect; 4) estimation of ED_x not limited to doses on test; 5) use of both measured and nominal concentrations; and 6) ability to compare dose-response curves across endpoints (Chapman et al. 1996, OECD 2003). The size of the resulting confidence intervals, indicating the precision of the estimated ED_x, is a function of the inherent variability in the response, and the number and spacing of the

concentrations tested. Guidelines often require five concentrations that are geometrically spaced and sublethal, plus a no-dose control. Thus, a range-finding test would be required to determine appropriate dietary concentrations.

170. Regression modeling is sufficiently flexible to handle a wide range of dose-response patterns, including nonmonotonic if enough doses within the affected region are represented. If only one or two responses are not either 0% or 100% affected and at least one is greater than 50% affected, the Spearman-Kärber nonparametric method can be used to estimate a median effective concentration (EC50). Finally, the regression approach can handle a wide range of responses, including continuous responses, counts, and quantal data, by re-expressing or transforming the data (e.g., $\log[y+c]$, $[y+c]^{1/2}$), and probit, respectively). At a minimum of intermediate responses, a significant linear slope would represent a significant average effect. Simple linear regression, however, would not be able to detect a low-dose effect that is not observed consistently at higher doses.

171. There is a significant lack of information, however, on determining a protocol based on a regression approach. As stated above, the number of replicates and doses required for testing has not been established to estimate an EC_x with a desired confidence interval width. Neither the percentage response required to be estimated nor appropriate dosing strategies have been determined. Other statistical considerations that need to be addressed are strategies for

- Determining the presence or absence of low dose effects;
- Selecting the number and range of treatments when many endpoints of varying response ranges; must be characterized simultaneously using the regression approach; and
- Detecting and fitting nonmonotonic responses.

172. A blending of the ANOVA approach with the regression approach may be the best way to anticipate potentially extreme responses. A low dose response can be statistically detected using ANOVA while nonlinear regression allows a comparison in functional response between endpoints. The difficulty is balancing the larger replication needs of the ANOVA design with the increased number of doses required by regression and the ultimate cost of the experiment. A design with a greater number of replicates in the control and lowest dose and a smaller number of replicates for the remaining doses would allow detection of a low-dose effect, regression analysis, and maintain low cost.

6.3.2 Statistical Versus Biological Significance

173. The determination of statistical significance in hypothesis testing means that a value greater than or equal to that observed of a specific statistic has a small probability of occurring by chance alone. The definition of small is used to define the level of significance and is usually set at 5%. The p-value is the probability that a statistic will have a value greater than or equal to the achieved value. It is calculated based on specific assumptions about the distribution of the statistic assuming the null hypothesis is true. The decision to reject a null hypothesis for a defined alternative is based on the level of significance chosen before analysis. However, presenting the achieved p-value allows one to assess the potential biological significance of the result as well.

174. The biological significance of a result is a function of the sample size and the sampling strategy as well as the biological implications of a result under a wider array of variables than the statistical inference. At a minimum, a biologically significant result would be expected to affect a population's viability. Implications of biological significance may be used to hypothesize community effects through food webs and ecological modeling. For statistical inference, the sample size must be large enough to

encompass the variability in the population for which the inference was intended or in other words the sample must be representative of the population. For biological inference, the sampled population must provide relevant information for a larger targeted inference space. Power calculations are recommended to determine the sample size required to detect a biologically significant effect and, thus, to reduce the number of statistically significant effects that are irrelevant biologically. Unfortunately, the knowledge needed for this calculation is often lacking.

175. Population modeling was used to evaluate the ecological relevance of the mysid bioassay (Kuhn et al. 2000, 2001). Survival and reproductive endpoints were highly correlated with a population model-derived contaminant concentration associated with zero population growth for 15 metals and 8 organic contaminants (see Section 3.3.2). Kuhn et al. (2001) found that population-level effects on mysids can be predicted from multigenerational tests. Further, Nimmo and Hamaker (1982) found that mysids were found to be as sensitive or more sensitive to toxic substances than other marine species. Thus, it is possible that protective actions based on mysids may be relevant to a wider group of species. However, EDSTAC (1998) points out that there are not sufficient data to infer what level of protection would be provided to other invertebrate groups.

6.4 DESCRIPTION OF ASSAY ENDPOINTS REFLECTIVE OF REPRODUCTIVE AND DEVELOPMENTAL IMPAIRMENT

176. In considering the list of potential endpoints presented in the following sections, it is desirable that the measured response is directly related to exposure to the potential EDC and that the response is the result of interruption of endocrine function (Ingersoll et al. 1999). Although many, if not all of the endpoints described in the following sections could indicate a response to an EDC, most also will vary in response to exposure to other stressors. Further confounding the interpretation of testing results is the interrelatedness of some endpoint measurements. Thus, a stressor could act on a mysid to reduce its swimming ability, thereby reducing its ability to feed, which then reduces its growth. Similarly, reduced fecundity could be an expression of reduced growth. Often, exposure to a compound that results in reduced fecundity also results in reduced growth (Nimmo et al. 1981); however, this is not always the case. Lussier et al. (1999) reported fecundity effects without reduced growth, and McKenney and Celestial (1996) found reduced reproductive measures at concentrations of methoprene below that which resulted in reduced growth. Various endpoint measures can have differing sensitivities to stressors. For example, reproductive parameters are often, but not always more sensitive measures of contaminant toxicity than is simple survival (Lussier et al. 1985). Hollister et al. (1980) found no reproductive impairment among test organisms, in spite of significant mortality due to exposure to alkaline effluent. Additionally, some reproductive measures can be affected by stressors, whereas others are not (e.g., Hollister et al. 1980, Lussier et al. 1985). Clearly, the possible interrelationships among potential endpoints and the varying sensitivities shown by some endpoints underscore the need to use multiple endpoints in EDC testing.

6.4.1 Growth, morphological and behavioral alterations

177. Measures of growth, morphological changes, and changes in test organism behavior as indicators of EDC effects are reviewed below.

6.4.1.1 Growth

178. Molting is one of the key arthropod physiological processes that is under hormonal control, and therefore, it is susceptible to the negative effects of EDCs. Molting is regulated primarily by the interaction of molt stimulating hormones (ecdysteroids) and nervous system secretions produced in the cephalothorax with molt inhibiting hormones produced in the eyestalks (e.g., Cuzin-Roudy and Saleuddin 1989, Fingerman 1997, Subramoniam 2000, Zou and Fingerman 1997). Molting is either directly or indirectly involved in the expression of the various endpoints that may be examined through toxicological testing. Because noticeable growth can only occur as a result of molting, any disruption of molting could result in alterations in growth. Many pesticides, generally classed as Insect Growth Regulators (IGR), have been developed that target insect development. IGRs function as ecdysone agonists, JH or anti-JH analogs and as chitin synthesis inhibitors. All can have detrimental effects on crustaceans, especially through interruption of the molt cycle, most likely by impacting endocrine function (Touart 1982). For this reason, the estimation of molting frequency may be a useful endpoint relevant to mysid EDC testing. The duration of the molt cycle of adult mysids (*Americamysis bahia*) was determined to be about 6.6 d (Touart 1982), with the duration for male being slightly less and that for females being slightly more. Juveniles are expected to molt within the first 24 h post release, but may delay molting until about 48 h (Touart 1982). Successive juvenile molts occur at increasingly longer durations. Touart (1982) found that at test conditions of 22°C and 20‰, sexual characters were noticeable after the fourth molt (9–12 d) and that mating occurred after the fifth or sixth molts (17–19 d). Higher test temperatures would likely shorten the duration between molts. Molt staging relies on changes in the integument and is most commonly divided into four stages: postmolt, intermolt, premolt, and molt (Verslycke et al. 2004a). Winkler and Greve (2002) found that temperature, body size, and age all are factors contributing to mysid growth rates. Touart also found that the pesticide diflubenzuron (Dimilin®) increased the duration of the molt cycle in A.

bahia, probably acting on the mysid endocrine system as a molt inhibitor. This was demonstrated by his experiment in which eyestalk-ablated mysids that were then exposed to dimilin had molt cycle durations that did not differ from those in control animals. In effect, the Dimilin may have replaced or mimicked the “natural” mysid molt inhibitor synthesized in the eyestalks. Zou and Fingerman (1997) suggest a certain degree of overlap between molting hormones and estrogen-mimicking xenobiotics. They tested this hypothesis using two compounds, DES and endosulfan. Molting time for daphnids exposed to DES and endosulfan was delayed because these compounds blocked the ecdysteroid receptors, preventing molting hormones from binding to the receptor and turning the receptor on, thereby slowing the molting process. However, it is unclear whether the delayed molting process was due to the exposure to these contaminants or just a general response to stressors. More pharmacokinetic studies are underway to investigate delayed molting further.

179. Because there are current immunoassay methods for quantifying ecdysteroids and also methods for molt staging, evaluations of potential interactions of EDCs and molting should be possible (Verslycke et al. 2004a).

180. Two direct measures of growth may be used in the assessment of sublethal effects on mysids. Probably the most common measurement is the determination of simple dry weight. At test termination, animals are briefly rinsed in deionized water to remove salt, and dried to constant weight. In published studies that employ this method, drying temperatures ranged from 60°C (e.g., McKenney 1982) to 105°C (Khan et al. 1992) and drying times ranged from 16 h (Khan et al. 1992) to 72 h (UCSC 1998). ASTM (1997) specified that growth should be measured by drying animals at 60°C for 72 h to 96 h, or to constant weight. ASTM also recommended that separate determinations be made for males and females. Ash-free dry weight (AFDW) is often used as the appropriate weight measurement for many invertebrates, because the technique reduces any inaccuracies introduced by inorganic constituents in the organism’s body. Inorganic components can originate from processes such as the development of skeletal components or feeding (the ingestion of sediment). As with some other small crustaceans, the small size of the mysids and their planktonic diet may make the removal of ash from the dry weight measurement an unnecessary step that would not greatly improve the accuracy of the measurement (EPA 2000).

181. Another direct measurement of growth is body length. Body length has been measured as the distance from the base of the eyestalks to the tip of the telson (Hunt et al. 1997; UCSC 1998) or to the tip of the exopod (Langdon et al. 1996; Winkler and Greve 2002). ASTM (1997) suggests that body length should be measured along the midline of the body from the tip of the carapace to the tip of the exopod (excluding the terminal setae). ASTM notes that it is difficult to make this measurement on preserved animals because of the body curvature that results from the fixation process. Langdon et al. (1996) reduced the potential inaccuracy in this measurement by relaxing the mysids prior to fixation and then determining length as the sum of a series of relatively straight-line measurements. Winkler and Greve (2002) anaesthetized mysids in soda water to relax organisms prior to collecting length measurements. Age-specific growth rates were calculated using duration of the successive intermolt periods in days and the growth factor at each molt. Comparisons of different temperatures (10°C and 15°C) different size classes, and other mysids provided a combination of results. The mysid *N. integer* reached maturity in a shorter time and at a smaller size than *P. flexuosus*. *P. flexuosus* grew faster at all stages but matured much later. Both species had temperature-dependent growth curves but in opposite ways. *N. integer* had a lower growth rate at 10°C than at 15°C.

182. The most common response to exposure to toxicants is reduced growth (several studies) although it may not be a particularly sensitive endpoint (Lussier et al. 1999). McKenney and Celestial (1996) reported reduced growth after exposure to methoprene, a juvenile hormone analog. Reduced growth has important connections to reproductive success because the size of females is directly related to fecundity (Verslycke et al. 2004a). However, reduced growth occurred only at the highest concentration in their

exposure series. It may not be useful to measure growth for all treatments. Because growth is a sublethal endpoint, its use has sometimes been restricted to treatments for which survival was not significantly less than that in control treatments (Hunt et al. 1997). Hunt et al. (1997) observed that individuals that survive high toxicant concentrations may often be larger than average. In a study conducted by Hunt et al. (2002), survival of the mysid, *Mysidopsis intii*, was the most sensitive of five endpoints measured during a 28-day chronic exposure to nickel. Growth is less sensitive in such cases because surviving mysids are large, more tolerant of toxicants, and use cannibalism as a nutritional source. The effect of contaminant exposure on growth may be related to test organism age in addition to contaminant concentration. McKenney (1986) found that older (9–16 d) juvenile mysids showed significantly reduced growth at a lower fenthion concentration than that found for younger (0–9 d) juveniles. Reduced growth in mysids possibly results from a transfer of energy from growth mechanisms as an organism attempts to counteract stress (McKenney 1985, 1989, McKenney and Matthews 1990; Section 6.4.3 below). Reduced growth has strong implications for reproductive success in mysids as several studies (e.g., Mees et al. 1994, Turpen et al. 1994) have shown that fecundity is directly related to female body size.

183. Molt stage has become an increasingly used technique for measuring growth. If molt stages are classified, duration of different stages under normal laboratory conditions, and then environmental effects on relative duration of stages can be evaluated, using the molt-stage technique (Gorokhova 2002). Molt cycle chronology is a prerequisite for the use of molt staging in growth studies. One study was conducted using laboratory reared juveniles and subadults of *Mysis mixta* and *Neomysis integer*; molt staging techniques were used to determine the main molt stages for each species under different regimes of temperature and feeding. Field application of the molt stage technique was applied for *Mysis mixta*, to determine molt cycle duration.

6.4.1.2 Morphology

184. Changes in morphology resulting from exposure to contaminants have been documented for many taxa, including arthropods. For example, chironomid (midge) larvae, which are commonly used in toxicological testing, can develop mouthpart deformities when exposed to chemicals that affect endocrine systems (Meregalli et al. 2001, Watts et al. 2003). The deformities result from physiological disruption during molting. Meregalli et al. (2001) found that sublethal concentrations of 4-*n*-nonylphenol, a known endocrine disruptor, caused mouthpart deformities in *Chironomus riparius*. Relatively low concentrations of 17 α -ethinylestradiol and bisphenol A (10 ng/L) caused mouthpart deformities. Typically, deformities include, loss, addition, fusing, and splitting and these are assigned a numerical score. When more than one deformity was observed, the scores were summed to provide an overall total score. Deformities were assessed relative to the normal arrangement of each of three structures (mandibles, the mentum, and the ephipharyngeal pecten). Observations were made by using a scanning electron microscope, and the most notable deformities were associated with the mentum. In the same study molting in response to these chemicals was also recorded and showed no statistically significant differences at the higher concentrations. This difference in endpoint sensitivity highlights the need to select multiple endpoints for assessment of potential EDCs and the need to understand the mechanism involved in selected effects noted for one mouth part over others such that screening tools could be developed for compounds such as EDCS that may demonstrate a specific deformity in response to those contaminants. Despite the similarities in molting physiology between insects and crustaceans, it does not appear that morphology has been widely considered as an appropriate and measurable endpoint in mysid toxicological studies. A laboratory study showed that exposure to cadmium resulted in the development of abnormal genitalia in male *A. bahia* and malformed carapaces in males and females (Gentile et al. 1982). The time to the first appearance of the abnormalities was positively related to cadmium concentration. One study based on field-collected data found that four populations of *N. integer* contained many individuals with abnormal telson morphology (Mees et al. 1995). Such abnormalities could arise through physiological perturbations that occur during molting, and they could provide a quantifiable measure of disruption in endocrine-related functions.

185. The determination of the degree of fluctuating asymmetry (FA) found among mysids could provide a quantifiable, repeatable measure of morphological perturbations resulting from exposure to possible EDCs. FA is the asymmetric development of a normally symmetrical bilateral structure in which there is no tendency for one side of the structure to have a larger value than the other (Palmer and Strobeck 1986, Palmer 1994, Leung et al. 2000). FA is thought to arise because of environmental or genetic stress during development (Clarke 1993, Leung and Forbes 1996) and may result from a shift in metabolic energy from systems that maintain developmental stability to those that help organism compensate for increased stress (Sommer 1996). Measurement of FA, once appropriate characters have been determined, is relatively easy and requires only use of a microscope. Characters that have been measured are readily observable and include midge larvae teeth (*Chironomus*, Clarke 1993), copepod body spines (*Tisbe*, Clarke 1992), and shrimp antennae (*Palaemon*, Clarke 1993). However, Leung et al. (2000) mentioned that the measurement of single characters does not always reliably indicate environmental stress, and they suggested that an approach involving the use of composite indices of FA could increase the probability of detecting environmental stress. Although some studies of FA have been done on crustaceans (e.g., Clarke 1992, 1993), none has focused on mysids. Before FA could be used to detect mysid developmental abnormalities resulting from exposure to EDCs, preliminary studies examining several suites of potential characters would have to be performed.

186. Laufer et al. (2002), using male spider crabs *Libinia emarginata*, showed that ecdysteroids and low methyl farnesoate concentration (controlled by standard eyestalk ablation experiments) promoted allometric growth (disproportionate growth of body parts; for example, propodus of male spider crabs becomes disproportionately larger than the carapace during reproduction) while ecdysone and higher concentrations of methyl farnesoate inhibited allometric growth. Again, researchers need to understand the mechanistic causes before these changes can be attributed to effects.

187. The use of histology for assessing impacts associated with EDCs has been largely unexplored as an endpoint for crustaceans. One study, using the amphipod *Gammarus pulex*, examined impacts associated with a sewage treatment plant. A highly significant number of females collected displayed abnormal structure of oocytes in vitellogenesis. Particularly noted were uneven ooplasm, few yolk bodies, and lipid globules that were present and unevenly distributed. In addition, both the males and females had decreased body size possibly related to interference with the ecdysteroid receptor, which could have impacts on the molt cycle and interfere with vitellogenesis (Gross et al. 2001).

6.4.1.3 Behavior

188. The disruption of mysid swimming behavior is one endpoint that has recently been investigated as potentially informative in documenting sublethal exposure to contaminants. In a series of studies, Roast et al. (1998, 2000a, 2000b, 2001) determined that sublethal concentrations of cadmium and chlorpyrifos (an organophosphorus pesticide) significantly altered the swimming behavior of *N. integer*. Perturbations included decreased ability to swim against a current, an increase in general activity with no improvement in ability to swim against the current, and a reduced tendency to maintain a position near the bottom. These changes in behavior could have important ecological impacts to the animals by causing them to be moved to an unfavorable habitat or by increasing their susceptibility to predation, thus indirectly resulting in a lethal response to contaminants at sublethal concentrations. However, if the disruption of swimming behavior is to be used as an endpoint in endocrine disruption studies, it should be applied with caution. Roast et al. speculated that the disruption of swimming that they observed was probably a nervous system effect related to interference with cholinergic pathways.

189. Reduced feeding has been noted as a sublethal response to exposure to some contaminants. For example, Nimmo et al. (1981) noted, but did not quantify, reduced mysid feeding in response to exposure to some pesticides. Although difficult to accurately quantify, reduced feeding no doubt is an important

sublethal effect of contaminant exposure that may be expressed in many of the life cycle parameters described below. Reduced feeding could lead to reduced growth, reduced time to maturity, and reduced egg production, among other factors.

190. One study using the amphipod *Gammarus pulex* looked at the combined effects of water and dietary uptake of 4-nonylphenol. Most feeding studies usually focus on one route of exposure and assume a constant feeding rate during the course of the study. The exposure route for benthic invertebrates includes direct contact with the sediment and porewater, uptake from the water column, and ingestion of food and other particles. 4-nonylphenol is more likely to be absorbed to the sediment particles because of its hydrophobic nature. The results of this study showed that feeding rates can be determined in a variety of ways depending upon how the data are analyzed (variable feeding rate versus mean feeding rate). The study showed that significant uptake of 4-nonylphenol occurred from food.

6.4.2 Measures of reproductive performance

191. There are several measures of reproductive performance that can be used to assess sublethal response. For example, sexual maturity, the time to first brood release, the time required for egg development, fecundity, and alterations in reproductive characteristics in populations have all been used as endpoints. Zou and Fingerman (1997) showed that conditions of crowding, food shortage, and change in length of day can trigger a change from asexual to sexual reproduction, with males being produced initially. Females receive environmental cues, which trigger the ovaries to lay male-producing eggs. Exogenous agents may interfere with male differentiation and may affect this process of environmental cues to the ovaries.

6.4.2.1 Sexual Maturity

192. Khan et al. (1992) asserted that sexual maturity, which they described as the presence of gonads or a brood pouch, is a feasible endpoint for reproductive tests, because gonad maturation is essentially the first step toward reproductive output. Maturity allows for the measurement of effects to both males and females. Khan et al. used a dissection microscope to examine live mysids for the presence of gonads at the test termination. They quantified maturity as the ratio of the number of sexually mature mysids to the number of surviving mysids in each test replicate, and demonstrated that maturity was a sensitive endpoint for tests of 96-h and 7-day duration that were initiated with 8-day-old mysids. Others used the time required for mysids to reach sexual maturity as a test endpoint (e.g., Gentile et al. 1983, Lussier et al. 1985). Gentile et al. (1983) used the development of testes or the presence of eggs in the oviduct to determine when sexual maturity was reached in male and female mysids, respectively. They reported that exposure to high levels of mercury significantly lengthened the time required for mysids to reach maturity, which then was expressed as delay in the appearance of eggs in the brood pouch and the release of young.

193. Exposure of *H. azteca* to 17 α -ethinylestradiol showed several reproductive and morphological effects including: smaller gnathopods in females; skewed sex ratios favoring females; histological aberrations of the reproductive system, such as indications of hermaphroditism, disturbed maturation of germ cells; and disturbed spermatogenesis of the post F1 generation males (Vandenbergh et al. 2003). For both gnathopod growth and secondary sex characteristics, more pronounced effects were noted at the lower concentration (U-shaped curve), which may suggest a receptor-mediated response. It could be that 17 α -ethinylestradiol causes disturbance of the androgen gland through interaction with the AGH or hormone metabolizing enzymes with subsequent changes in AGH activity.

6.4.2.2 Time to First Brood Release

194. In uncontaminated systems, the length of time to the release of the first brood is primarily related to environmental temperature, with some influence by salinity and an interaction between the two factors (McKenney 1996). McKenney (1996) determined that the shortest time to release of the first brood for *A. bahia* was about 16 days at a temperature of 28°C and a salinity of 28‰. The time to release of the first brood increases with decreasing temperature and salinity. Because this parameter is measured as the number of days from hatching of the mysids used in the test until they release their first brood, it can represent the expression of more than one factor, including the length of time it takes a mysid to reach sexual maturity and the time required for eggs to develop in the brood pouch before being released. Most contaminant effects are likely to lengthen the time to release of the first brood. Lussier et al. (1985) found that several metals (mercury, zinc, nickel) significantly increased the time to first brood release, whereas others (e.g., cadmium, copper, silver) did not. Gentile et al. (1983) found that very high levels of mercury caused mysids to abort eggs that had been deposited in the brood pouch.

6.4.2.3 Egg Development Time

195. Egg development time is measured as the number of days between the first appearance of eggs in the brood pouch and the first release of juveniles. Gentile et al. (1983) found that mercury did not significantly affect the brood duration, although several other reproductive parameters were affected. Winkler and Greve (2002) calculated incubation time for *N. integer* as the difference between the date of laying eggs in the brood pouch and the date of release of the juveniles from the marsupium. Data were collected by removing females with fertilized eggs from culture jars and placing them in individual containers. Temperature affected the start of maturation. At 15°C, development was much shorter (1.5 months) than at 10°C (3 months). The reproduction rate of *N. integer* increased at higher temperatures because of declining incubation periods plus an increasing number of neonates released per brood (more juveniles released as the female gets larger). Also, at higher temperatures the number of released juveniles per brood was highly variable, most likely due to the successive oviposition of the females. Overall, *N. integer* had double the reproduction success of *P. flexuosus* because of the longer incubation times and release of fewer juveniles.

6.4.2.4 Brood Size (Fecundity)

196. Brood size can be measured as the number of eggs per brood (Khan et al. 1992) or as the number of young produced, expressed either as total young per female or as young produced per available female reproductive-day (Gentile et al. 1982, Lussier et al. 1985). The latter measure is used to normalize differences in the number of females available per test concentration (Gentile et al. 1982). The number of available female reproductive-days is calculated by multiplying the number mature females by the number of days survived. McKenney (1996) showed that the number of eggs in the first brood was related to salinity and temperature, and that the largest number of eggs was produced at a temperature of 25°C and 31‰ salinity. Because it is an important measure of reproductive success, any reduction in brood size can be interpreted as an indication of reproductive toxicity (Khan et al. 1992). However, brood size is also directly related to female size. Therefore, reduced fecundity in response to exposure to EDCs needs to be carefully evaluated to distinguish direct interruption of reproductive processes from a simple reduction in growth. Khan et al. (1992) also stated that the use of fecundity without supporting parameters to indicate reproductive impairment is not advisable, because fecundity is labor-intensive to determine, requires trained personnel, and ignores toxic effects on males. The most likely effect of contaminants is a reduction in fecundity (Hollister et al. 1980, Lussier et al. 1985), which in some cases is the only response to contaminant exposure (Lussier et al. 1999). Contaminant exposure can also result in the abortion of broods (Gentile et al. 1983). Lussier et al. (1999) reported, but could not explain, a seasonal difference in fecundity: more eggs were produced in the fall than in winter or summer. This phenomenon should be

considered when comparing tests conducted at different times of the year. Lussier et al. (1999) concluded that fecundity was nonetheless a sensitive and useful endpoint.

6.4.2.5 *Intersexuality and Sex Determination*

197. Exposure to EDCs can result in profound alterations in the reproductive characteristics of populations, expressed as physiological or morphological changes in individuals. For example, the most commonly reported phenomenon is a condition, pseudohermaphroditism, in which female mollusks develop male reproductive structures in response to exposure to tributyltin (LeBlanc et al. 1999). Among the Crustacea, cases in which individuals showed intersexuality have been reported for several different taxa (see references cited in LeBlanc et al. 1999). Mees et al. (1995) reported intersexuality in natural populations of *N. integer* collected in northern Europe. Ford et al. (2004a, 2004b) found intersexuality in males and females of *Echinogammarus marinus*, an estuarine amphipod. Ford et al. (2004a) found that the polluted sites had a higher incidence of intersexuality than reference sites in Scotland. When compared to “normal” individuals, intersex amphipods matured more slowly and showed reduced fertility and fecundity (Ford et al. 2004b). Reduced fitness was attributed to difficulties in mating resulting from the larger size of intersex individuals and the abnormal morphology associated with the condition. However, links between such phenomena in crustaceans and EDCs have not been established (LeBlanc et al. 1999, Ford et al. 2004a, 2004b). Regardless, Ford et al. (2004a) identified a distinctive morphometry associated with intersex males and suggested that it might be a useful biomarker of endocrine disruption. In some cases, abnormal sex ratios could be the result of EDC exposure, as has been seen in populations of copepods (Moore and Stevenson 1991, 1994). Studies of daphnid crustaceans have demonstrated that the population sex ratios found under good environmental conditions may be altered by juvenile hormone analogs (Olmstead and LeBlanc 2002, Tatarazako et al. (2003). Daphnids reproduce parthenogenetically, producing female young when environmental conditions are favorable, but produce male offspring when conditions become unfavorable. Five juvenile hormone analogs were shown to alter normal sex ratios in *D. magna* by stimulating the production of males (Tatarazako et al. 2003).

198. Relatively few field studies have been conducted addressing endocrine disruption to organisms found in estuaries, which have been shown to contain a variety of contaminants including sewage, chlorinated hydrocarbons, metals, and radioactive materials. Some of these contaminants have endocrine disruption potential (e.g., sewage effluents containing steroidal estrogens, DDT and its metabolites, and TBT) (Oberdörster and Cheek, 2000).

6.4.3 *Biochemical measures*

199. Five biochemical measures are explored below as possible endpoints for EDC-exposure tests: metabolic disruption, steroid metabolism, vitellogenin induction, and the levels of cytochrome P450 enzymes and of blood glucose. Care must be taken when interpreting results of bioenergetic biomarkers such as those mentioned here, because many abiotic as well as toxic stressors affect metabolic processes.

6.4.3.1 *Metabolic Disruption (O:N ratios)*

200. Many of the perturbations expressed in the parameters described in Sections 4.3.1 and 4.3.2 could be related to changes in energy pathways resulting from chronic exposure to contaminants. McKenney (1985, 1989) and McKenney et al. (1991) showed that exposure to contaminants caused increased respiration rates in juvenile mysids, often with as little as 24-h exposure. The increased general metabolic demands related to contaminant exposure reduced growth by decreasing the amount of energy available to produce new somatic tissues. Normally, energy allocations to metabolism, growth, and reproduction occur, so that if there are changes in allocations of energy this may be an indicator of stress attributable to a toxicant or an environmental change (e.g., temperature, oxygen levels) (Verslycke et al. 2004a). It is

important that bioenergetic endpoints include a baseline assessment of what is considered normal for mysids. Further, alterations to energy metabolism can be assessed by using tools like the cellular energy allocation assay. Changes in metabolic usage can also be monitored using oxygen and nitrogen ratios, protein and lipid content, or the carbon to nitrogen ratios.

201. One way of predicting effects of abiotic stress on energy metabolism is the cellular energy allocation (CEA) assay. Verslycke et al. (2003b) conducted a study using *Neomysis integer* exposed to varying concentrations of tributyltin chloride (10, 100, and 1000 ng TBT Cl/L). Effects were noted using the CEA assay such as less overall consumption of energy and lower respiration rates than control mysids. These results show that TBT interferes with the energy metabolism by disrupting energy production process. Overall there was a loss of protein, lipids, and sugars in the organisms exposed to higher concentrations of TBT versus the control. Later, Verslycke et al. (2004b) showed that exposure to sublethal doses of nonylphenol and methoprene significantly increased energy consumption, resulting in less energy available for reproduction and growth. Using the CEA assay to assess the energy budget quantitatively for a particular organism helps to elucidate potential modes of action for toxicants. Methods for CEA measurements include spectrophotometric measures of lipid, protein, and sugar (Verslycke and Janssen 2002). Additionally, electron transport activities were measured by using INT (p-iodonitrotetrazolium violet) as an electron acceptor mimic in the electron transport system, which can provide a measure of oxygen consumption (Owens and King 1975 as cited by Verslycke and Janssen 2002).

202. Increased metabolic demands caused by exposure to contaminants could also impair mysid growth and reproductive capability. Exposure to relatively high concentrations (0.072 $\mu\text{g/L}$, 0.1 $\mu\text{g/L}$) of the pesticide chlorpyrifos caused increased oxygen consumption rates and reduced egestion rates in *Neomysis integer* (Roast et al. 1999). These responses resulted in reduced energy available for growth (i.e., lower “scope for growth”). Young mysids typically use high-energy lipids to meet metabolic demands, but change to metabolize proteins as they mature, thereby leaving more lipid material available for the production of gametes (McKenney 1989). Gorokhova and Hansson (2000) found that a 6% decrease in the carbon:nitrogen ratios occurred as juvenile *Mysis mixta* matured to gravid adults and asserted that this was evidence that maturation and reproduction are fueled primarily by lipids. Increased metabolic demands caused by exposure to contaminants is met by greater lipid metabolism, which reduces the lipids available to meet reproductive needs (McKenney 1985, 1989). These changes in metabolic substrate usage can be measured by monitoring the oxygen:nitrogen (O:N) ratio of test organisms. The O:N ratio indicates the relationship between the amount of oxygen consumed by an organism to the amount of nitrogen excreted, and shows the relative role protein catabolism plays in the organism’s energy budget (Carr et al. 1985; McKenney 1985). *Americamysis bahia* showed a change toward lipid metabolism after only 4 days of exposure to high concentrations of cadmium (Carr et al. 1985). Mysids showed increased metabolic demands after four days of exposure to the herbicide thiobencarb (McKenney 1985) or five days of exposure to the defoliant DEF (McKenney et al. 1991). High O:N ratios occurred among maturing mysids exposed to the compound, indicating a shift to lipid metabolism that would have reduced the lipids available for gamete production. All three studies concluded that changes the O:N ratio measured among test mysids was a sensitive indicator that could provide for the relatively early detection of reproductive impacts by contaminants.

6.4.3.2 Steroid Metabolism

203. Although the role estrogens play in crustacean reproduction is not known (Baldwin et al. 1995), these steroids are important in other invertebrate groups. Because of the likelihood that crustaceans could be exposed to environmental estrogens such as dichlorodiphenyl trichloroethane (DDT), polychlorinated biphenyls (PCBs), and nonylphenols (nonionic surfactants), there is the potential for these compounds to disrupt steroid metabolism. Baldwin et al. (1995, 1997, 1998), studied the effects of three environmental

estrogens, diethylstilbestrol (DES), 4-nonylphenol (4NP), and nonylphenol polyethoxylate (NPPE), on the steroid metabolism of the freshwater daphnid, *Daphnia magna*. Their work focused on the disruption of the metabolic elimination of testosterone after short- and long-term exposure to the test compounds and sought to determine whether such an analysis could be used as an early indication of reproductive impairment. They measured differences in the glucose conjugation, sulfate conjugation, hydroxylated and reduced/dehydrogenated metabolites of ¹⁴C-labeled testosterone in daphnids exposed to sublethal concentrations of the test compounds. They found that the different compounds had different effects on testosterone metabolism. For example, DES increased glucose conjugation, but did not affect sulfate conjugation, whereas 4NP reduced both of these elimination processes. In their two earlier studies (Baldwin et al. 1995, 1997), Baldwin's group proposed that changes in testosterone metabolism could provide an early indication of potential reproductive toxicity after sublethal exposure to suspected EDCs. However, in their 1998 paper, Baldwin et al. studied NNPE, a nonionic surfactant that degrades to nonylphenol, and did not find significant disruption of steroid metabolism after short-term exposure. They did report some effects after chronic exposure and postulated that those could have resulted from the degradation of NPPE to NP. Therefore, they cautioned that use of short-term exposures as an early warning indicator might underestimate chronic effects resulting from bioaccumulation and bioactivation of the test compounds.

204. Verslycke et al. (2002) studied testosterone metabolism in *Neomysis integer*. Significantly, they detected endogenously-produced testosterone in male and female mysids. They also found an anabolic steroid, β -boldenone—the first known occurrence of the compound in an invertebrate—in mysids exposed to testosterone added to the test medium, although the metabolic pathway by which it is formed was not discovered. A vertebrate estrogen, 17 α -estradiol was not detected. Verslycke et al. (2002) showed that testosterone metabolism in mysids involves phase I (oxido-reduced/hydroxylated) and phase II (conjugated) derivatives. The study proposed that changes in invertebrate steroid metabolism might be used to indicate exposure to EDCs. To evaluate this idea, Verslycke et al. (2003b) examined the effect of exposure to tributyltin on the elimination of testosterone by the mysids. The authors summarized the effect of TBT exposure on testosterone elimination by calculating the metabolic androgenization ratio—the ratio of the eliminated oxido-reduced products to the polar products (hydroxylated plus conjugated). TBT exposure changed testosterone metabolism by inducing reductase activity at low, but not high, concentrations and reducing sulfate conjugation, although the mechanisms by which these occurred were not identified. To further refine the idea that changes in testosterone metabolism could be used to indicate exposure to an EDC, Verslycke et al. (2004b) subjected individuals of *N. integer* to seven compounds (testosterone, flutamide, 17 α -ethinestradiol, precocene, nonylphenol, fenoxycarb, methoprene) suspected of having endocrine disruption properties. All seven were very toxic to *N. integer* with 96-h LC₅₀ values ranging from 0.32 mg/L (methoprene) to 1.95 mg/L (testosterone). The effects of sublethal doses of nonylphenol and methoprene on testosterone metabolism were investigated and found to be significant. Methoprene and nonylphenol affected phase I testosterone metabolism in a manner similar to that of TBT (induced reductase activity at low concentrations of the chemical). The two compounds had opposite effects on phase II metabolism. Glycosylation was increased at a high (100 μ g/L) dose of nonylphenol, but was significantly reduced at the same dose of methoprene. The metabolic androgenization ratio showed a dose-dependant increase after exposure to methoprene, but after exposure to nonylphenol was significantly increased only at the lowest dose tested, 10 ng/L.

6.4.3.3 Vitellogenin

205. Depledge (unpublished, cited in Depledge and Billingham 1999) found that exposure to 4-nonylphenol induced the production of vitellogenin in decapods. Vitellogenin production is most likely controlled by, the hormones primarily involved in molting, although in some crustacean groups this is likely not the case (Subramoniam 2000). The molt-inhibiting hormone and 20-hydroxyecdysone (20HE) are involved in the production and regulation of vitellins and there are several feedback loops (positive and

negative) involved. To date, because of the complexity of vitellogenesis, it has not been used as a biomarker of endocrine disruption (Oberdörster and Cheek 2001). Little research has been conducted on vitellin expression in crustaceans, possibly due to the lack of antibodies, which have been shown to have cross-reactivity with other species. Most often polyclonal antibodies are used and available for crustaceans with the drawback that these antibodies often cross-react with other nonvitellin proteins. One such study, using grass shrimp exposed to pyrene, showed up-regulation of vitellin, and when the females were transferred to clean seawater, there was an increase in embryo mortality, suggesting that the vitellin can transport lipophilic contaminants to developing embryos.

206. As pointed out by Verslycke et al. (2004a), to understand the potential effects of xenobiotics on reproduction, there must be accurate measures of vitellogenin and vitellin in crustaceans. A few studies had been done to either develop an ELISA or to characterize and purify vitellin. Additional studies need to be conducted to evaluate the utility of such measures.

6.4.3.4 *Cytochrome P450 Enzymes*

207. Cytochrome P450 enzymes (CYPs) are commonly occurring proteins that are involved in the metabolism (i.e., detoxification) of many exogenous and endogenous compounds (Snyder 2000, Snyder and Mulder 2001, Verslycke et al. 2004a). Snyder and Mulder (2001) measured CYP45 levels, a family of P450 proteins found in the lobster *Homarus americanus*, and thought they must be involved in the molting cycle, in response to exposure to the pesticide heptachlor, a known EDC. They found that peak levels of ecdysteroid hormones, and accordingly, molting, occurred later in heptachlor-exposed larvae than in control larvae, indicating that heptachlor disrupts steroid molting hormone metabolism. They also found CYP45 levels in lobster larvae exposed to heptachlor on Days 1, 2, or 3 after hatching to be 15 times higher than they were in those exposed to control solutions. Levels of CYP45 typically peaked 1 to 2 days after exposure, then decreased. Snyder and Mulder suggested that it could be a useful early biomarker of exposure to EDCs, because it showed a dramatic and rapid increase in levels after exposure to heptachlor. In their testosterone metabolism study, Verslycke et al. (2002) showed that *Neomysis integer* has many P450 enzymes that comprise its complex steroid hydroxylation system. They suggested that changes in P450 activity could be used as a biomarker indicating exposure of this mysid to EDCs.

208. It is known that most P450 activities occur in the hepatopancreas, but other tissues, such as the gills, stomach, intestines, and antennal glands, have demonstrated some P450-like roles. Oxidative metabolism, most notably of PAHs, has been demonstrated in many crustacean species. A detailed review of studies conducted examining the metabolic changes (mostly caused by benzo(a)pyrene) in crustaceans can be found in James and Boyle (1998). This paper also offers suggestions of P450 involvement and or metabolism relative to drugs, steroids, and pesticides. Several studies have shown the presence of testosterone and progesterone in gonadal tissues and hemolymph. The source of these steroids is not well known but it is likely that several cytochrome P450-dependent steps are involved. Future directions should include genomic information on the crustacean P450 system, the definitive identification of an aH receptor, and a best understanding of the regulation of the P450 system, particularly for steroid synthesis.

6.4.3.5 *Blood Glucose Levels*

209. Levels of glucose in crustacean blood is regulated by a hormone, crustacean hyperglycemic hormone (CHH) that is produced in the sinus gland (Fingerman et al. 1998). Release of CHH increases blood glucose levels. Some exogenous compounds have been shown to affect the levels of glucose in the bloodstreams of several crustacean taxa, probably be stimulating (e.g., naphthalene) or inhibiting (e.g., cadmium) CHH synthesis (Fingerman et al. 1998). Measurement of changes in blood glucose levels in mysids exposed to potential EDCs could be indicative of hormonal perturbation other than that associated directly with reproduction or molting. Hyperglycemia is a common response to environmental or toxicant

interactions, and changes in blood glucose levels in response to presence of EDCs may be an indication of interference with hormonal activities (Verslycke et al. 2004a). However, many abiotic and toxic responses affect energy metabolism, and researchers must be able to make distinctions between natural variations in hormone levels and variations that are caused by EDCs.

210. Methyl farnesoate (MF) is the unepoxidated form of the insect juvenile hormone (JHIII) and is produced in the mandibular organ. Recent studies have suggested that MF functions in a similar manner in crustaceans as JH does in insects (Laufer and Biggers 2001). MF is involved in reproductive maturation and in morphological changes most notable from late juvenile stages to adult. Various investigators have conducted eyestalk ablation studies in which crustaceans such as shrimp, crayfish, and fiddler crabs were stimulated to reach ovarian maturation. JH analogs such as methoprene have inhibited early larval and post larval development in the shrimp *Palaemonetes pugio*; they actually enhanced premetamorphic stage development.

7.0 RESPONSE TO ECDYSTEROID AGONISTS AND ANTAGONISTS

211. Concern has often been expressed in recent years about the disruption of endocrine systems in aquatic organisms by the action of organic and inorganic contaminants (e.g., Snyder and Mulder 2001; Depledge and Billingham 1999; Fingerman 1997). In a review by Hutchinson et al. (1999a), it was suggested that based on estimated figures, the concentration of EDCs expected to be introduced to the United States' aquatic environment could be as high as 2.16 ng/L for 17 α -ethinylestradiol-derived contraceptives, and 41.5 ng/L for conjugated estrogens used in hormone-replacement therapy. Although there has been considerable research conducted on the health of fish exposed to EDCs, there is little information available for crustaceans (Baldwin et al. 2001, Hutchinson et al. 1999a, Hutchinson et al. 1999b).

212. Developmental abnormalities and toxicity to daphnid embryos were noted (at levels far below concentrations causing toxicity to maternal organisms) when the maternal organisms were continuously exposed and also when the embryos were collected from unexposed parents and exposed directly to testosterone. These developmental abnormalities and delays in molt frequency of neonates were mitigated when the daphnids were co-exposed to 20-hydroxyecdysone. These findings suggest that testosterone may function as an anti-ecdysteroid in crustaceans (Mu and LeBlanc 2002a). This study suggests that testosterone was able to block the activity of 20-hydroxyecdysone when both steroids were provided. Ecdysteroids for the early embryo stage originate from the mother and are transferred to the egg. Ecdysteroids present in the late embryo are in part synthesized from the embryo. Thus, when the embryo is exposed to an anti-ecdysteroid antagonist it could affect both pools (maternal and embryo) of ecdysteroids. This could be manifested by early- and late-stage abnormalities. When direct exposure of embryos is conducted, it would have no impact on the maternal pool of ecdysteroid resident in the embryo, but it would affect production of ecdysteroids within the embryo itself, most likely noted by late-stage developmental abnormalities.

213. In the environment most chemicals are present as mixtures, yet little research has been conducted to evaluate the potential endocrine disruption of complex mixtures. Toward that end, the toxicity of a chemical mixture of fenarimol and testosterone was evaluated using *Daphnia magna* to ascertain if combined exposure would result in greater than additive toxicity. By itself, fenarimol causes late-stage developmental abnormalities in embryos while testosterone interferes with early and late-stage embryo abnormalities (Mu and LeBlanc 2004). Fenarimol is a known ecdysteroid synthesis inhibitor of endogenous hormones; when combined with testosterone, a known ecdysteroid antagonist, fenarimol effectively inhibited hormone synthesis, paving the way for testosterone to bind to ecdysteroid receptors. By exposing only embryos to fenarimol, this would result in perturbations in late embryo development since maternal ecdysteroids would be present and active, whereas exposure of testosterone to isolated embryos would cause both early- and late-stage developmental abnormalities. Results showed that fenarimol increased the toxicity of testosterone, while testosterone had no effect on the toxicity of fenarimol. Additionally, a model was used to predict combined effects using algorithms. The predictive model was very effective in estimating the joint toxicity of these compounds for the independent action and synergy of both compounds. Further studies are needed to evaluate the role of complex mixtures on crustacean endocrine systems.

214. The endocrine and reproductive effects of EDCs mimic the effects of natural hormones, antagonize the effects of hormones, alter the pattern of synthesis and metabolism of hormones, and modify hormone receptor levels (Depledge and Billingham 1999). The ability of some environmental contaminants to bind to steroid hormone receptors as agonists or antagonists in a recognized mechanism of toxicity to endocrine-related processes has been documented (LeBlanc and McLachlan 1999).

215. The literature is vague with respect to gender differences from exposure to ecdysteroids. Cuzin-Roudy and Saleuddin (1989) discussed possible differences in effects to male and female mysids, *Siriella armata*. This study showed that secondary vitellogenesis starts at the beginning of the molt cycle for this organism, when ecdysteroid levels are low. There is a striking difference between males and females at this point: in females, ecdysteroid levels were 10 times higher than those in males, but the response of the epidermis for molt preparation was the same. Females also had much higher levels of 20-E, ecdysone, and high polarity products, which are probably linked to the storage of ecdysteroids in oocytes during secondary vitellogenesis. Embryonic and post-embryonic development occurs in the marsupium of the females. Juveniles are released shortly before ecdysis, after which the adult female lays a new batch of eggs in the marsupium. A secondary vitellogenic cycle starts for a new batch of oocytes on the second day of the female molt cycle. Secondary vitellogenesis is strictly linked to the molt cycle. During development, gonads and gonoducts differentiate before the appearance of secondary sexual characteristics (Cuzin-Roudy and Saleuddin 1989).

216. Crustaceans are in general fast-growing and slow-breeding organisms. Integration between molting and reproduction is a physiological necessity in females. Ecdysteroid, the chief hormone in molting, is thought to be involved with control of female reproductive activities. However, this is controversial. Investigations using amphipods have shown that levels of vitellogenin fluctuate with hemolymph ecdysteroid levels (see for example Cuzin-Roudy and Saleuddin 1989, Depledge and Billingham 1999).

217. In crustacean females, sequestered ecdysteroids may be passed on to the eggs for possible elimination and to function as morphogenetic hormones partaking in the control of embryogenesis and early development. The ovary in many crustaceans accumulates ecdysteroid for possible use during embryogenesis (Subramoniam 2000). Molting and reproduction are more evident in the female, because vitellogenesis is the central event of the female reproductive cycle along with secretion of a new cuticle during molting. Hormones play a role in the nutritive supply for molting and vitellogenesis. The ovaries eliminate ecdysteroids by forming ecdysonic acid as a necessary way to eliminate ecdysteroid in the eggs and embryos. They also form conjugates as a means of elimination. In embryos, there are concentrations of the three ecdysteroids—ecdysone, 20-E, and PoA—and their conjugates. There are fluctuations in embryonic ecdysteroids, as evidenced by one shrimp species, *Sicyonia ingentis*, in which the eggs after spawning contain low levels of ecdysteroid. The levels then rise through development, probably by the synthesis of this hormone by the embryo's Y-organ. The endogenous accumulation of ecdysteroid within the ovary is also known to function in the induction of meiotic maturation of the oocyte (Subramoniam 2000).

218. In one experiment, Subramoniam (2000) removed eggs from the pleopods of the freshwater prawn, *Macrobrachium nobilii*. The release of eggs quickened the next molting and reproductive cycle. In another experiment, Subramoniam (2000) found that although the ovarian cycle begins during the intermolt stage, vitellogenesis (serum levels) progresses into the next premolt stage. Premolt starts with the release of the larvae, and the next spawning occurs after ecdysis. Among penaeid shrimp, free spawning occurs during the premolt stage, followed by ecdysis (Subramoniam 2000). There are few *in vitro* studies available that focus on specific mechanisms involved in disruption in arthropods. The ecdysone receptor is in the same gene family as the thyroid receptor found in vertebrates.

219. In another experiment conducted by Bodar et al. (1990), daphnids were exposed to varying concentrations of cadmium and separately to exogenous ecdysone and 20-hydroxyecdysone to monitor any changes to the molt or reproductive cycle and to evaluate the role of ecdysteroids relative to molting and reproduction. The study showed dose-dependent effects on molting and reproduction for the ecdysone and cadmium exposures. The effects of the higher ecdysteroid concentrations included unsuccessful exuviations, incomplete molting, and eventually death such that animals died before the age of potential reproduction was reached. The daphnids did not molt after treatment to ecdysteroids, and it was speculated that they spent a disproportionate amount of energy on molting which negatively impacted the reproductive physiology.

220. For the cadmium exposures, a dose-dependent effect of cadmium on ecdysteroid titers was observed. At a cadmium concentration of 5 $\mu\text{g/L}$ there was an increase in levels above control (~210 pg ecdysone eq/mg dry weight) after 2-day exposure; these high levels declined after 8 days to levels approaching the control. As the cadmium concentration increased to 20 $\mu\text{g/L}$, there was a linear increase in ecdysone concentrations to around 750 pg ecdysone eq/mg, which is three times higher than controls. A pronounced decline in growth occurred under cadmium exposure. Also, a stimulatory effect on steroid hormones was seen such that increasing cadmium levels corresponded with increasing hormone titers. This stimulatory effect of cadmium on ecdysteroids has been observed for other organisms. Two theories are put forth: either the cadmium caused increased ecdysteroid levels which led to molt and reproductive impairment, or the cadmium interfered with the metal regulatory system through the metallothioneins and metalloenzymes that are involved in the molt cycle.

221. Incubation of ecdysteroid synthesis tissues *in vitro* is a method to detect endocrine modulators on molting hormone synthesis. This method was investigated as a potential biomarker using the midge *Chironomus riparius*. This method can be used to determine if a particular chemical causes endocrine disruption at the suborganismal level. Detailed methods for conducting ecdysteroid biosynthesis and subsequent measurement using radioimmunoassay techniques is described in Hahn and Schulz (2002). Male and female midges responded quite differently to this technique such that ecdysteroidogenic activity was significantly increased above controls for the males, while the opposite trend was found for the females. Also, exposed males developed faster than controls, whereas the treated females showed slower development than the controls. Even further, there is speculation that ecdysteroid metabolism is regulated by different processes in males and female midges during the fourth stage of larval development.

222. Block et al. (2003) used two crustaceans—a copepod (*Amphiascus tenuiremis*) and an amphipod (*Leptocheirus plumulosus*)—to determine ecdysteroid concentrations at different life stages. They also used a method known as fluorescence-based enzyme immunoassay (EIA) to quantify and compare ecdysteroid titers in such small organisms. Detailed methods for conducting the assay, including tissue collection and extraction of small sample volumes, is described. The overall synthesis, regulation, and metabolism of ecdysteroids used within and across species is most likely associated with variations in growth (molting), mating, and life cycles. Therefore, it is important to have a precise measure of ecdysteroid levels capable of detection at the femtomolar (10^{-15} molar) level.

223. In summary, detailed studies of crustacean response to ecdysteroids are lacking. Future studies that address sequence determination of vitellogenic genes and their hormonal activity could provide interesting insight into the vitellogenic process in this taxonomic group. A genomic and nongenomic effect of ecdysteroid on ovarian maturation is a potential area of work. Synergistic and antagonistic actions of the X-organ sinus-molt and gonad-inhibiting neuropeptides, and the mandibular organ control over molting and reproduction are other areas requiring further study as a basis for use of crustaceans for EDC testing in the future.

8.0 ANDROGENIC AND ESTROGENIC RESPONSES

8.1 ANDROGENIC RESPONSES

224. Vertebrate-type steroidal androgens have been measured in some crustaceans, but androgen receptors have not been documented. Presently, the androgenic gland has been identified only for malacostracans (Block et al. 2003). Steroidal androgens can function directly as hormones in ways that do not require receptors, or they can be present as inactive components of steroid metabolic pathways (LeBlanc and McLachlan 1999). There is currently no published research that evaluates the androgenic hormones and their effect specifically on mysids. Administration of testosterone to shrimp has resulted in hypertrophy and hyperplasia of the androgenic gland. The androgenic gland is associated with the testis, and is responsible for the secretion of the androgenic hormone. This hormone is nonsteroidal and is responsible for masculinization. Testosterone administered to shrimp and crab results in the increase in testis size and in the conversion from ovaries to testes in females (LeBlanc and McLachlan 1999). Experiments using *Daphnia magna* showed that testosterone, acting as an antagonist to 20-hydroxyecdysone, caused toxicity to neonates by interfering with the control of molting by ecdysteroids (Mu and LeBlanc 2002a). These studies, and the recent discovery of endogenous testosterone in *Neomysis integer* (Verslycke et al. 2002), suggested that studies designed to measure androgenic effects in mysids could be conducted. To date, an androgen receptor has not been identified nor cloned in crustaceans; research should be directed at identification and characterization of this receptor to aid in determining the usefulness of sex steroids as an evaluation tool for crustaceans (Verslycke et al. 2004a).

8.1.1 Endpoint Sensitivity

225. Vertebrate-type steroid hormones are found in the ovaries, testis, mandibular organ, and hemolymph of crustaceans. Many of these steroid hormones exhibit fluctuations during gonadal development, suggesting a role in reproduction of crustaceans.

226. The identification of physiological targets of EDC in invertebrates is the approach taken by LeBlanc and McLachlan (1999). One example is that diethylstilbestrol and endosulfan have been shown to inhibit molting in immature daphnids, but to have no effect on the mature animals' fecundity. These effects may indicate that chemicals that are estrogenic to vertebrates could affect molting and reproduction in crustaceans, interfering with the proper function of the ecdysone receptor. In a study designed to examine antiandrogens, *Daphnia magna* was exposed to the compound cyproterone acetate, to determine whether it interferes with the androgen receptor as it does with vertebrates (LeBlanc and McLachlan (1999). The results showed an impairment to growth. The exposed organisms were smaller, and there was a reduction in number of offspring. The latter was most likely due to the smaller size of the organisms, which would not have been able to accommodate a more normal number of brood in the pouch. The effects of steroidal androgens and chemicals that cause metabolic androgenization are consistent with interference to the delivery or packaging of nutrients into the developing eggs. Ecdysteroids, juvenoids, progestogen, and crustacean androgens have all been shown to influence vitellogenin or lipid production in arthropods. Androgens may interfere with one or more of the hormonally regulated processes that provide nutrients to embryos.

227. Baldwin (1997, 1998) conducted a series of experiments using the daphnid. During one experiment, the daphnid was exposed to 4NP, which resulted in changes in rates of elimination of

testosterone and a corresponding decrease in glucose-conjugated testosterone, and an increase in the rate of production of various androgenic derivatives of testosterone. This is called metabolic androgenization, which is found to reduce fecundity of exposed daphnids associated with developmental abnormalities and high mortality of offspring. Results from a separate experiment with exposure to NP revealed no significant evidence of changes in steroid elimination processes, except at the highest concentration, which reduced elimination of glucose- and sulfate-conjugates and increased elimination of oxido-reduced derivatives. Effects were seen at sublethal levels for 4NP and approaching acute levels for nonylphenol. It has not yet been determined whether there is an androgen receptor in crustaceans; therefore, more studies are needed to determine the functional role of steroidal androgens.

228. It is possible that endogenous androgens may be precursors to other hormones, and that large doses of exogenously added androgens could elicit activity through other receptors. In crustaceans, testosterone is converted to androstenedione at various rates (LeBlanc and McLachlan 1999). Future studies may reveal that the conversion is affected by age, reproductive state, or photoperiod. It is possible that alteration in testosterone metabolism could serve as a biomarker, because effects are observed at concentrations less than those eliciting reproductive response.

229. Verslycke et al. (2003b) examined the elimination rates of testosterone by monitoring a specific set of metabolites (polar hydroxylated, nonpolar oxido-reduced, and glucose- and sulfate-conjugated). Various theories surround imposex of neogastropods exposed to TBT: inhibition of the cytochrome P450 system; blocking phase II sulfate conjugation; and interference directly with neurohormonal system, leading to changes in steroid titers. Further work is needed to confirm which mechanism(s) is responsible.

8.1.2 Gender Differences

230. Currently, there is no documented research that discerns gender differences in mysids as a result of androgenic-type hormone response. Detailed mechanistic and anatomical studies would need to be conducted on mysids to ascertain whether differences in gender relative to EDCs can be observed.

231. In the 1980s, the condition of imposex (the imposition of male sex organs including a penis and vas deferens) was observed with increasing frequency on marine gastropods exposed to tributyltin (TBT) (Depledge and Billingham 1999). The mode of action of TBT giving rise to imposex is currently under investigation. Female snails exposed to TBT have elevated testosterone in the hemolymph, and injections of TBT into females induced penis formation (Depledge and Billingham 1999). Lee (1991) thought that many of the observed effects in mollusks are related to enzymes involved in TBT metabolism. Inhibition of a cytochrome P450-dependent aromatase (which normally converts 17α -estradiol to testosterone) could result in the accumulation of testosterone, which would otherwise be metabolized.

232. Studies of the shrimp, *Palaemon serratus*, showed that eyestalk ablation resulted in rapid maturation of the ovaries (reviewed by Fingerman et al. 1998). It was later shown that this effect is caused by the sinus gland containing a gonad-inhibiting hormone (GIH). This system is present in male crustaceans as well, and eyestalk ablation to induce gonadal maturation is a common practice on shrimp farms worldwide (Fingerman et al. 1998). The presence of gonad-stimulating hormones (GSH) was demonstrated in decapod crustaceans. In female crustaceans, the GIH and GSH acted directly on ovaries, which then secreted the ovarian hormone. Ovaries are a source of ovarian hormone, which induces the development of secondary female sexual characteristics. In male crustaceans, GIH and GSH acted on the androgenic gland. Two experiments were conducted to determine the role of the androgen gland using *Macrobrachium rosenbergii*. When the androgen gland was removed, the male became feminized, and when the androgen gland was implanted into a female, the female became masculinized (Fingerman et al. 1998).

233. In their review, Fingerman et al. (1998) reported that parasitism of crustaceans by rhizocephalans induced castration. The castration of the males often involved additional impairment to testicular function by modification of the secondary sexual characteristics, causing the males to take on female appearance. For example, the narrow male abdomen of crabs became wider, resembling that of a female. Several authors, such as Fingerman et al. (1998) have reported that in male shore crabs, *Carcinus maenas*, spermatogenesis occurred nonetheless in the testes of specimens found with feminized abdomens.

8.2 ESTROGENIC RESPONSES

234. Billinghamurst et al. (1998, 2000, 2001) examined the effects of two estrogens, 4-*n*-nonylphenol and 17 β -estradiol on larval settlement and the production of a larval storage protein (cypris major protein, CMP) in *Balanus amphitrite*. Cyprids use CMP during settlement and the early post-settlement development. Because CMP is structurally related to vitellin, which is analogous to vitellogenin, it can be used as a biomarker of estrogen exposure in lower vertebrates. The expectation in these studies was that cyprid settlement might be affected by the stimulation of CMP synthesis after larval exposure to environmental estrogens. The results of the 1998 study, however, showed reduced settlement after exposure to both estrogens, but that the cause was not related to endocrine disruption. The second study (Billinghurst et al. 2000) measured levels of CMP and found that they were elevated after exposure of nauplii to low levels of the estrogens. The third study (2001) measured effects of 4-*n*-nonylphenol and 17 β -estradiol to larvae of *Eliminius modestus* (nauplii and cyprids). Specifically a disruption of the timing of larval development was noted, but this disruption was not consistent for different trials of this same experiment. This is in contrast to the 1998 study, but the studies were carried out at different times of year and with species that have different reproductive cycles. The variable response of different species to the same chemical reinforces the concept that development of larval crustaceans is subject to precise mechanisms and exposure to contaminants may depend on season and species. Further, this study showed that varying the timing of exposure of 4-*n*-nonylphenol and 17 β -estradiol or the duration was critical. For example, organisms exposed for 12 months were significantly smaller than control organisms. As reported for other studies, Billinghamurst et al. (2001) found that intermediate concentrations of NP are more disruptive than higher or lower concentrations.

235. Hutchinson et al. (1999a, 1999b) found that exposure to several steroids had no effect on the survival and development of copepod (*Tisbe battagliai*) nauplii and cautioned against extending the reported effects of steroid exposure in some species of crustaceans to the group as a whole. At about the same time, Bechmann (1999) showed that high levels (>62 $\mu\text{g/L}$) of nonylphenol were acutely toxic to *T. battagliai*, but that exposure to a low level (31 $\mu\text{g/L}$) did not affect any of the measured life-table parameters (survival, sex ratio, fecundity) measured. Breitholtz and Bengtsson (2001) did not find evidence of endocrine disruption in the harpacticoid copepod *Nitocra spinipes* after exposure to the estrogens 17 β -estradiol, 17 α -ethinylestradiol, and diethylstilbestrol. This contrasts with another study that exposed *Hyalella azteca* to 17 α -ethinylestradiol at concentrations of 0.1 $\mu\text{g/L}$ to 10 $\mu\text{g/L}$ in a multigenerational experiment (Vandenbergh et al. 2003). Results showed that F1 males exposed from gametogenesis to adulthood developed significantly smaller second gnathopods; this response generated a U-shaped dose response curve suggesting a receptor-mediated response because effects were found at lower concentrations but masked at higher concentrations. Post F1-generation males exhibited histological aberrations of the reproductive tract (i.e., hermaphroditism, disturbed maturation of germ cells, and disturbed spermatogenesis); again these responses were more pronounced at the lower concentrations, suggesting a receptor-mediated response. Also noted, but not statistically significant, was that the populations exposed to 17 α -ethinylestradiol for more than two generations tended to favor females.

236. Additional experiments conducted by Oberdörster et al. (2000) using *Palaemonetes pugio* in a 6-week pyrene exposure showed a significant increase in vitellin at 63 ppb and a significant increase in embryo mortality at 63 ppb. The increase in VTN could be a countermeasure against lipophilic

compounds such as pyrene, because vitellins may be able to bind lipophilic compounds and transfer them to developing embryos.

237. In the shrimp, *Penaeus monodon*, 17α -estradiol and progesterone in free and conjugated forms increase in the ovary during vitellogenesis (Fairs et al. 1990). Metabolic precursors such as pregnenolone and dehydroepiandrosterone also increase and show a peak during the major vitellogenic stages, suggesting a pathway in crustaceans that is similar to that in vertebrates. Fairs et al. (1990) also reported that 17α -estradiol and progesterone levels in the hemolymph showed fluctuations resembling that of serum vitellogenin levels during ovarian maturation. Estrogen could possibly control the stimulation of yolk synthesis, whereas the progesterone could control the prophasic meiotic maturation, causing germinal vesicle breakdown in the post vitellogenic oocytes. Exogenous injections of steroidal hormones induced vitellogenesis in the prawn, *P. japonicus*. In a study of a marine shrimp, *P. semisulcatus*, it has been shown that both the vitellogenin synthesis in the hepatopancreas and vitellin synthesis on the oocytes are coded by one gene (Subramoniam 2000).

238. Female crustaceans synthesize and secrete the protein vitellogenin into the hemolymph at the onset of oogenesis. There are a limited number of enzyme-linked immunosorbent assays (ELISA) that have been developed for measurement of vitellogenin or vitellin (VTN) in crustaceans but all are designed for detection in larger decapod species. A new method has been proposed by Volz and Chandler (2004) for measurement of VTN in small microquantities in a sediment-dwelling copepod, *Amphiascus tenuiremis*. This ELISA uses VTN-specific polyclonal antibodies from *Leptocheirus plumulosus*, which show specificity toward female copepod proteins. Quantities of purified VTN, used as a standard, were collected from grass shrimp eggs because it can be collected in relatively large quantities and it reacts well with anti-VTN antibodies from *L. plumulosus*. The results using this ELISA showed significant discrimination between gravid females and male samples. The final working range for the ELISA was from 31.25 to 1000 ng/mL with intra-assay and interassay CVs of 3.9 and 16.8%. Further, the detection limit was 2 ng/mL and the ability to quantify VTN in small numbers (four or more) copepods makes this ELISA a promising tool for further research for monitoring endocrine activity of toxicants to copepods and other crustaceans.

239. Studies (Pounds et al. 2002) to determine mode of action of several selected natural and synthetic steroids and xenoestrogens were examined using a combination of the *Tisbe battagliai* life-cycle test and also the B_{II} haemocyte line of *Drosophila melanogaster*, which is a screening assay to examine the agonist and antagonist effect of compounds to the cell line. One steroid tested, 20 HE, demonstrated reproductive effects on *Tisbe battagliai* at 26.9 μ g/L and also demonstrated agonist activity to the ecdysteroid receptor, suggesting that the *in vivo* response was mediated via the receptor. The other steroids tested showed no response to either assay, indicating that even at high concentrations the synthetic and natural vertebrate steroids do not interact with the ecdysteroid receptor. Receptor-binding assays provide information relative to specific interaction of a compound to the endocrine system but do not give predictive information about how this compound will affect the whole organism.

9.0 RESPONSE TO OTHER HORMONAL DISTURBANCES

240. In their review paper, Fingerman et al. (1998) described other hormonal responses and disturbances in crustaceans, such as color-changing hormones, retinal pigment hormones, pericardial hormones, and blood glucose hormones. Each of these will be briefly described below, relative to crustaceans in general. Specific hormone disturbances to mysids as well as endpoint sensitivity and gender differences await further study.

241. The sinus gland is the storage and release site for color-change hormones, among others. The sinus gland is located proximal to the eye and lies next to the large hemolymph sinus (Fingerman 1997). For sessile crustaceans, the sinus gland is located in the head close to the optic centers. Investigators believe that 90% of axonal terminals that compose the sinus gland belong to neurons whose cell bodies lie in the medulla terminalis X-organ. Therefore, the medulla terminalis X-organ sinus-gland complex is similar to the vertebrate hypothalamo-neurohypophyseal complex (Fingerman 1997).

242. Color change is affected by cells called chromatophores, which are located in the integument. They are responsible for color change through their dispersion and aggregation. In an early experiment, the hemolymph of a dark prawn specimen was transferred to a pale one (Fingerman 1997). When this organism was kept on a white background, it turned dark. The researcher then cut through the exoskeleton to sever any peripheral nerves that should innervate the chromatophores to determine whether color change was related to the endocrine system or to the nervous system. The incision had no effect on color change. Histological examination failed to show any innervation of the chromatophores, which led to the conclusion that color changes of this prawn are due to hemolymph-borne pigment concentrating substances.

243. For a variety of decapod crustaceans, chromatophores that cause integumentary color changes are controlled by antagonistically acting pigment-dispersing and pigment-concentrating neurohormones. For example, in the fiddler crab *Uca pugilator*, the neurohormone, 5-HT-serotonin triggers the release of red pigment-dispersing hormone, but has no effect on the black chromatophores (Fingerman 1997). The black chromatophores are triggered by norepinephrine, which releases a black pigment-dispersing hormone (BPDH). Studies have shown that the eyestalks of *U. pugilator* contained four times as much BPDH as did the control after exposure to naphthalene, due to naphthalene's inhibition of norepinephrine release. The opposite mechanism was observed for cadmium (Fingerman 1997). The eyestalks of control organisms contained three times more BPDH than did the cadmium-exposed crabs, which indicated that cadmium inhibited the synthesis of BPDH.

244. Retinal tissue contains pigments that control the amount of light striking the rhabdom (the photosensitive part of each ommatidium that compose the compound eye) through changes in position. Three types of retinal pigments have been categorized: the distal, proximal, and reflecting (Fingerman 1997). Migration of the distal pigment is controlled by the light-adapting hormone (LAH) and the dark-adapting hormone (DAH). Most studies of retinal pigments use the distal pigment, because techniques for its use are noninvasive. Several studies were conducted using *Palaemonetes vulgaris*. In one experiment conducted by Fingerman in 1959, this species was kept under constant illumination and then injected with extracts of eyestalks or sinus glands (described in Fingerman 1997). Because there is an initial light-adapted response followed by a dark-adapted response, Fingerman was able to induce a dark-adapting

response from an organism kept under constant illumination, suggesting that eyestalks contain both LAH and DAH.

245. The sinus gland contains the source of CHH, which causes elevation of blood glucose levels for crustaceans. CHH is similar in structure to MIH. It has been determined that MIH and CHH also show similar activity (Fingerman 1997). Future studies should address the specific roles of these hormones in mysids and other crustaceans. Exposure of the freshwater prawn *Macrobrachium kistnensis*, and several species of crabs, to cadmium caused hyperglycemia. Similarly, exposure of *U. pugilator* to naphthalene caused hyperglycemia, although the mode of action is apparently different for the two compounds. Cadmium inhibits CHH synthesis, whereas naphthalene stimulates CHH synthesis; 5-HT apparently triggers release of CHH (Fingerman 1997).

246. Pericardial organs lie in the venous sinus that surrounds the heart, and the axon terminals could be part of the neuroendocrine system that releases hormones affecting the heart. Experiments showed that the hearts of three species, *Cancer pagurus*, *Homarus vulgaris*, and *Squilla mantis* responded to pericardial organ extracts with increase in both frequency and amplitude of the heart beat. Efforts to identify substances in the pericardial organs have revealed 5-HT, dopamine, and octopamine (Fingerman 1997).

10.0 CANDIDATE PROTOCOLS FOR AQUATIC ARTHROPODS LIFE CYCLE AND TWO-GENERATION TOXICITY TESTS

247. Chronic testing protocols for freshwater and estuarine or marine arthropod species have been developed by several regulatory agencies. Examples of protocols developed by national regulatory agencies and by OECD on Daphnids, on Chironomids and other sediment dwelling invertebrates, on Mysids and on Copepods are highlighted below.

10.1 PROTOCOLS USING DAPHNID SPECIES

248. EPA (2002a) developed a chronic protocol for use in the National Pollutant Discharge Elimination System (NPDES) Permits Program to identify effluents and receiving waters containing toxic materials in chronically toxic concentrations. The method includes chronic testing procedures for a daphnid (*Ceriodaphnia dubia*) that, although not designed specifically for EDC testing, could be modified use in such a program. The method also may be adapted for other daphnid species, such as *Daphnia major* and *D. pulex*. Test conditions and endpoints from the EPA protocol are presented in Table 5. The test is continued until 60% of the control animals produce three broods of young. If this does not occur by 8 days, the test must be repeated. Environment Canada (1992) also has developed a similar chronic protocol for *C. dubia* that includes reproductive endpoints.

249. OECD revised its original daphnid test procedure (TG 201), separating the acute and chronic components into two guideline documents. The new chronic guideline (TG 211), which was written specifically to test *Daphnia magna*, was adopted in September 1998 (OECD 1998). OECD recommends that the data from the acute test be available for use in determining the appropriate test substance concentrations for the chronic evaluation. Newly hatched daphnids (<24-h old) are exposed to the test substance for 21 days. The primary endpoint measured is the total number of living young produced per adult daphnid that is still alive at the termination of the test. Any young produced by adults that die during the test are not included in the endpoint calculation. Secondary endpoints may include adult survival, growth, and the time to first brood. Japan (OECD 2003b) has proposed a draft enhancement to TG 211 that includes endpoints specifically intended to detect endocrine disruption in *Daphnia*. These are the offspring sex ratio, which is used as an endpoint for juvenile hormone-like chemicals; and molt inhibition, which is used as an endpoint for molting hormone-like chemicals. These endpoints are based on data reported by Baldwin et al. (2001), Olmstead and LeBlanc (2002, 2003), and Tatarazako et al. (2003).

250. ASTM reapproved its standard guide for conducting life-cycle assays using *Daphnia magna* (ASTM 2004). Newly hatched daphnids (<24-h old) are exposed to the test substance for 21 days. Biological data collected include mortality (recorded daily), the number of young produced (determined three times per week), size (dry weight) of the first-generation individuals still living at the end of the test, time to first reproduction, and behavioral abnormalities. Second-generation daphnid responses, such as survival, development, and behavior, may be obtained by observing these daphnids for an additional four days or more.

10.2 PROTOCOLS USING CHIRONOMID SPECIES AND OTHER SEDIMENT-DWELLING ORGANISMS

251. *Chironomus tentans* and *Hyalella azteca*: To test the potential sublethal toxicity of contaminated sediments on freshwater invertebrates, the EPA (2000) developed chronic testing methods for the freshwater-dwelling larvae of the midge, *Chironomus tentans*, and an amphipod, *Hyalella azteca*. Both species are ecologically important sediment dwellers in freshwater systems (Section 4.1). The protocol includes survival, growth, and reproductive endpoints (Table 5). Both species have been used successfully in EDC testing (Section 3.2), although not necessarily following procedures similar to this protocol. The EDC testing already performed on these species highlights the utility of including sediment-dwelling species in EDC evaluations. Similar protocols have been developed by Environment Canada (1997a, 1997b) for both species.

252. *Leptocheirus plumulosus*: EPA (2001) developed a chronic protocol to evaluate the potential sublethal toxicity of contaminated sediments to an estuarine amphipod, *Leptocheirus plumulosus*. The procedure runs for 28 days and includes survival, growth rate, and reproductive endpoints (Table 5). The test requires sediment as a substrate for the amphipods, but can be used with sediments having a wide variety of porewater salinities (1‰–35‰). This protocol could be adapted to the EDC program to test the potential effects of sediment-bound EDCs (created by spiking sediments) on a sediment-dwelling estuarine organism.

253. OECD has prepared draft technical guidelines for testing the effects of chemicals on chironomids. The guidelines focus on routes of exposure to sediment-dwelling invertebrates and include a spiked-sediment test (Technical Guideline 218; OECD 2001a) and a spiked-water test (Technical Guideline 219; OECD 2001b). Both tests are similar in approach (Table 6) and can be used for *Chironomus riparius*, *C. tentans*, or *C. yoshimatsui*. Both tests are chronic exposures lasting from 20 to 28 days for *C. riparius* and *C. yoshimatsui* or from 28 to 65 days for *C. tentans*. The endpoints for both tests are the total number of adults that emerge at the end of the exposure and development time. Additional short-term endpoints, survival and growth, can be obtained if additional replicates are included in the test design. Both protocols could be adapted to an EDC evaluation program with little modification.

10.3 PROTOCOLS USING COPEPOD SPECIES

254. Marine Copepods: OECD is preparing two technical guidance documents to standardize procedures for evaluating the effects of chemicals on calanoid (OECD 2004b) and harpacticoid (OECD 2004c) copepods. The calanoid guideline (Table 6) is designed specifically for *Acartia tonsa* and involves exposure of the animals from the egg to adult-producing stage (F₀ generation). Endpoints for this portion of the test include early life stage development, the onset of egg production and stable egg production, survival, sex ratio, and copepod length. The early life stages of the F₁ generation are exposed and the effects of the test substance on development are evaluated. The primary endpoint associated with this portion of the test is the Larval Development Ratio, which compares the total number of copepodites produced to the total number of early-stage individuals (nauplii + copepodites). The first portion of the test lasts 14–17 days, whereas the second runs 5–7 days. The harpacticoid guideline (Table 6) suggests that *Amphiascus tenuiremis*, *Nitocra spinipes*, and *Tisbe battagliai* are appropriate species to use for this procedure. Newly hatched larvae (<24-h old) are placed individually into microwell chambers (or other suitable containers) and exposed to the test substance. When the larvae mature (about 10–15 days), they are examined to determine gender, mating pairs are created, and placed into individual microwell chambers. The pairs are exposed to the test substance for 7–14 days, during which mating occurs. The test continues until the females have produced at least one brood. The numbers of young produced during exposure to the test material are compared to those produced in the control treatments. Additional endpoints that should be recorded are adult survival and the time to the first brood. Brood size, the

proportion of infertile eggs, and the time between broods may also be used in the evaluation of a test substance.

Table 5. Example US EPA protocols for evaluating chronic toxicity in Aquatic Arthropods

	EPA-821-R-02-013 October 2002	EPA/600/R-99/064 March 2000	EPA/600/R-99/064 March 2000	EPA/600/R-99/064 March 2000	EPA 600/R-01/020 March 2001
Test Species:	<i>Ceriodaphnia dubia</i>	<i>Chironomus tentans</i>	<i>Chironomus tentans</i>	<i>Hyalella azteca</i>	<i>Leptocheirus plumulosus</i>
Holding Conditions:	Hold at conditions similar to test	Hold at conditions similar to test or acclimate gradually to test conditions (Temperature at 1 °C/1-2 h)	Hold at conditions similar to test or acclimate gradually to test conditions (Temperature at 1 °C/1-2 h)	Hold at conditions similar to test or acclimate gradually to test conditions (Temperature at 1 °C/1-2 h)	Hold at conditions similar to test or acclimate gradually to test conditions (A change in temperature or salinity not exceeding 3 °C or 3 ‰ per 24 h)
Test Setup:					
Test organism age:	<24 h and all within 8 h of the same age	1 d (<24 h)	1 d (<24 h)	7 to 8 d	Neonates: age-selected (<48 h old) or size-selected: retained between 0.25 mm and 0.6 mm mesh screens
Duration:	Maximum of 8 d Until 60% or more of surviving control females have three broods	50–65 d depending on emergence. Each treatment may need to be terminated separately	50–65 d depending on emergence. Each treatment may need to be terminated separately	42 d	28 d
Test Material:	Effluents and receiving waters	Sediment	Sediment	Sediment	Sediment
Endpoint(s):	Survival and reproduction	20-d survival and weight; female and male emergence, adult mortality, the number of egg cases laid, the number of eggs produced, and the number of hatched eggs	20-d survival and weight; female and male emergence, adult mortality, the number of egg cases laid, the number of eggs produced, and the number of hatched eggs	28-d survival and growth; 35-d survival and reproduction; and 42-d survival, growth, reproduction, and number of adult males and females on Day 42	Survival, growth, and reproduction
Number of Treatments:	Effluents: 5 and a control Receiving Water: 100% receiving water (or a minimum of 5) and a control				
Concentration Series:	Effluents: ≥ 0.5 Receiving Waters: None or ≥ 0.5				

	EPA-821-R-02-013 October 2002	EPA/600/R-99/064 March 2000	EPA/600/R-99/064 March 2000	EPA/600/R-99/064 March 2000	EPA 600/R-01/020 March 2001
Dilution Water:	Uncontaminated source of receiving or other natural water, synthetic water prepared using Millipore milli-Q® or equivalent deionized water and reagent grade chemicals or DMW				
Solvent:	N/A				
Flow Conditions:		2 volume additions/d; continuous or intermittent (e.g., one volume addition every 12 h)	2 volume additions/d; continuous or intermittent (e.g., one volume addition every 12 h)	2 volume additions/d; continuous or intermittent (e.g., one volume addition every 12 h)	Siphon off and replace 400 mL 3 times/week
Number of Replicates:	10 (required minimum)	16 (12 at Day -1 and 4 for auxiliary males on Day 10)		12 (4 for 28-d survival and growth and 8 for 35- and 42-d survival, growth, and reproduction)	5 for toxicity test; ≥2 dummy chambers for pore water ammonia (Day 0 and Day 28)
Test Chamber:	30-mL borosilicate glass beakers or disposable polystyrene cups (recommended because they fit in the viewing field of most stereoscopes).	300-mL high-form lipless beaker		300-mL high-form lipless beaker	1-L glass beaker or jar with 10cm inner diameter
Test Volume:	15 mL (minimum)	100 mL sediment, 175 overlying water		100 mL sediment, 175 mL overlying water in the sediment exposure from Day 0 to Day 28 (175 to 275 mL in the water-only exposure from Day 28 to Day 42)	175 mL (about 2cm depth) sediment; approximately 725 mL water (fill to the 900mL mark on jar)
Number of organisms/rep:	1 Assigned using blocking by known parentage	12		10	20
Other Setup Notes:	New test solutions are prepared daily, and the test organisms are transferred to the freshly prepared solutions				
Test Conditions:					
Light:	Ambient laboratory illumination 10–20 μE/m ² /s, or 50–100 ft-c	Wide-spectrum fluorescent lights; About 100 to 1000 lux	Wide-spectrum fluorescent lights; About 100 to 1000 lux	Wide-spectrum fluorescent lights; About 100 to 1000 lux	Wide-spectrum fluorescent lights; About 500 to 1000 lux
Photoperiod:	16L: 8D	16L: 8D		16L: 8D	16L: 8D
Temperature:	25 °C±1 °C	23 °C ± 1 °C		23 °C ± 1 °C	Daily limits: 25 °C (±3 °C); 28-d mean: 25 °C (±2 °C).
pH:					7.0 to 9.0

	EPA-821-R-02-013 October 2002	EPA/600/R-99/064 March 2000	EPA/600/R-99/064 March 2000	EPA/600/R-99/064 March 2000	EPA 600/R-01/020 March 2001
Dissolved Oxygen:	4.0 mg/L	>2.5 mg/L	>2.5 mg/L	>2.5 mg/L	Daily Limits: ≥ 3.6 mg/L (50% saturation) 28-d mean: ≥ 4.4 mg/L (60% saturation)
Aeration:	used only as a last resort	if DO level falls below 2.5 mg/L: 1 bubble/second	if DO level falls below 2.5 mg/L: 1 bubble/second	if DO level falls below 2.5 mg/L: 1 bubble/second	28-d mean: ≥ 4.4 mg/L (60% saturation)
Salinity:	N/A	N/A	N/A	N/A	Daily limits: 5‰ ($\pm 3\%$) if pore water is 1‰ to 10‰, 20‰ ($\pm 3\%$) if pore water is 10‰ to 35‰; 28-d mean: 5‰ ($\pm 2\%$) or 20‰ ($\pm 2\%$)
Monitoring:					
WQ Frequency:	DO, temperature, pH: daily, prior to renewals in at least one test chamber at each concentration and control pH is measured in the effluent sample each day before new test solutions are made Conductivity, alkalinity, and hardness measured in each new sample and in the control	Hardness, alkalinity, conductivity, and ammonia at the beginning, on Day 20 and at the end of the test. Temperature daily (ideally continuously). DO and pH three times/week. Conductivity weekly. Measure DO more often if DO has declined by more than 1 mg/L since previous measurement.	Hardness, alkalinity, conductivity, and ammonia at the beginning and end of a sediment exposure (day 0 and 28). Temperature daily. Conductivity weekly. DO and pH three times/week. Measure DO more often if DO has declined by more than 1 mg/L since the previous measurement.	Hardness, alkalinity, conductivity, and ammonia at the beginning and end of a sediment exposure (day 0 and 28). Temperature daily. Conductivity weekly. DO and pH three times/week. Measure DO more often if DO has declined by more than 1 mg/L since the previous measurement.	Daily temperature in water bath or test or surrogate chamber, daily min/max recommended; salinity, temperature, DO, and pH at test initiation and termination, and in one replicate per sediment treatment preceding water renewal during the test (three times per week); aeration rate daily in all containers; total ammonia on Days 0 and 28 in one replicate per treatment.
Observation Frequency:	Daily	Daily to assess test organism behavior such as sediment avoidance	Daily to assess test organism behavior such as sediment avoidance	Daily to assess test organism behavior such as sediment avoidance	3 times/week in each test chamber preceding water renewal for condition and activity
Feeding:	0.1 mL each of YCT and algal suspension per test chamber daily	1.5 mL of Tetrafin® (4 mg/mL dry solids) to each beaker. If fungal or bacterial growth develop from excess food, feeding should be suspended for one or more days. If feeding is suspended in one treatment, it should be suspended in all treatments.	1.5 mL of Tetrafin® (4 mg/mL dry solids) to each beaker. If fungal or bacterial growth develop from excess food, feeding should be suspended for one or more days. If feeding is suspended in one treatment, it should be suspended in all treatments.	YCT food, fed 1.0 mL (1800 mg/L stock) daily to each test chamber. If fungal or bacterial growth develop from excess food, feeding should be suspended for one or more days. If feeding is suspended in one treatment, it should be suspended in all treatments.	3 times/week after water renewal Days 0–13, 20mg TetraMin® per test chamber; Days 14–28, 40mg TetraMin® per test chamber.

	EPA-821-R-02-013 October 2002	EPA/600/R-99/064 March 2000	EPA/600/R-99/064 March 2000	EPA 600/R-01/020 March 2001
Other Monitoring Notes:		At 20 d, 4 of the initial 12 reps. are selected for use in growth and survival measurements. AFDW of midges should be determined for the growth endpoint. Emergence traps are placed on the reproductive replicates on day 20		
Termination Notes:	Because of the rapid rate of development of <i>Ceriodaphnia dubia</i> , at test termination all observations on organism survival and numbers of offspring should be completed within two hours.	Clean sediment will typically require 40–50 d from initial setup to completion (emergence). Environmental stressors will reduce growth and delay emergence. For treatments in which emergence has occurred, the treatment (not the entire test) is ended when no further emergence is recorded over a period of 7 d.	Growth can be reported as either length or weight. Length should be measured \pm 0.1 mm from the base of the first antenna to the tip of the third uropod along the curve of the dorsal surface. Dry weight of amphipods in each replicate can be determined on Day 28 and 42.	Missing adult organisms should be recorded as dead. Growth rate of amphipods can be reported as daily change of average individual length or weight. Growth Rate (mg/individual/day) = (mean adult dry weight - mean neonate dry weight)/28. Count offspring within 2 weeks of termination.
Test Validity Criteria:	Mean control survival \geq 80%; 60% of surviving control females must produce at least three broods, with an average of 15 or more young per surviving female.	Average size of <i>C. tentans</i> in the control at 20 d \geq 0.6 mg/surviving organism dry weight or 0.48 mg/surviving organism AFDW. Emergence \geq 50%; mean number of eggs/ egg case \geq 800; percent hatch \geq 80%. Hardness, alkalinity, and ammonia in overlying water should not vary by more than 50% during the test, DO > 2.5 mg/L	Mean control survival \geq 80% on Day 28. Hardness, alkalinity, and ammonia in the overlying water typically should not vary by more than 50% during the sediment exposure, and dissolved oxygen should be maintained above 2.5 mg/L in the overlying water.	Mean control survival \geq 80% at the end of the test, with no single replicate having 60% survival or less.

Table 6. Example OECD protocols for evaluating chronic toxicity in aquatic arthropods

	OECD TG 211 September 1998	OECD TG218/219 February 2001	OECD Draft Calanoid TG	OECD Draft Harpacticoid TG
Test Species:	<i>Daphnia magna</i>	<i>Chironomus riparius</i> , <i>C. tentans</i> , <i>C. yoshimatsui</i>	<i>Acartia tonsa</i>	<i>Amphiascus tenuiremis</i> , <i>Nitocra spinipes</i> , <i>Tisbe battagliai</i>
Holding/ Culturing Conditions:	Hold at conditions similar to test	Hold at conditions similar to test	Hold at conditions similar to test	Hold at conditions similar to test
Test Setup:				
Test organism age:	<24 h; must not be first brood progeny	1 st instar larvae	1 st stage nauplii	Newly hatched nauplii (<24-h old)
Duration:	21 d	20–28 d; 28–65 d depending on species. If midges emerge earlier, test can be terminated five days after emergence of the last adult in the control.	19–25 d	20–25 d
Test Material:	Chemical in water	Spiked Sediment (218) or spiked water (219)	Seawater (natural or artificial)	Seawater (natural or artificial)
Endpoint(s):	Primary: Number of offspring per adult alive at end of test Other: parent survival; time to first brood Draft enhanced endpoints for endocrine disruption evaluations: offspring sex ratio; molting inhibition	Total adult emergence; development time	F ₀ : (14–17 d) early life-stage development, onset of egg production and stable egg production, survival, sex ratio, body length F ₁ : (5–8 d) development (larval development ratio, LDR)	Reproductive output of exposed animals versus control(s) Parent animal survival Time to first brood should also be reported. Also, brood size, infertile or unhatched eggs, time interval between successive broods and possibly intrinsic or instantaneous rates of population increase (r_m or λ), may be examined
Number of Treatments:	≥5 and a laboratory control	≥5 and a laboratory control	≥5 and a laboratory control	≥5 and a laboratory control

	OECD TG 211 September 1998	OECD TG218/219 February 2001	OECD Draft Calanoid TG	OECD Draft Harpacticoid TG
Concentration Series:	Geometric series with separation factor ≤ 3.2 ; based on acute test or range-finding test	Factor between concentrations should be ≤ 2 for EC_x studies; ≤ 3 for LOEC/NOEC studies	Geometric series with separation factor ≤ 3.3 Test concentrations should not include any that have a statistically significant effect on survival of the F_0 generation since the main objective is to measure sublethal effects	Geometric series with separation factor ≤ 3.2 Test concentrations should not include any that have a statistically significant effect on survival since the main objective is to measure sublethal effects
Dilution Water:		Any suitable natural or synthetic water may be used. The water has to be aerated before use. May be used to make stock for dosing test (e.g., acetone, ethanol, methanol)	DO saturation value $>70\%$; pH = 8.0 ± 0.3 before use	Constant quality; DO saturation value $>70\%$; pH = 8.0 ± 0.3 before use
Solvents:	May be used to make stock for dosing test (e.g., acetone, ethanol, methanol)			
Dispersants:	May assist in accurate dosing and dispersion (e.g., Cremophor RH40, HCO-40)			
Flow Conditions:	Semi-static; flow-through allowable	Static	Static-renewal	Static-renewal
Number of Replicates:	10 (semi-static); 4 (flow through)	≥ 3 for EC_x study; ≥ 4 for LOEC/NOEC study	Control: minimum 6, 12 recommended for LDR test design; 4 (linear regression), 6 (ANOVA)	Minimum 3 microplates/concentration (48–144 wells)
Test Chamber:	Glass beakers; volume not specified	600-mL, 8-cm diameter glass beaker	100-mL beaker	Microwell (300–5000 μL total volume) test chambers preferred
Test Volume:	50–100mL	1.5–3.0 cm sediment depth	80 mL	<50 mL
Number of organisms/rep:	1 (semi-static); 10 (flow through)	20 for <i>C. riparius</i> and <i>C. yoshimatsui</i> ; 12 for <i>C. tentans</i>	10	48 animals added to three replicate test vessels (microplates) per concentration and control(s)

	OECD TG 211 September 1998	OECD TG218/219 February 2001	OECD Draft Calanoid TG	OECD Draft Harpacticoid TG
Other Setup Notes:	Renew test solution at least 3 times/week; more frequently if not stable (<80% initial concentration over 3 d) Prepare new test vessels and transfer adults to them	Formulated sediment (also called reconstituted, artificial or synthetic sediment) should be used. Spike sediment for TG218; spike water for TG219	LDR test: renew 50–80% on Day 3 or increase volume Basic exposure: renew 50–80% every 2 nd day or increase volume Prepare new test vessels and transfer adults to them	Depends on stability of test substance, should be at least three times per week Prepare new test vessels and transfer adults to them With microplates, evaporation losses ≤ 7% daily; correct by adding deionized or distilled water
Test Conditions:				
Light:	Ambient laboratory illumination <15-20 μE/m ² /s	About 500 to 1000 lux	Wide-spectrum fluorescent lights; low intensity; ~5-10 μmol/s/m ²	Not specified
Photoperiod:	16L:8D	16L:8D	12L:12D	<i>N. spinipes</i> –0L:24D <i>A. tenuiremis</i> –12L:12D <i>T. battagliai</i> –16L:8D
Temperature:	18–22°C; should not vary >2°C during any one test	20°C ± 2°C; 23°C ± 2°C for <i>C. tentans</i>	15–20°C; vary < 2°C during test	20-25°C; vary ≤ 2°C during test
pH:	6–9; should not vary >1.5 units	6–9 at start of test	8.0 ± 0.3	± 0.3 units of control
Dissolved Oxygen:	3 mg/L	>60% saturation value	>70% saturation value	>70% saturation value (dilution water)
Aeration:	None allowed	Gentle aeration; 1 bubble/second	if DO level falls below 70% saturation: 1 bubble/second	None during test
Hardness/ Salinity:	>140 mg/L CaCO ₃	<200 mg/L CaCO ₃	~20 ‰ ± 2 ‰	Use same as culture water; <i>N. spinipes</i> 1–35‰ <i>A. tenuiremis</i> 15–35‰ <i>T. battagliai</i> 20–35‰
Monitoring:				

	OECD TG 211 September 1998	OECD TG218/219 February 2001	OECD Draft Calanoid TG	OECD Draft Harpacticoid TG
WQ Frequency:	DO, temperature, hardness, pH: once per week in fresh and old media, controls, and highest test concentration Determine concentration of test substance regularly	DO measured daily in all test vessels Hardness and ammonia at the beginning and at the end of the test in controls and one vessel of highest concentration. Temperature and pH at start and end of test.	Measure dissolved oxygen, pH, salinity, and temperature in controls and all test concentrations each time test medium is renewed. As a minimum, measurements shall be made in the control(s) and highest test concentration. Temperature preferably monitored continuously.	Measure dissolved oxygen, pH, and temperature should be measured in the controls and all test concentrations each time test medium is renewed. As a minimum, measurements should be made in the control(s) and highest test concentration. Temperature preferably monitored continuously in at least one test vessel
Observation Frequency:	Not specified	At least three times per week; visual assessment of abnormal behaviour (e.g. leaving sediment, unusual swimming), compared with the control.	Daily for egg production test; not specified for LDR test	Not specified
Feeding:	At least 3 times per week, preferably daily (semi-static); concentrated algal suspension	At least 3 times per week, preferably daily; 0.25-0.5 mg fish-food (a suspension in water or finely ground food) per larvae per day for the first 10 days. Slightly more food for older larvae; 0.5-1.0 mg per larvae per day Reduce food ration in all treatments and control if fungal growth occurs or if mortality is observed in controls. If fungal development cannot be stopped the test is to be repeated.	LDR test: 5×10^4 cells/mL of medium added on days 0 and 3 Basic exposure: 5×10^4 cells/mL of medium added on day 0 and subsequently to every renewal of the medium. Egg production: 5×10^4 cells/mL of medium added daily.	<i>A. tenuiremis</i> : 2 μ L of 1:1:1 volume (10^7 cells/mL) chlorophyte, chrysophyte, diatom suspension every 6 d during maturation/development and reproduction. <i>N. spinipes</i> : ($\sim 10^7$ cells/mL) chrysophyte, preferably <i>Rhodomonas salina</i> or <i>R. baltica</i> , every third day during maturation/development and reproduction; final concentration in each test chamber is 10^5 cells/mL. <i>Tisbe battagliai</i> : ($\sim 3 \times 10^7$ cells/mL) chrysophyte, preferably <i>Rhodomonas reticulata</i> , every third day during maturation/development and reproduction, so that the final concentration in each test chamber is 2×10^5 cells/mL.

	OECD TG 211 September 1998	OECD TG218/219 February 2001	OECD Draft Calanoid TG	OECD Draft Harpacticoid TG
Other Monitoring Notes:	Record parent animal mortality daily Remove and count offspring daily Record presence of aborted eggs or dead offspring daily	During the period of expected emergence, a daily count of emerged midges is necessary.		
Termination Notes:		Record number of fully emerged male and female midges	Fix animals in Lugol's solution; score biological parameters (count, measure, determine gender, etc.)	Fix animals in Lugol's solution or formaldehyde (4%); score biological parameters (count, measure, determine gender, etc.)
Test Validity Criteria:	Parent animals (female <i>Daphnia</i>) mortality $\leq 20\%$ at the end of the test Mean number of live offspring produced per parent animal surviving at the end of the test > 60	Control mortality $\leq 30\%$ at end of test <i>C. riparius</i> and <i>C. yoshimatsui</i> emergence to adults occurs 12–23 days after initiation; <i>C. tentans</i> emergence to adults occurs 20–65 days after initiation Mean emergence in controls 50–70 per cent DO concentration $\geq 60\%$ air saturation value at the end of the test pH of overlying water 6–9 range in all test vessels Water temperature vary ≤ 1.0 °C between vessels during test Temperature held within the ranges specified for test species	Mortality in mass culture control(s) $\leq 20\%$ at end of exposure DO concentration $\geq 70\%$ air saturation value throughout the exposure period Temperature vary $\leq 2^\circ\text{C}$ during the test pH vary ≤ 0.3 units from average control pH Salinity vary $\leq 10\%$ from the control start value Sex ratio in controls = 40:60 to 60:40 Hatching success in hatching control $\geq 80\%$ Average mortality in the control(s) on day of LDR observation $\leq 20\%$ Average number of eggs/female /d in controls ≤ 30 at Day 12 and later Control mortality of the isolated females in the egg production test (Day 7–12) $\leq 20\%$.	Control mortality $\leq 20\%$ at the end of the exposure period DO concentration $\geq 70\%$ air saturation value throughout the exposure period Temperature vary $\leq 2^\circ\text{C}$ during the test pH vary ≤ 0.3 units from average control pH Salinity vary $\leq 10\%$ from the control start value Sex ratio in controls = 40:60 to 60:40 Average number of offspring per clutch in the controls ≥ 5 Average number of days to first brood in the control(s) \leq number of days = 50% of total exposure period (at a temperature of 20°C)

10.4 PROTOCOLS USING MYSID SPECIES

255. Specific guidance for conducting short-term toxicity tests with species other than *A. bahia* has been published. It is possible that these protocols can be modified to allow longer life cycle testing.

256. *Holmesimysis costata*.—Chapman et al. (1995) described a 7-day test protocol designed to measure growth and survival in tests using the west coast mysid species, *H. costata*. In addition to recommended test conditions, guidance in culturing the animals and analyzing the data are presented. The protocol describes the ecological and culture requirements for *H. costata*; this information could be used to modify the ASTM and OPPTS protocols described above to allow longer life cycle testing. The test protocol was evaluated by means of a series of intra- and interlaboratory comparisons (Hunt et al. 1997), which concluded that this test had sufficient sensitivity and precision to make it useful in testing possible contaminant impacts. A summary of the test conditions recommend by this protocol is presented in Table 7.

Mysidopsis intii.—A short-term toxicity test protocol for a west coast species, *M. intii*, was developed with support from the EPA (Langdon et al. 1996). The protocol concisely describes the test conditions required to conduct a 7-day toxicity test to measure survival and growth of this species. Initial test development was performed using zinc sulfate as the toxicant. The test protocol was evaluated by means of an interlaboratory comparison that employed sodium dodecyl sulfate as the toxicant (Harmon and Langdon 1996). Harmon and Langdon (1996) also compared *M. intii* test with those using *A. bahia* and *H. costata*, and reported that its sensitivity was equal to that of the *A. bahia* test, but that it was lower than that of the *H. costata* test. A summary of the test conditions recommended by this protocol is presented in Table 7.

Table 7. Recommended test holding conditions for *Holmesimysis costata* and *Mysidopsis intii*

	Chapman et al. (1995)	Langdon et al. (1996) Harmon and Langdon (1996)
Test Species:	<i>Holmesimysis costata</i>	<i>Mysidopsis intii</i>
Holding Conditions:	WQ similar to test conditions 4, 1000-L tanks ½ volume changed twice/week <20 mysids/L Feed: ≤24 h old <i>Artemia</i> (5–10/mysid/d); plus 10–20 mg ground Tetramin [®] /100 mysids/d	WQ similar to test conditions 3-L Pyrex glass beakers 2 L volume; 90% exchange every 2 d 15/beaker Feed: 2-d old enriched <i>Artemia</i> + <i>Tigriopus californicus</i>
Test Setup:		
Test organism age:	3–4 d post hatch	0 or 2 d growth 6 d reproductive condition 15 d reproductive output
Duration:	7 d	7 d
Test Material:	Effluent	NS ^a
Endpoint(s):	Survival; growth	Survival; growth, reproductive condition, reproductive output
Number of Treatments:	NA ^b	NA
Concentration Series:	Minimum of 5 and 1 control	No guidelines specified (see reference toxicant section)
Dilution Water:	Uncontaminated 1-µm-filtered natural seawater or hypersaline brine prepared from natural seawater	NS

	Chapman et al. (1995)	Langdon et al. (1996) Harmon and Langdon (1996)
Flow Conditions:	Static renewal	Static renewal
Delivery System:	Manual remove/replace	
Flow Rate:	75% renewal at 48 h and 96 h	90% change every 2 d
Calibration:	NA	NA
Calibration Check:	NA	NA
# Replicates:	5	3
Test Chamber:	1000 mL	1-L Pyrex glass beaker
Test Volume:	200 mL	1 L
#organisms/rep:	5	15
Other Setup Notes:	NS	NS
Test Conditions:		
Light:	10–20 $\mu\text{E}/\text{m}^2/\text{s}$ (ambient laboratory)	1000 lux
Photoperiod:	16 h light :8 h dark	16 h light :8 h dark
Temperature:	13 °C \pm 1 °C (N of Pt. Conception) 15 °C \pm 1 °C (S of Pt. Conception)	20 °C \pm 2 °C
pH:	NS	8.0 \pm 0.3
Dissolved Oxygen:	>4.0 mg/L	> 60% saturation (at test conditions)
Aeration:	None unless needed to maintain above limit; then < 100 bubbles/min.	None unless needed to maintain above limit.
Salinity:	34 ‰ \pm 2 ‰	34 ‰ \pm 2 ‰
Monitoring:		
WQ Frequency:	At beginning and end of exposure in one chamber per treatment; temperature daily in two chambers per environmental control system. (WQ includes pH).	Before and after each water exchange
Observation Frequency:	Daily; count and record number of live mysids; remove dead mysids and excess food	number of live mysids recorded at each water exchange
Feeding:	\leq 24-h-old <i>Artemia</i>	\leq 24-h-old <i>Artemia</i> + <i>Tigriopus californicus</i> Varies with mysid age
ration	40/mysid/day (two feedings/day, 20 each)	
Other Monitoring Notes:	NS	NS
Termination Notes:	Remove and record dead mysids Count survivors (considered alive if visibly respond to stimulus) Screen surviving mysids to remove from chamber, count, rinse in deionized water, transfer to tared weighing boat, dry at least 24 h at 60 °C	Growth and Survival: Remove and count surviving mysids, place in 10 mL dilution water, and refrigerate at 5° C for 24 h Fix in 5% buffered formaldehyde Measure body length (base of eyestalk to posterior edge of exopod) of 10 individuals + Reproductive Condition: Determine percent females carrying eggs or juveniles in brood sac + Reproductive Output: Determine number of juveniles released per final number of females
Test Validity Criteria:	A test is valid if \geq 75% control survival \geq 0.40 μg dry wt in control; survival MSD $^{\circ}$ <40%; growth MSD <50 μg ; NOECs , 100 $\mu\text{g}/\text{L}$ with zinc.	A test is valid if \geq 85% control survival \geq 3.0 mm total body length for controls
Reference Toxicant:	Zinc sulfate	NS (protocol development tested zinc sulfate and sodium dodecyl sulfate)

	Chapman et al. (1995)	Langdon et al. (1996) Harmon and Langdon (1996)
Concentration Series	0, 10, 18, 32, 56, 100 µg/L total zinc	Zinc sulfate: 0, 37, 50, 70, 101, 230 µg/L
number of reps	5	
test volume	NS	<u>SDS:</u>
test containers	NS	0, 1.58, 2.51, 3.97, 6.30, 10.0 mg/L
# of animals/rep	NS	
Termination Notes:	NS	NS
Test Validity Criteria:	NS	NS

a) NS Not specified.

b) NA Not applicable.

c) MSD Minimum significant difference.

10.5 CANDIDATE PROTOCOL FOR EDC-DETERMINATION STUDIES AND ADDITIONAL DATA NEEDS

257. In this section, we recommend the test species and the protocol for its use that would be most suitable for EDC-determination studies. Gaps in the current knowledge are evaluated, and necessary supplementary studies are recommended.

258. The utility of invertebrates in general can be justified on the basis of the ease of culture of many species from different phyla that have relatively short life cycles. Although there is limited documentation to date of invertebrate responses to EDCs, we have reviewed in previous sections the studies that have been conducted primarily in the lab, but also a few that were done *in situ*. The reproductive system of invertebrates appears to be particularly vulnerable to EDCs. Of particular interest in using *aquatic* invertebrate species as test organisms is that the neuroendocrine centers in the eyestalk, brain, and thoracic ganglia produce substances that regulate ion movements in tissues of crustaceans in freshwater and saltwater habitats.

259. The test protocol presented here is a proposed guideline submitted to OECD (Appendix B) for consideration for further development, validation, and acceptance. The protocol provides general guidance for conducting a two-generation toxicity test in which the parent generation (F0) is exposed to the toxicant and the parent and first offspring (F1) generations are monitored for 7 days after the mean date of release of the second brood by the control treatment parent generation (F0). The guideline also allows for the exposure of the F1 generation to the test substance. The guideline is applicable to *Americamysis bahia*, but could be modified to suit other mysid species. The test is an extension of previous mysid life-cycle test procedures produced by ASTM (ASTM 1997) and EPA (Nimmo et al. 1977, 1978a, 1978b, McKenney 1986, 1998).

10.5.1 Preferred species

260. The preferred mysid species for use in the testing of potential EDCs is *Americamysis bahia* (Molenock 1969). The primary reasons for its selection are that it is commercially cultured and readily available year-round, it has been the subject of many toxicity tests, it has a short generation time, and its testing requirements and biology are well known. It can be cultured and maintained easily by testing laboratories. These advantages outweigh the disadvantage that in some situations, EDC testing with this species may mean using a test organism that is not indigenous to the geographic area of interest.

10.5.2 Description of the method

261. The two-generation test is initiated when healthy < 24-h-old mysid juveniles are placed randomly into replicate test chambers. These original juveniles comprise the parent (F0) mysid stock for the test. Development, sexual maturation, reproduction, and growth are observed in the F0 and F1 generations. The assay is conducted with at least five toxicant concentration treatments and appropriate control treatments (typically a 0% concentration and a solvent control, if one was used to deliver the toxicant to the test treatments). The highest exposure concentration should be equal to the lowest concentration that caused adverse effects in the acute test or 1/10 the LC₅₀ (EC₅₀). Toxicant and control treatments are delivered to the mysids in water, which is continuously or intermittently delivered to the test chambers via a proportional diluter system or infusion pumps. The specific exposure duration will vary, but is at least 7 days longer than the median second-brood-release date by the original parent mysids in the control treatments. During the test, chambers are examined for mysid mortality, the presences of molted exuviae, the presence of ovigerous females, and the release of young, all of which are recorded. Young from the

first brood (= F1') release by the parent stock are held for four days, after which they are counted and measured. The parent stock is allowed to produce a second brood (= F1''), after which the parent mysids are counted and measured. The F1'' mysids are exposed to the test materials and maintained until their first brood is released (= F2), after which the F1'' mysids are counted and measured. The test is terminated with the release of the F2 mysids, which are counted. Surviving organisms may be analyzed biochemically, as appropriate.

10.5.2.1 General Procedures and Equipment

262. The test mysids shall be obtained from a single batch of juveniles obtained from the same brood stock. The brood stock must have been hatched and raised in the testing laboratory or obtained by the laboratory prior to sexual maturity and held at test environmental conditions for at least 14 days. Food during holding must be the same as that used during the test. Animals selected for testing must be <24 h old and must not exhibit abnormal behavior or morphology. Brood stock holding tanks must contain no more than 20 mysids/L and be free of other organisms.

263. The test generally should consist of five toxicant concentration treatments and appropriate control treatments. The highest exposure concentration should be equal to the lowest concentration causing an adverse effect in an acute test or equal to 1/10 the LC₅₀ (EC₅₀). Each concentration, except for the control and the highest concentration, should be at least 50% of the next highest concentration. The definitive test concentration series can be determined by using known results from acute toxicity studies or by conducting a range-finding test in which the concentrations tested are widely separated, such as values of 1, 10, and 100 mg/L. The experimental unit for the mysid test is the test chamber and is defined as the smallest unit to which treatments can be independently applied. The suggested definitive test chamber that is used is a 98 × 14 × 15 cm container that can hold the suggested 21-L test volume and two retention baskets, constructed of 15-cm glass Petri dishes to which a 15 cm high cylinder of nylon mesh screen (210 μm mesh) has been attached. ASTM (1997) and some other protocols call for the use of relatively large tanks that are subdivided into compartments. However, the toxicant concentration in such cases is not applied independently to all compartments, which are therefore not considered experimental units or replicates.

264. The number of replicates used in various protocols and tests ranges from two to eight. Although it may be desirable to determine the number of replicates based on the expected variation, desired detection limit, and selected power, as suggested by ASTM (1997), the number selected will most likely be constrained by the capacity of the diluter system. However, a minimum of three replicate chambers per concentration, including the control treatment, are included in the recommended design.

265. Exposed Parental mysids (F0).—Fifteen mysid juveniles (< 24 hours old) are randomly assigned to reproduction/survival retention baskets within each replicate tank. Within each replicate tank, one additional retention basket is initiated with 15 mysid juveniles that can be subsampled weekly for growth measurements, or used to measure optional biochemical endpoints. Upon reaching maturity (approximately 13–16 d), one male and one gravid female are randomly assigned to each of the brood cups within each replicate (with a maximum of 7 male/female pairs possible per replicate). Individuals in the reproduction/survival retention baskets not paired in brood cups will be maintained and observed within the retention baskets for survival and sex determination until they are paired or until the test is terminated (one week after the mean day of release of the second brood).

266. Unexposed or Exposed F1 generation.—Offspring from the first brood (F1'), the second brood (F1''), or both, are transferred to clean dilution water or exposed water for all treatments (dependant on availability of 15 young from each treatment on the same day). The F1 generation evaluation is initiated with 15 newly released mysid juveniles (< 24 hours old) randomly assigned to F1 generation reproduction/survival retention baskets within a separate tank for each of the F0 exposure conditions.

Mixing of young released on the same day, but across replicate chambers, is encouraged to minimize reproduction between siblings. An additional F1 generation retention basket is initiated with 15 newly released mysid juveniles within each replicate that can be subsampled weekly for growth measurements, or used to measure optional biochemical endpoints. Upon reaching maturity (approximately 13–16 days), one male and one gravid female are randomly assigned to separate F1 generation brood cups within each replicate (with a maximum of 7 male/female pairs possible per replicate). Individuals in the F1 generation reproduction/survival retention baskets not paired in brood cups will be maintained and observed within the F1 generation retention baskets for survival and sex determination until they are paired or until the test is terminated (one week after the mean day of release of the second brood).

267. The requirements for a given test facility should accommodate the use of continuous or intermittent flow-through or recirculating tanks for holding and/or testing. The ideal would be to use proportional diluters with an elevated head box to allow for gravity-fed dilution water into the brood tanks or chemical mixing chambers. A metering system could be used to mix and deliver test concentrations to the appropriate testing chamber. The test chamber must be maintained at a constant temperature using either temperature-controlled water or recirculating water baths. The water and air going into the testing system should be strained of any particulate matter using either strainers or air and water filters capable of filtering material through a 0.20- μ m bacterial filter (ASTM 1997, EPA 1996, OECD 2004a).

268. Lighting conditions for testing require the use of timers capable of delivering light for a 14-h light and 10-h dark cycle with a 15–30-minute transition period. The transition period is important, because mysids may become stressed by instantaneous changes in light. In the natural environment, the normal vertical migration of mysids allows for gradual acclimation to light changes. Under laboratory conditions, instantaneous change in light has been observed to cause jumping or impingement on the sides of the testing container (ASTM 1997).

269. The test facility should be well ventilated and free of fumes. Laboratory ventilation systems should be checked to ensure that return air from chemistry laboratories or sample handling areas is not circulated to culture or testing rooms, or that air from testing rooms does not contaminate culture rooms. Air pressure differentials between rooms should not result in a net flow of potentially contaminated air to sensitive areas through open or loose-fitting doors. Air used for aeration must be free of oil and fumes. Oil-free air pumps should be used where possible. Filters to remove oil, water, and bacteria are desirable. Particles can be removed from the air using filters such as BALSTON Grade BX (Balston, Inc., Lexington, Massachusetts) or equivalent, and oil and other organic vapors can be removed using activated carbon filters (e.g., BALSTON C-1 filter) or equivalent (EPA 2000). During phases of rearing, holding, and testing, test organisms should be shielded from external disturbances such as rapidly changing light or pedestrian traffic (EPA 2000).

270. Equipment and supplies that contact stock solutions, sediment, or overlying water should not contain substances that can be leached or dissolved in amounts that adversely affect the test organisms. In addition, equipment and supplies that contact sediment or water should be chosen to minimize sorption of test materials from water. Glass, type 316 stainless steel, nylon, and high-density polyethylene, polypropylene, polycarbonate, and fluorocarbon plastics should be used whenever possible to minimize leaching, dissolution, and sorption. Concrete and high-density plastic containers may be used for holding and culture chambers, and in the water-supply system. These materials should be washed in detergent, acid-rinsed, and soaked in flowing water for a week or more before use. Cast-iron pipe should not be used in water-supply systems, because colloidal iron will be added to the overlying water and strainers will be needed to remove rust particles. Copper, brass, lead, galvanized metal, and natural rubber must not contact overlying water or stock solutions before or during a test. Items made of neoprene rubber and other materials not mentioned above should not be used unless it has been shown that their use will not adversely affect survival, growth, or reproduction of the test organisms (EPA 2000).

271. For administration of test substance via water, the recommended equipment consists of proportional diluters, metering systems, pumps, or other suitable systems to be used to deliver test substances to the testing chambers. The system that is chosen should permit the mixing of test material with dilution water before its entrance into the testing chambers, and should supply the selected concentrations in a reproducible fashion (ASTM 1997).

272. The system must be calibrated before use to determine the flow rate into the chambers and the test concentration entering each chamber. It is advisable to allow the test solutions to flow through the system for a time sufficient to allow concentrations of the test concentrations to reach a steady state. Then two sets of samples should be taken at least 24 h apart. The chemical analysis of the concentrations should verify that the test concentrations have reached a steady state before organisms are placed into testing chambers. The measured test concentrations should be within 30% of estimated or nominal concentrations. The delivery system should provide for at least a 90% volume exchange every 8 to 12 h for continuous-flow operations, or 70% exchange every 5 h for intermittent-flow operations. The flow rate through any two chambers should not differ by more than 10%. The general operation of the test delivery system should be functioning properly for 48 h prior to the initiation of a test (OECD 2004a) and should be checked twice per day, usually morning and afternoon during the test (ASTM 1997, EPA 1996).

10.5.2.2 Test Validity

273. The test acceptability criteria generally follow those listed in EPA (1996), with some guidance from the ASTM (1997) procedure. Basic principles of experimental design must be followed. All test chambers must be identical, treatments must be randomly assigned to test chamber locations, all appropriate control treatments must be included, and individual mysids must be allocated randomly or impartially into the test chambers. Tests conditions (e.g., water quality) must be within the specified guidelines. Control treatment survival for first generation mysids must be at least 70%. At least 75% of the F0 or F1 females in the control treatments must produce young, and the average number of young produced per control female in the first two broods must be at least eight.

Table 8. Recommended Mysid Life Cycle Toxicity Test Conditions

	ASTM E1191 (ASTM 1997)	OPPTS 850.1350 (EPA 1996)
Test Species:	<i>Americamysis bahia</i> <i>A. bigelowi</i> <i>A. almyra</i>	<i>Americamysis bahia</i>
Holding Conditions:	Hold at conditions similar to test or acclimate gradually to test conditions (Temperature at 3°C/12 h salinity at <3‰/24 h) 76-L aquaria Flow through or recirculating system 14 h light:10 h dark, or 16 h light: 8 h dark, with 15–30 min transition period Gentle aeration Feed excess ≤24 h old <i>Artemia</i> ; 150/mysid/day; may supplement with algae or other food	Hold at conditions similar to test or acclimate gradually to test conditions (Temperature at 1°C/24 h; salinity at <5‰/24 h) Flow through or recirculating system 14 h light:10 h dark, with 15–30 min transition period Aeration if needed
Test Setup:		
Test organism age:	≤24 h	≤24 h
Duration:	≥7 days after median first brood release in controls	28 days
Test Material:	Reagent grade or better	NS ^a
Endpoint(s):	Survival, growth, reproduction	Survival, growth, young produced
Number of Treatments:	≥5 plus control (add solvent control if necessary)	≥5 plus control (add solvent control if necessary)

	ASTM E1191 (ASTM 1997)	OPPTS 850.1350 (EPA 1996)
Concentration Series:	Test concentrations should bracket the highest concentration at which there is not an unacceptable effect; each concentration should be at least 50% of the next highest concentration	5 or more concentrations chosen in geometric ratio between 1.5 and 2.0.
Dilution Water:	Natural or reconstituted seawater acceptable to saltwater mysids; uniform quality during test; should not affect test outcome	Natural (>20- μ m-filtered) or artificial seawater
Solvent:	Must allow satisfactory survival, growth, and reproduction If solvent used, ≤ 0.1 mL/L concentration.	If solvent used, ≤ 0.1 mL/L concentration
Flow Conditions:	Flow through	Flow through
Delivery System:	Proportional diluter	Proportional diluter
Flow Rate:	>5 volume additions/24 h (must be capable of 10 additions/24 h)	5 x chamber volume/24 h
Calibration limit:	<10%/chamber/time	<10%/chamber/time
Calibration/ Check:	Prior to test; visual check twice daily	Prior to test; twice daily
Number of Replicates:	Variable, estimated according to expected variation, desired detection limit, and selected power.	5+ (minimum 40 mysids/treatment)
Test Chamber:	e.g., 300 mm x 450 mm x 150 mm deep with adequate compartments (to provide 30 cm ² /mysid).	Volume NS; materials must minimize sorption of test chemicals; loosely covered
Test Volume:	Solution depth ≥ 100 mm (in above specified chamber)	NS
Number of organisms/rep:	NS (recommends 1 male-female pair/ compartment, but can't determine gender for ~12 d)	8 (maximum)
Other Setup Notes:	NS	NS
Initiation Notes:	NS	NS
Test Conditions:		
Light:	NS	NS
Photoperiod:	14 h light:10 h dark, or 16 h light:8 h dark, with 15–30 min transition period	14 h light:10 h dark with 15–30 min transition period
Temperature:	27°C (for <i>A. bahia</i>); 3°C individual measurements: $\pm 1^\circ\text{C}$ time-weighted average; <2°C difference between any two jars measured concurrently.	25°C $\pm 2^\circ\text{C}$
pH:	6.6–8.2	NS
Dissolved Oxygen:	A concentration between 60–100% of saturation is best.	60–105% saturation
Aeration:	yes	
Salinity:	15–30‰; variation among treatments should be <5, must be <10 ‰	20‰ ± 3 ‰
Monitoring:		
Test Concentration	Twice prior to test, at 24 h apart; Measured concentration $\leq 30\%$ of nominal concentration. During test frequently enough to establish average and variability, at least weekly.	At Day 0, 7, 14, 21, 28; should vary <20% among replicates/concentration
WQ Frequency:	Salinity daily; temperature in one chamber hourly or min/max measured daily; pH at start and end of test and weekly in control, include highest concentration; dissolved oxygen in at least one test chamber at start and end and weekly	Weekly (includes pH)

	ASTM E1191 (ASTM 1997)	OPPTS 850.1350 (EPA 1996)
Observation Frequency:	Daily Count, determine gender, remove dead G1 mysids; count live mysids; record number live females Record day of brood release; count and remove young daily Record abnormal development and aberrant behavior	Periodically; record number dead on Day 7, 14, 21, 28
Feeding:	Live brine shrimp nauplii at least once daily; may supplement Dead brine shrimp should be removed daily before feeding occurs.	Recommend 48-h-old <i>Artemia</i> . Frequency and amount not specified.
Other Monitoring Notes:	Weekly determinations of particulate matter, total organic carbon, and total dissolved gasses desirable	NS
Termination Notes:	Count live G1 mysids and determine gender Desirable to measure total body length (anterior tip of carapace to tip of uropod) Obtain dry weight of surviving G1 (males and females separate); remove any brine shrimp present; rinse mysids in deionized water, dry at 60°C for 72–96 h Morphological observations at end of test may be desirable May be desirable to hold G2 mysids for 4+ day longer to observe possible effects	Record number of dead on Day 7, 14, 21, 28 Record number of males & females and measure body length (anterior tip of carapace to tip of uropod) when distinguishable and on Day 28. Count and separate G1 offspring as produced, hold at test concentrations. If possible (i.e., by Day 28), count, determine gender and measure G2 mysids. Record abnormal behavior or morphology.
Test Validity Criteria:	A test is valid if General test requirements are met ≥70% G1 control survival ≥75% G1 control females produce young ≥3 average number of young/female	A test is valid if ≥75% G1 control females produce young ≥3 average number of young/female/day
Range-Finding Test		
Concentration Series	NS	Widely spaced; e.g., 1, 10, 100mg/L
number of samples		1
test volume		NS
test containers		NS
number of animals/rep		Minimum 10/concentration
duration		NS, allow estimate of test concentrations
Termination Notes:	NS	NS
Test Validity Criteria:	NS	NS
Reference Toxicant:	NS	NS
Concentration Series	NS	NS
number of reps		
test volume		
test containers		
number of animals/rep		
Termination Notes:	NS	NS
Test Validity Criteria:	NS	NS

a) NS Not specified.

10.5.3 Endpoints: appropriateness and preferred method for quantification

10.5.3.1 Reproductive and Developmental Endpoints

274. Because potential EDCs may elicit more than one response and the responses may vary with the chemical tested, several endpoints are included in the testing program. Many of the adverse responses to

exposure to toxicants that are typically measured in toxicity testing programs (e.g., survival, growth, reproductive biology) are important in determining whether or not a potential EDC could cause significant impacts to wild populations. However, all of these responses might not be attributable to the disruption of endocrine processes. Therefore, to conclude that a particular toxicant is an endocrine disruptor, it is also necessary to establish that exposure to toxicants has affected endocrine systems.

275. The primary endpoints that can indicate possible impacts to populations that should be included in a toxicity test are described below.

276. *Survival* is recorded as the proportion of individuals still living at the termination of the appropriate phase of the test. Mysids are considered dead if they are opaque white in color, immobile (especially regarding respiratory and feeding appendages), and do not respond to gentle prodding. Missing animals should be considered dead. Survival is determined for F0 and F1 mysids.

277. *Growth* is a sublethal developmental endpoint that can be reduced by exposure to EDCs or other stressors, and is therefore appropriate to include. The inclusion of growth as an endpoint may help distinguish between the responses. Growth is measured as the difference in the dry weight of test organisms before and after exposure to test and control treatments. Dry weight should be measured by rinsing all surviving mysids carefully in deionized water to remove salt. Any *Artemia* nauplii that may be caught in female brood pouches should be removed and all amphipods should be dried at 60°C until constant weight is reached (usually <72–96 h). Weight measurements should be made to the nearest 0.01 mg (as suggested in EPA 2000, OECD 2004a). Growth, at Day 7 and Day 14, is measured by replicate for F0 and F1 mysids.

278. *Reproduction* measurements including sexual maturity, time to first brood, brood size, sex ratio, and offspring produced (total and average per female), described in detail as follows:

- *Time to sexual maturity* is calculated as the duration, in days, of the interval between the initiation of the test (F0 mysids), or the release of the F1 mysids, and the appearance of structures defining maturity. Maturity is defined as the appearance of oostegites (marsupium) in the female and by the development of testes in the male. Some authors have used the presence of eggs in the oviduct as the indication of maturity in female mysids. Time to sexual maturity is determined for F0 and F1 mysids.
- The *time to first brood* is the time, measured in days, from hatching of the test organisms to the release of the first brood of young. Time to first brood is determined for F0 and F1 mysids. *Time to second brood* is determined for F0 and F1 mysids. *Interbrood duration* is the time between the release of the first and second broods.
- The *total number of offspring* is the total number of young produced by each replicate population for F0 and F1 mysids.
- *Sex ratio*, the ratio of females to males in the replicate treatment populations is determined for F0 and F1 mysids.
- The percentage of females that are reproductively active is determined for the F0 and F1 generations.

10.5.3.2 Optional Biochemical Endpoints

279. Several metabolic measurements have been strongly associated with alterations in endocrine-related processes after exposure to sublethal concentrations of toxicants. However, some pre-validation studies may need to be conducted to allow inclusion of these endpoints (see section 6.4.3). Detailed descriptions and methods for measuring those that should be considered for inclusion are presented in Section 10.5.3.

280. *Metabolic disruption* occurs as stress induces changes in the substrates used in metabolism. It is determined by measuring the O:N ratios of the test organisms. This ratio indicates the relationship between the amount of oxygen consumed by an organism to the amount of nitrogen excreted, and shows the relative role protein catabolism plays in the organism's energy budget (Carr et al. 1985; McKenney 1985). Changes in the O:N ratio measured among test mysids is a sensitive indicator that could provide for the relatively early detection of reproductive impacts by contaminants.

281. Disruption in steroid metabolism by EDCs can be determined by studying metabolic elimination of testosterone by mysids after exposure to the test compounds. Difference in metabolic by-products such as glucose conjugation, sulfate conjugation, hydroxylated and reduced/dehydrogenated metabolites of ¹⁴C-labeled testosterone in mysids exposed to sublethal concentrations of the test compounds can be measured. Different EDCs can affect testosterone metabolism in varying ways. For example, DES increased glucose conjugation, but did not affect sulfate conjugation, whereas 4NP reduced both of these elimination processes (Baldwin et al. 1995, 1997, 1998). Therefore, tests should not rely on measurements of only one by-product. Recent work with mysids (Verslycke et al. 2002) indicated that these measurements will be useful in studies of the effects on potential EDCs on that group.

282. *Vitellogenin induction* in crustaceans is probably controlled by ecdysteroids. However, whether or not this is true for mysids is not known. Differences in vitellogenin production among treated and nontreated mysids could provide evidence of endocrine system disruption and should be explored during pre-validation studies.

283. *Cytochrome P450 enzyme levels* may be affected by exposure of the crustaceans to EDCs. Measurements of differences in CYP levels between treated and nontreated mysids could provide direct evidence of disruption of steroid molting hormone levels.

284. *Blood glucose levels* in crustaceans are regulated by crustacean hyperglycemic hormone produced in the sinus gland. Changes in blood glucose levels in mysids exposed to potential EDCs could indicate disruption of hormonal activity other than that associated with molting or reproduction.

10.5.4 Exposure protocol

285. Because there is no validated two-generation mysid toxicity test, the exposure protocol selected for the EDC mysid testing is one based primarily on OECD (2004a), but also includes suggested improvements derived from the OPPTS protocol (EPA 1996), ASTM (1997), and Lussier et al. (1988), among other procedures. The suggested protocol and notes about some of the conditions are presented in Table 9.

286. One concern regarding the OECD design is the low minimum number of replicates (three) recommended by the guideline. The restriction of the test to three replicates may be a result of "spatial" constraints imposed by the relatively large test chambers and test volume (21 L) required to house the individual "baskets" used in the design. The often high variability in the response of organisms to potential EDCs may mask the ability of such a design to detect differences among treatments.

Table 9. Mysid two-generation toxicity test conditions recommended for conducting tests on potential endocrine disrupting chemicals

	Recommended Protocol	Notes
Test Species:	<i>Americamysis bahia</i> (Molenock 1969) Mysids used in test must originate from laboratory cultures	Restricted species to <i>A. bahia</i>; see text (OECD 2004a)
Holding Conditions:	Hold at conditions similar to test or acclimate gradually to test conditions (Temperature at 1 °C/24 h; salinity at ≤5%/24 h) Source for holding/culture water must be the same as for dilution water Holding facility should have same background colors and lighting intensities as testing areas Facility should be well ventilated and free of fumes that could affect test organisms Flow through or recirculating system; latter with ability to filter water as necessary 14 h light:10 h dark, or 16 h light:8 h dark, with 15–30 min transition period Gentle aeration Feed excess <i>Artemia</i> from reference cysts for which fatty acid content is known; # 150/mysid/d; may supplement with algae or other food	OECD (2004a); EPA (1996); ASTM (1997). Removed ASTM restriction on holding tank size to allow more flexibility. Feeding ration from OECD (2004a) and ASTM (1997).
Definitive Test		
Test organism age:	≤24 h	EPA (1996); OECD (2004a)
Duration:	7 days after mean day of second brood release in controls Expected duration could be 60 d or longer	OECD (2004a); time estimate is based on expectations for controls; treatments may be delayed in responses so test should extend beyond control response
Test Material:	Reagent grade chemical or better	EPA (1996); NS ^a in OECD (2004a)
Reproductive and Development Endpoints:	Survival (F0, F1) Growth—dry weight (Days 0, 7, 14; F0, F1) Time to Maturation (F0, F1) Time to First Brood Release (F0, F1) Time to Second Brood Release (F0, F1) Interbrood duration (F0, F1) Number of young per female in each of first two broods (F0, F1) Total number of young per brood in each replicate population (F0, F1) Total number of young in the first two broods in each replicate population (F0, F1) Sex ratio (F0, F1) Percentage of females that are reproductively active (F0, F1)	See Section 6.4 for descriptions of endpoints.
Optional Biochemical Endpoints:	Metabolic Disruption (F0, F1) Steroid Metabolism (F0, F1) Vitellogenin Induction (F0, F1) Cytochrome P450 Enzymes (F0, F1) Blood Glucose (F0, F1)	See Sections 6.4 and 11.0 for description of endpoints and data gaps. Biochemical endpoints should be considered for inclusion as they may provide information not available from reproductive and growth endpoints
Number of Treatments:	Minimum 5 concentrations, plus control (add solvent control if necessary)	OECD (2004a); EPA (1996)

	Recommended Protocol	Notes
Concentration Series:	Use series in which concentrations are stepped down from the highest by a factor of at least 2.0. Use range-finding test (described below) or results from acute test to determine definitive test concentrations. Highest concentration should equal the lowest concentration causing an adverse effect in acute test or 1/10 LC ₅₀ (EC ₅₀)	OECD (2004a); EPA (1996)
Dilution Water: Solvent:	Natural (<20-µm-filtered) or reconstituted seawater acceptable if mysids can survive and reproduce in it; at test initiation, dilution water should have: DO: 80% – 100% saturation Salinity: 18‰ – 22‰ pH: 7.6 – 8.2 With cationic test material, TOC ≤5 mg/L If solvent used, ≤50 µL/L; triethylene glycol recommended as solvent.	OECD (2004a); EPA (1996)
Flow Conditions:	Continuous or intermittent flow through	OECD (2004a); EPA (1996); ASTM (1997)
Delivery System: Flow Rate: Calibration limit: Calibration/Check:	Proportional diluter 90% exchange every 8–12 h for continuous system; 70% exchange every 5 h for intermittent system <10% variation in flow/chamber/time Prior to test system must be functioning well for 48 h; check twice daily	OECD (2004a); EPA (1996) Calibration limit recommended from EPA as OECD does not specify
# Replicates:	3 (minimum); more, if possible, are highly recommended	The number of replicates used in various protocols and tests ranges from 2 to 8. OECD (2004a) suggests a minimum of 3 replicates, however, this number is not likely to provide sufficient power for the test. EPA (1996) suggests at least 5. Although it may be desirable to determine the number of replicates based on the expected variation, desired detection limit, and selected power as suggested by ASTM (1997), the number selected may be constrained by the capacity of the diluter apparatus.
Test Chamber: Cover:	≥ 98 cm × 14 cm × 15 cm tank to hold retention baskets of 15-cm glass Petri dish with attached 15-cm high nylon mesh screen (210 µm mesh) May be advisable	OECD (2004a) Test chamber may be covered and have a screened overflow port.
Test Volume:	21 L	OECD (2004a)
Organisms/replicate	≥15 (≥45/treatment) for F0 generation Note: each replicate also requires 15 mysids for growth for a total of 30 mysids per replicate	OECD (2004a)

	Recommended Protocol	Notes
Initiation Notes:	Collect 8 groups of 5 Day 0 mysids; rinse briefly in deionized water; place into tared weighing boats; dry at 60°C for 24 h, weigh to nearest 0.1 µg Two retention baskets in each replicate tank—one for survival/reproduction endpoints, one for growth endpoint; place 15 mysids into each basket	OECD (2004a)
Test Conditions:		
Light:	Wide-spectrum fluorescent bulbs	OECD (2004a)
Photoperiod:	14 h light:10 h dark, with 15–30 min transition period	EPA (1996); OECD (2004a)
Temperature:	Test average: 25 °C ± 1 °C Day-to-Day: 25 °C ± 3 °C	EPA (1996); OECD (2004a); higher temperature may decrease embryo and larvae survival (Wortham-Neal and Price 2002)
pH:	7.6–8.2	OECD (2004a)
Dissolved Oxygen:	≥4.9 mg/L (=67% saturation at test conditions)	OECD (2004a)
Aeration:	None, unless DO <4.9 mg/L	
Salinity:	20‰ ± 2‰	OECD 2004a; EPA (1996) suggests 20 ‰ ± 3 ‰; note: higher salinity improves growth and reproduction (Lussier et al. 1988; McKenney and Celestial 1995; McKenney 1996; ASTM 1997)
Monitoring:		
Test Concentration	Stock Solution and test concentrations on Days 0, 7, 14, 21, 28, etc., and last day of test	EPA (1996); OECD (2004a)
WQ Frequency:	Salinity, temperature daily in one replicate chamber per concentration pH at start and end of test and weekly in control, include highest concentration DO in at least one test chamber per concentration at test initiation and termination; and three times per week during the exposures	NS in OECD (2004a); EPA (1996) suggests weekly for all parameters, may not be adequate for salinity and temperature; DO should be monitored frequently because aeration is not supplied unless the concentration drops below 4.9 mg/L
Observation Frequency:	Daily; Assess survival Watch for developing embryos Count F1 mysids daily until 7 days after mean day of second brood (F1 ¹) release Note date of appearance of marsupial pouch	OECD (2004a)
Feeding:	24-h old <i>Artemia</i> nauplii daily at rate of: Days 1–3: 1800/basket Days 4–6: 2250/basket Days 7–9: 2700/basket Days 10–12: 3150/basket Days >12: 3600/basket Brood Cups: 900/cup throughout test Dead brine shrimp should be removed daily before feeding occurs.	OECD (2004a); based on McKenney (1987)
Other Monitoring Notes:	Day 7 and Day 14 remove 8 groups of 5 mysids from growth baskets in each replicate for dry weight determinations (F0, F1)	OECD (2004a)

	Recommended Protocol	Notes
Data Collection/ Termination Notes:	Terminate test 7 days after mean day of second brood release in control F0 Count live F0 mysids and determine gender No dry weight determination at termination	OECD (2004a) Data collection points vary according to generation. See text for details.
Test Validity Criteria:	A test is valid if General test requirements, including water quality requirements, are met ≥70% control survival ≥75% control females (F0 or F1) produce young ≥8 average number of young/female in first two broods of controls	OECD (2004a)
	Range-Finding Test	
Concentration Series # samples test volume test containers # animals/sample duration	Widely spaced; e.g., 1, 10, 100 mg/L 1 400 mL 500-mL wide mouth jar Minimum 10/concentration 48–96 h	EPA (1996); OECD (2004a) Test volume, container size, numbers of animals, and duration are not specified in OECD (2004a) or EPA (1996); items listed here are suggestions; test should run long enough to allow estimation of test concentrations
Notes:	Perform test in manner similar to intended definitive test; report results along with those of definitive test	OECD (2004a)
Test Validity Criteria:	Response should be sufficient to allow estimation of appropriate definitive test concentration series	NS in EPA (1996) or OECD (2004a)
	Reference Toxicant Test	
Toxicant Concentration Series # reps test volume test containers # animals/rep	Copper sulfate 0, 150, 200, 300, 400 µg/L Cu 3 400 mL 500-mL wide mouth jar 10	NS in EPA (1996) or OECD (2004a); the approach here is suggested if a reference toxicant test is to be included to characterize the sensitivity of the test population
Termination Notes:	Count surviving mysids	NS in EPA (1996) or OECD (2004a)
Test Validity Criteria:	90% survival in controls; data sufficient to calculate LC ₅₀	NS in EPA (1996) or OECD (2004a)

^a NS, not specified.

10.5.5 Results and reporting

10.5.5.1 Interpretation of Results

287. Testing a matrix spiked with known concentrations of contaminants can be used to establish cause-and-effect relationships between chemicals and biological responses. Results of toxicity tests with test materials spiked into sediments or water at different concentrations may be reported in terms of an LC_x, an EC_x, an IC_x, or as an NOEC or LOEC. The most common techniques for statistical analysis of data sets include graphical displays, pairwise comparisons, trend analysis, analysis of variance (ANOVA), multiple regression techniques, and nonlinear dose-response analysis (ASTM 1997, Appendix). Graphical displays should be produced every time a test is performed to assess the structure of the data and reveal unanticipated relationships, influential observations, or anomalous data points (ASTM 1997, Appendix).

288. The assumptions of normality and homogeneity of variance should be assessed and appropriate data transformations performed if necessary. Nonparametric analysis can be used when assumptions

cannot be met. Some endpoints, such as the proportion surviving, can be transformed by the arcsine-square root, whereas others, such as growth and reproduction, can be transformed by logarithmic methods.

289. Some experiments are set up to compare more than one treatment with a control, whereas others compare the treatments with one another either assuming a trend or all pair-wise comparisons. The basic design is similar. After the applicable comparisons are determined, the data must be tested for normality and whether the variances of the treatments are equal to determine whether parametric or nonparametric statistics are appropriate. If normality of the data and equal variances are established, then ANOVA can be performed to address the hypothesis that all the treatments including the control are equal (EPA 2000). Williams' parametric test of ordered alternatives (Williams 1971) or the Jonckheere-Terpstra nonparametric trend test (Jonckheere 1954; Terpstra 1952) are used to determine the significance of a specified trend in the response associated with the treatments and estimate the NOEC and LOEC.

290. Commonly used approaches to assess the dose response using regression techniques are the graphical and linear interpolation method, probit analysis, trimmed Spearman-Kärber, and other nonlinear regression models. In general, results from these methods should yield similar estimates. Data for at least five test concentrations and the control should be available to calculate an LC₅₀, although some methods can be used with fewer concentrations. Survival in the lowest concentration must be at least 50%, and an LC₅₀ or EC₅₀ should not be calculated unless at least 50% of the organisms respond in at least one of the serial dilutions. When less than a 50% response occurs in the highest test concentration, the LC₅₀ or EC₅₀ is expressed as greater than the highest test concentration.

10.5.5.2 Reporting Requirements

291. The report should contain all pertinent information that is suggestive or predictive of chronic toxicity. The record of the results of an acceptable test should include the following information either directly or by referencing available documents:

- Name of test and investigator(s), name and location of laboratory, and dates of start and end of test
- If applicable, source of test water or sediment, and method for collection, handling, shipping, storage, and disposal of sediment
- Source of test material, lot number if applicable, composition (identities and concentrations of major ingredients and impurities if known), known chemical and physical properties, and the identity and concentration(s) of any solvent used
- Source of the dilution water, its chemical characteristics, and a description of any pretreatment, and results of any demonstration of the ability of an organism to survive or grow in the water
- Source, history, and age of test organisms; source, history, and age of brood stock, culture procedures; and source and date of collection of test organisms, scientific name, name of person who identified or cultured the organisms and the taxonomic key used, age or life stage, means and ranges of weight or length, observed diseases or unusual appearance, treatments used, and holding procedures
- Source and composition of brine shrimp; concentrations of test material and other contaminants in the brine shrimp; any added supplements; procedure used to prepare food; and feeding methods, frequency and ration

- Description of the experimental design and test chambers, the depth and volume of solution in the chamber, the number of mysids, the number of replicates, the loading, the lighting, and test substance delivery system and the flow rate as volume additions per 24-h period
- Methods used for physical and chemical characterization of water or sediment and the measured concentrations of test substances in test chambers. Information should include schedule for obtaining samples for analysis, the results of the analysis of test concentrations
- Range including minimum, maximum and average of measured water quality parameters (dissolved oxygen, salinity, temperature, and pH)
- A table of the biological data for each test chamber for each treatment, including the control(s), in sufficient detail to allow independent statistical analysis, including such measurements as time to sexual maturity (F0, F1); length of time to first brood (F0, F1); length of time to second brood (F0, F1), the interbrood duration, the average, and respective confidence intervals, for dry weight of males and females at Day 7 and Day 14 (F0, F1); the cumulative number of young produced for the first two broods (F0, F1), the sex ratio of each replicate population (F0, F1), and the number of surviving F0 and F1 mysids.
- Definition(s) of the effects used to calculate LC₅₀ or EC₅₀, biological endpoints for tests, and a summary of general observations of other effects
- A summary table of data on survival, growth, and reproduction of mysids in each test chamber, treatment, and control that includes the mean, standard deviation, and range for each endpoint
- Methods used for and results of the statistical analyses of data
- Summary of general observations on other effects or symptoms
- The results of the analytical determinations of the stock and test concentrations
- Anything unusual about the test, any deviation from these procedures, and any other relevant information
- Published reports should contain enough information to clearly identify the methods used and the quality of the results.

11.0 DATA GAPS, RESEARCH NEEDS AND OTHER CONSIDERATIONS

11.1 SIGNIFICANT DATA GAPS

292. Most of the testing with mysids has focused on short-term first generations studies. Specifics regarding carrying these tests for longer duration and incorporating a second generation need to be evaluated. For example, which F1 cohorts should be used for producing F2s, the length of time for observing the parent and offspring generations, and the appropriate performance criteria for the validation of a successful test all need to be determined. These issues could be addressed during pre-validation studies.

293. In addition, the necessity of a two-generation exposure duration should be assessed. Is the production of F2 offspring a more sensitive indicator than production of F1 offspring? Likewise, is the production of F2 offspring alone adequate or would survival or sex ratio or other endpoints be more sensitive still? Pre-validation studies are needed to determine the most cost-efficient and sensitive test duration and endpoints to be included in an optimized protocol suitable for progressing through full validation.

294. The goal of designing and conducting detailed chronic toxicity tests with mysids would be to determine whether specific endpoint responses can be determined for different classes of compounds that affect ecdysteroids, juvenile hormone analogs, vertebrate androgens, vertebrate estrogens, or other hormonal axes. Table 10 is an example of the type of information that might be obtained from such studies.

Table 10. Measurement of effects of three classes of hormones

	O:N Ratio	Survival	Growth	Reproduction				Steroid Metabolism
				Time to first Brood	Brood Size	Offspring Viability	Sex Ratio	
Ecdysteroid Agonist								
Ecdysteroid Antagonist								
Juvenile Hormone Analog Agonist								
Juvenile Hormone Analog Antagonist								
Vertebrate Androgen Agonist								
Vertebrate Androgen Antagonist								
Vertebrate Estrogen Agonist								
Vertebrate Estrogen Antagonist								
Other								

11.2 RESEARCH NEEDS

295. Because growth and development endpoints could be affected directly or indirectly by a variety of stress factors, such as environmental and biological, as well as chemical, it can be difficult to attribute the effects to a specific causal agent or mechanism with certainty. Therefore, biochemical studies must be included in the experimental regime to help verify that the observed endpoints resulted from disturbance of hormone systems. However, pre-validation studies about these biochemical metrics as they relate to mysids still needs to be done. For example, detailed mechanistic studies should be conducted using mysids to determine whether observed endpoint effects are caused by EDCs. Such studies could include receptor blocking/binding studies, endpoint response to specific classes of compounds, and enzyme studies, for example.

296. Detection of alterations in *steroid metabolism* could provide evidence supporting the impact of a potential EDC on mysid endocrine systems. Studies of testosterone metabolism in daphnid crustaceans showed that differences in glucose conjugation, sulfate conjugation, hydroxylated, and reduced/dehydrogenated metabolites of ¹⁴C-labeled testosterone exposed to test compounds versus control treatments provided evidence for disruption of endocrine processes. The general procedure (e.g., Baldwin et al. 1998) for measuring testosterone metabolites after exposure to test treatments is to place mysids in small containers that contain solutions having the same concentration of test substance to which they were exposed and to which radio-labeled testosterone has been added. Mysids are then homogenized on ice in distilled water, centrifuged, and the supernatant collected to estimate the soluble protein concentration.

The soluble protein values are used to normalize rates of testosterone metabolism among treatments. Details of the methods that should be applied to the mysid toxicity tests need to be developed.

- The general procedure for quantifying *glucose- and sulfate-conjugation* metabolites is by hydrolyzation with β -glucosidase (glucose conjugates) or sulfatase (sulfate conjugates) followed by thin layer chromatography (e.g., Baldwin et al. 1998).
- The general procedure for quantifying *hydroxylated and oxido-reduced/dehydrogenated* testosterone metabolites is by ethyl acetate extraction, steam evaporation, followed by thin layer chromatography and quantification via scintillation spectrophotography and comparison to known standards (e.g., Baldwin et al. 1998).

297. Changes in *cytochrome P450 enzymes*, which function in the detoxification of many exogenous and endogenous compounds, may be associated with disruption of the hormonally-regulated molting process and are therefore appropriate to measure in EDC studies. The general procedure for determining levels of cytochrome P450 in mysids involves homogenization of whole animals, centrifugation, and the collection of the resulting supernatant. Quantification of the levels occurs via gel electrophoresis. Snyder and Mulder (2001) described a specific method for determining CYP45 (a family of CP450 enzymes) levels in daphnid crustaceans.

298. It would be necessary to develop biomarkers for the selected species; promising biomarkers could be the induction of vitellogenesis in males, and the inhibition of aromatase in females (Depledge unpublished, cited in Depledge and Billinghamurst 1999). As more evidence for endocrine disruption responses is gathered from experimental research, mechanistic studies would be required to determine the specific ways in which chemicals can disturb hormones. That is, it becomes important to distinguish between endocrine disruption and metabolic toxicity, and to determine which is a primary, and which is a secondary effect. The final step in the strategy would be to conduct field surveys to detect and confirm that endocrine disruption effects occur *in situ* (Depledge and Billinghamurst 1999).

11.3 IMPLEMENTATION CONSIDERATIONS

299. To implement the recommended protocol, regulatory and other legal requirements must be met, and long-term goals for public health and safety should be kept in mind. Following the general principles put forth by ICCVAM, pre-validation studies should be initiated. None of the biochemical endpoints suggested in the recommended protocol have been through a validation process and have not been routinely used by laboratories. Based on available information, selection of the appropriate biochemical endpoints would be difficult. It is therefore recommended that a pre-validation study be performed that would evaluate the biochemical endpoints as a marker for endocrine disruption in mysids. Validation of the study design through interlaboratory comparisons should be conducted once preferred endpoints have been identified using compounds that span the possible endocrine effects, including strong and weak androgen receptor agonists and antagonists, estrogen receptor agonists and antagonists, and thyroid agonists and antagonists

11.4 ANIMAL WELFARE

300. Legislation governing the care and use of laboratory animals in the United States is contained in the Animal Welfare Act, passed in 1966 and amended in 1970, 1976, and 1985. The Animal Welfare Act covers all warm-blooded animals except mice, rats, birds, and horses, and other farm animals when they are not used for research. It spells out requirements for veterinary care, adequate food and water, protection from temperature extremes, shelter from outdoor elements, sanitation, and record keeping. Because mysids are cold-blooded, there are no current laws governing the culturing or testing of these

organisms. Nationally recognized protocols such as ASTM E1191 (ASTM 1997) and OPPTS 850.1350 (EPA 1996) will be followed for testing. These protocols state that all mysids used in testing must be destroyed at the end of the test using humane methods. One humane method that may be used is placing the mysids into a solution of oxygenated MS 222 (euthanizing agent).

11.5 RECOMMENDED EQUIPMENT AND CAPABILITY

301. To ensure interlaboratory comparability and the general accessibility of the protocol to a broad number of testing laboratories, the following essential equipment and or capabilities are recommended to properly conduct the mysid chronic test:

- Diluters or other flow-through systems with capability of meeting flow and dilution precision requirements
- A high-quality microscope system for anatomical measurements (e.g., male-female characteristics, eyestalk structure); a low-intensity dual-channel laser-scanning confocal microscope is recommended as one of the best systems to use
- Analytical measurement capacity – chemical analysis of test compounds
- Protein testing – electrophoresis, chromatography capability, vitellin-probes such as biodipy
- Culture and maintenance of mysids (laboratories can purchase <24- h-old mysids from commercial suppliers).

11.6 TESTING WITH NON-NATIVE SPECIES

302. Interest in the occurrence and impacts of introduced marine and estuarine species has increased since the 1980s. Recently, Ruiz et al. (2000) documented that about 300 introduced species (invertebrates and algae) have become established in United States coastal waters. Introduced species are significant stressors to coastal ecosystems (Ruiz et al. 1999) and the damage they cause to coastal ecosystems is well documented (e.g., Grosholz and Ruiz 1995, 1996). Governmental initiatives have been implemented to reduce the likelihood of new introductions (Federal Register 1999). Because the primary species recommended for the EDC testing program is not indigenous to some of the geographic regions where testing may occur, all testing laboratories should take appropriate precautions to reduce the possibility that an accidental introduction to a local ecosystem could occur. All test mysids should be destroyed in an appropriate manner at the completion of each test (ASTM 1997).

12.0 REFERENCES

1. Anctil, M. 2000. Evidence for Gonadotropin-Releasing Hormone-Like Peptides in a Cnidarian Nervous System. *General and Comparative Endocrinology* 119(3):317-328.
2. Andersen, H.R., Wollenberger, L., Halling-Sorensen, B., and Kusk, K.O. 2001. Development of Copepod Nauplii to Copepodites: a Parameter for Chronic Toxicity Including Endocrine Disruption. *Environmental Toxicology and Chemistry* 20(12):2821-2829.

3. Andries, J.C. 2001. Endocrine and Environmental Control of Reproduction in Polychaeta. *Canadian Journal of Zoology Revue Canadienne de Zoologie* 79(2):254-270.
4. Arnold, S.F., Klotz, D.M., Collins, B.M., Vonier, P.M., Guillette, L.J. Jr., and McLachlan, J.A. 1996. Synergistic Activation of Estrogen Receptor with Combinations of Environmental Chemicals. *Science* 272(5267):1489-1492.
5. ASTM (American Society for Testing and Materials). 1997. *Standard Guide for Conducting Life-cycle Toxicity Tests with Saltwater Mysids*. E 1191-97. American Society for Testing and Materials. West Conshohocken, Pennsylvania.
6. ASTM (American Society for Testing and Materials). 2000. *Standard Guide for Conducting Sediment Toxicity Tests with Polychaetous Annelids*. E 1611-00. American Society for Testing and Materials. West Conshohocken, Pennsylvania.
7. ASTM (American Society for Testing and Materials). 2004. *Standard Guide for Conducting Daphnia magna Life-cycle Toxicity Tests*. E 1693-97 (Reapproved 2004). American Society for Testing and Materials. West Conshohocken, Pennsylvania.
8. Axiak, V., Micallef, D., Muscat, J., Vella, A., and Mintoff, B. 2003. Imposex as a Biomonitoring Tool for Marine Pollution by Tributyltin: Some Further Observations. *Environment International* 28:743-749.
9. Baldwin, W.S., Bailey, R., Long, K.E., and Klaine, S. 2001. Incomplete Ecdysis is an Indicator of Ecdysteroid Exposure in *Daphnia magna*. *Environmental Toxicology and Chemistry* 20(7):1564-1569.
10. Baldwin, W.S., Graham, S.E., Shea, D., and LeBlanc, G.A. 1998. Altered Metabolic Elimination of Testosterone and Associated Toxicity Following Exposure of *Daphnia magna* to Nonylphenol Polyethoxylate. *Ecotoxicology and Environmental Safety* 39(2):104-111.
11. Baldwin, W.S., Graham, S.E., Shea, D., and LeBlanc, G.A. 1997. Metabolic Androgenization of Female *Daphnia magna* by the Xenoestrogen 4-Nonylphenol. *Environmental Toxicology and Chemistry* 16(9):1905-1911.
12. Baldwin, W.S., Milam, D.L., and LeBlanc, G.A. 1995. Physiological and Biochemical Perturbations in *Daphnia magna* Following Exposure to the Model Environmental Estrogen Diethylstilbestrol. *Environmental Toxicology and Chemistry* 14(6):945-952.
13. Bechmann, R.K. 1999. Effect of the Endocrine Disrupter Nonylphenol on the Marine Copepod *Tisbe battagliai*. *The Science of the Total Environment* 233(1-3):33-46.
14. Bejarano, A. C. and Chandler, G. T. 2003. Reproductive and Developmental Effects of Atrazine on the Estuarine Meiobenthic Copepod *Amphiascus tenuiremis*. *Environmental Toxicology and Chemistry* 22(12):3009-3016.
15. Benoit, D.A., Sibley, P.K., Juenemann, J.L., and Ankley, G.T. 1997. *Chironomus tentans* Life Cycle Test: Design and Evaluation for Use in Assessing Toxicity of Contaminated Sediments. *Environmental Toxicology and Chemistry* 16:1165-1176.

16. Billinghamurst, Z., Clare, A.S., and Depledge, M.H. 2001. Effects of 4-N-Nonylphenol and 17 β -Oestradiol on Early Development of the Barnacle *Elminius modestus*. *Journal of Experimental Marine Biology and Ecology* 257(2):255-268.
17. Billinghamurst, Z., Clare, A.S., Fileman, T., McEvoy, M., Readman, J., and Depledge, M.H. 1998. Inhibition of Barnacle Settlement by the Environmental Oestrogen 4-Nonylphenol and the Natural Oestrogen 17 β -Oestradiol. *Marine Pollution Bulletin* 36(10):833-839.
18. Billinghamurst, Z., Clare, A.S., Matsumura, K., and Depledge, M.H. 2000. Induction of Cypris Major Protein in Barnacle Larvae by Exposure to 4-N-Nonylphenol and 17 β -Oestradiol. *Aquatic Toxicology* 47(3-4):203-212.
19. Block, D.S., Bejarano, A.C., and Chandler, G.T. 2003. Ecdysteroid Concentrations Through Various Life-Stages of the Meiobenthic Harpacticoid Copepod, *Amphiascus tenuiremis* and the Benthic Estuarine Amphipod, *Leptocheirus plumulosus*. *General and Comparative Endocrinology* 132(1):151-160.
20. Bodar, C.Wm., Voogt, P.A., and Zandee, D.I. 1990. Ecdysteroids in *Daphnia magna*: Their Role in Molting and Reproduction and Their Levels Upon Exposure to Cadmium. *Aquatic Toxicology* 17(4):339-349.
21. Borst, D.W., Laufer, H., Landau, M., Chang, E.S., Hertz, W.A., Baker, F.C., and Schooley, D.A. 1987. Methyl Farnesoate and Its Role in Crustacean Reproduction and Development. *Insect Biochemistry* 17:1123-1127.
22. Bousfield, E.L. 1973. *Shallow-Water Gammaridean Amphipoda of New England*. Cornell University Press. Ithaca, NY.
23. Brandt, O.M., Fujimura, R.W., and Finlayson, B.J. 1993. Use of *Neomysis mercedis* (Crustacea, Mysidacea) for Estuarine Toxicity Tests. *Transactions of the American Fisheries Society* 122(2):279-288.
24. Breitholtz, M. and Bengtsson, B.E. 2001. Oestrogens Have No Hormonal Effect on the Development and Reproduction of the Harpacticoid Copepod *Nitocra spinipes*. *Marine Pollution Bulletin* 42(10):879-886.
25. Breitholtz, M., Wollenberger, L., and Dinan, L. 2003. Effects of Four Synthetic Musks on the Life Cycle of the Harpacticoid Copepod *Nitocra spinipes*. *Aquatic Toxicology* 63(2):103-118.
26. Brown, R.J., Conradi, M., and Depledge, M.H. 1999. Long-Term Exposure to 4-Nonylphenol Affects Sexual Differentiation and Growth of the Amphipod *Corophium volutator* (Pallas, 1766). *The Science of the Total Environment* 233(1-3):77-88.
27. Brusca, R.C. and Brusca, G.J. 1990. *Invertebrates*. Sinauer Associates, Inc. Sunderland, Massachusetts.
28. Buratini, S.V., Bertoletti, E., and Zagatto, P.A. 2004. Evaluation of *Daphnia similis* as a Test Species in Ecotoxicological Assays. *Bulletin of Environmental Contamination and Toxicology* 73:878-882

29. Candia Carnevali, M.D., Bonasoro, F., Patrino, M., Thorndyke, M.C., and Galassi, S. 2001a. PCB Exposure and Regeneration in Crinoids (Echinodermata). *Marine Ecology Progress Series* 215:155-167.
30. Candia Carnevali, M.D., Galassi, S., Bonasoro, F., Patrino, M., and Thorndyke, M.C. 2001b. Regenerative Response and Endocrine Disrupters in Crinoid Echinoderms: Arm Regeneration in *Antedon mediterranea* After Experimental Exposure to Polychlorinated Biphenyls. *Journal of Experimental Biology* 204(5):835-842.
31. Carr, R.S., Montagna, P.A., and Kennicutt, M.C. Jr. 1998. Sediment Quality Assessment of Storm Water Outfalls and Other Selected Sites in the Corpus Christi Bay National Estuary Program Study Area. CCBNEP-32. Corpus Christi Bay National Estuary Program, Texas.
32. Carr, R.S., Williams, J.W., Saksa, F.I., Buhl, R.L., and Neff, J.M. 1985. Bioenergetic Alterations Correlated With Growth Fecundity and Body Burden of Cadmium for Mysids *Mysidopsis bahia*. *Environmental Toxicology and Chemistry* 4(2):181-188.
33. Castro, B.B., Guilhermino, L., and Ribeiro, R. 2003. In Situ Bioassay Chambers and Procedures for Assessment of Sediment Toxicity With *Chironomus riparius*. *Environmental Pollution* 125(3):325-335.
34. Chandler, G.T., Cary, T.L., Volz, D.C., Walse, S. S., Ferry, J. L., and Klosterhaus, S. L. 2004. Fipronil Effects on Estuarine Copepod (*Amphiascus tenuiremis*) Development, Fertility, and Reproduction: a Rapid Life-Cycle Assay in 96-Well Microplate Format. *Environmental Toxicology and Chemistry* 23(1):117-124.
35. Chang, E.S. 1993. Comparative Endocrinology of Molting and Reproduction: Insects and Crustaceans. *Annual Reviews of Entomology* 38:161-180.
36. Chang, E.S. 1997. Chemistry of Crustacean Hormones that Regulate Growth and Reproduction. Pp 163-178. In: Fingerman, M., Nagabhushanam, R., and Thompson, M.F. (eds). *Recent Advances in Marine Biotechnology Vol. 1: Endocrinology and Reproduction*. Science Publishers, Inc., Enfield, NH.
37. Chang, E.S., Chang, S.A., and Mulder, E.P. 2001. Hormones in the Lives of Crustaceans: an Overview. *American Zoologist* 41(5):1090-1097.
38. Chapman, G.A., Denton, D.L., and Lazorchak, J.M. (eds.). 1995. Short-Term Methods for Estimating the Chronic Toxicity of Effluents and Receiving Waters to West Coast Marine and Estuarine Organisms. First edition. EPA/600/R-95-136. U.S. Environmental Protection Agency, Office of Prevention, Pesticides and Toxic Substances. Washington, DC.
39. Chapman, P.F., Crane, M., Wiles, J., Noppert, F., and McIndoe, E. 1996. Improving the Quality of Statistics in Regulatory Ecotoxicity Tests. *Ecotoxicology* 5:169-186.
40. Charmantier, G., Charmantier-Daures, M., and Van Herp, F. 1997. Hormonal regulation of growth and reproduction in crustaceans. Pp 109-161. In: Fingerman, M., Nagabhushanam, R., and Thompson, M.F. (eds). *Recent Advances in Marine Biotechnology Vol. 1: Endocrinology and Reproduction*. Science Publishers, Inc., Enfield, NH.
41. Chigbu, P. 2004. Assessment of the Potential Impact of the Mysid Shrimp, *Neomysis mercedis*, on *Daphnia*. *Journal of Plankton Research* 26(3):295-306.

42. Clarke, G.M. 1992. Fluctuating Asymmetry: A Technique for Measuring Developmental Stress of Genetic and Environmental Origin. *Acta Zoologica Fennica* 191:31-35.
43. Clarke, G.M. 1993. Fluctuating Asymmetry of Invertebrate Populations as a Biological Indicator of Environmental Quality. *Environmental Pollution* 82(2):207-.
44. Cochran, W.G. and Cox, G.M. 1957. *Experimental Designs, Second Edition*. Wiley. New York, NY.
45. Coeurdassier, M., De Vaufleury, A., Scheifler, R., Morhain, E., and Badot, P.M. 2004. Effects of Cadmium on the Survival of Three Life-Stages of the Freshwater Pulmonate *Lymnaea stagnalis* (Mollusca : Gastropoda). *Bulletin of Environmental Contamination and Toxicology* 72(5):1083-1090.
46. Crane, M. and Newman, M.C. 2000. What Level of Effect Is a No Observed Effect? *Environmental Toxicology and Chemistry* 19:516-519.
47. Cripe, G.M., Carr, R.S., Foss, S.S., Harris, P.S., and Stanley, R.S. 2000. Effects of Whole Sediments from Corpus Christi Bay on Survival, Growth, and Reproduction of the Mysid, *Americamysis bahia* (Formerly *Mysidopsis bahia*). *Bulletin of Environmental Contamination and Toxicology* 64(3):426-433.
48. Cripe, G.M., McKenney, C.L., Jr., Hoglund, M.D., and Harris, P.S. 2003. Effects of Fenoxycarb Exposure on Complete Larval Development of the Xanthid Crab *Rhithropanopeus harrisi*. *Environmental Pollution* 125:295-299.
49. Crisp, T.M., Clegg, E.D., Cooper, R.L., Wood, W.P., Anderson, D.G., Baetcke, K.P., Hoffmann, J.L., Morrow, M.S., Rodier, D.J., Schaeffer, J.E., Touart, L.W., Zeeman, M.G., and Patel, Y.M. 1998. Environmental Endocrine Disruption: an Effects Assessment and Analysis. *Environmental Health Perspectives* 106:11-56.
50. Cuzin-Roudy, J. and Saleuddin, A.S.M. 1989. The Mysid *Siriella armata*, a Biological Model for the Study of Hormonal-Control of Molt and Reproduction in Crustaceans - a Review. *Invertebrate Reproduction & Development* 16(1-3):33-42.
51. Czech, P., Weber, K., and Dietrich, D.R. 2001. Effects of Endocrine Modulating Substances on Reproduction in the Hermaphroditic Snail *Lymnaea stagnalis* L. *Aquatic Toxicology* 53(2):103-114.
52. Daly, K.L. and Holmquist, C. 1986. A Key to the Mysidacea of the Pacific Northwest. *Canadian Journal of Zoology* 64:1201-1210.
53. Dauvin, J.-C., Iglesias, A., and Lorgeré, J.-C. 1994. Circalittoral Suprabenthic Coarse Sand Community from the Western English Channel. *Journal of the Marine Biological Association of the United Kingdom* 74:543-562.
54. Delbecque, J.P., Weidner, K., and Hoffman, K.H. 1990. Alternative Sites for Ecdysteroid Production in Insects. *Invertebrate Reproduction & Development* 18:29-42.
55. Depledge, M.H. and Billinghurst, Z. 1999. Ecological Significance of Endocrine Disruption in Marine Invertebrates. *Marine Pollution Bulletin* 39(1-12):32-38.

56. Domingues, P.M., Turk, P.E., Andrade, J.P., and Lee, P.G. 1999. Culture of the Mysid, *Mysidopsis almyra* (Bowman), (Crustacea: Mysidacea) in a Static Water System: Effects of Density and Temperature on Production, Survival and Growth. *Aquaculture Research* 30(2):135-143.
57. Domingues, P.M., Turk, P.E., Andrade, J.P., and Lee, P.G. 2001a. Effects of Different Food Items on the Culture of the Mysid Shrimp *Mysidopsis almyra* (Crustacea : Pericaridea) in a Static Water System. *Aquaculture International* 9(5):393-400.
58. Domingues, P.M., Turk, P.E., Andrade, J.P., and Lee, P.G. 2001b. Effects of Enriched Artemia Nauplii on Production, Survival and Growth of the Mysid Shrimp *Mysidopsis almyra* Bowman 1964 (Crustacea : Mysidacea). *Aquaculture Research* 32(7):599-603.
59. Domingues, P.M., Turk, P.E., Andrade, J.P., and Lee, P.G. 1998. Pilot-Scale Production of Mysid Shrimp in a Static Water System. *Aquaculture International* 6(5):387-402.
60. Draper, N.R. and Smith, H. 1981. *Applied Regression Analysis, Second Edition*. John Wiley and Sons. New York, NY.
61. Duan, Y., Guttman, S.I., and Oris, J.T. 1997. Genetic Differentiation Among Laboratory Populations of *Hyalella azteca*: Implications for Toxicology. *Environmental Toxicology and Chemistry* 16(4):691-695.
62. Duan, Y.H., Guttman, S.I., Oris, J.T., and Bailer, A.J. 2000a. Genetic Structure and Relationships Among Populations of *Hyalella azteca* and *H. montezuma* (Crustacea : Amphipoda). *Journal of the North American Benthological Society* 19(2):308-320.
63. Duan, Y.H., Guttman, S.I., Oris, J.T., and Bailer, A.J. 2000b. Genotype and Toxicity Relationships Among *Hyalella azteca*: I. Acute Exposure to Metals or Low Ph. *Environmental Toxicology and Chemistry* 19(5):1414-1421.
64. Duan, Y.H., Guttman, S.I., Oris, J.T., Huang, X.D., and Burton, G.A. 2000c. Genotype and Toxicity Relationships Among *Hyalella azteca*: II. Acute Exposure to Fluoranthene-Contaminated Sediment. *Environmental Toxicology and Chemistry* 19(5):1422-1426.
65. Dunham, J.S. and Duffus, D.A. 2002. Diet of Gray Whales (*Eschrichtius robustus*) in Clayoquot Sound, British Columbia, Canada. *Marine Mammal Science* 18(2):419-437.
66. EDSTAC (Endocrine Disruptor Screening and Testing Advisory Committee). 1998. *Final Report*. EPA/743/R-98/003. U.S. Environmental Protection Agency. Washington, D.C.
67. Emery, V.L., Moore, D.W., Gray, B.R., Duke, B.M., Gibson, A.B., Wright, R.B., and Farrar, J.D. 1997. Development of a Chronic Sublethal Sediment Bioassay using the Estuarine Amphipod, *Leptocheirus plumulosus* (Shoemaker). *Environmental Toxicology and Chemistry* 16(9):1912-1920.
68. Environment Canada. 1997b. Biological Test Method: Test for Survival and Growth in Sediment using the Freshwater Amphipod *Hyalella azteca*. EPS 1/RM/33.
69. Environment Canada. 1997a. Biological Test Method: Test for Survival and Growth in Sediment using the Larvae of Freshwater Midges *Chironomus tentans* or *Chironomus riparius*. EPS 1/RM/32.
70. Environment Canada. 1992. Biological Test Method: Test of Reproduction and Survival using the Cladoceran *Ceriodaphnia dubia*. EPS 1/RM/21.

71. EPA (U.S. Environmental Protection Agency). 1996. *Ecological Effects Test Guidelines OPPTS 850.1350 Mysid Chronic Toxicity Test. Public Draft*. EPA 712-C-96-120. U.S. Environmental Protection Agency, Office of Prevention, Pesticides and Toxic Substances. Washington, D.C.
72. EPA (U.S. Environmental Protection Agency). 2001. *Methods for Assessing the Chronic Toxicity of Marine and Estuarine Sediment-associated Contaminants with the Amphipod Leptocheirus plumulosus - First Edition*. EPA/600/R-01/020. U.S. Environmental Protection Agency, Office of Prevention, Pesticides and Toxic Substances. Washington, DC.
73. EPA (U.S. Environmental Protection Agency). 2002a. *Methods for Measuring the Acute Toxicity of Effluents and Receiving Waters to Freshwater and Marine Organisms. Fifth Edition*. EPA-821-R-02-012. U.S. Environmental Protection Agency, Office of Prevention, Pesticides and Toxic Substances. Washington, DC.
74. EPA (U.S. Environmental Protection Agency). 2000. *Methods for Measuring the Toxicity and Bioaccumulation of Sediment-Associated Contaminants with Freshwater Invertebrates*. EPA/600/R-99/064. U.S. Environmental Protection Agency, Office of Prevention, Pesticides and Toxic Substances. Washington, DC.
75. EPA (U.S. Environmental Protection Agency). 2002b. *Short-term Methods for Measuring the Chronic Toxicity of Effluents and Receiving Waters to Freshwater Organisms. Fourth Edition*. EPA-821-R-02-013. U.S. Environmental Protection Agency, Office of Prevention, Pesticides and Toxic Substances. Washington, DC.
76. EPA (U.S. Environmental Protection Agency). 2002c. *Short-term Methods for Measuring the Chronic Toxicity of Effluents and Receiving Waters to Marine and Estuarine Organisms. Third Edition*. EPA-821-R-02-014. U.S. Environmental Protection Agency, Office of Prevention, Pesticides and Toxic Substances. Washington, DC.
77. EPA/USACE (U.S. Environmental Protection Agency/U.S. Army Corps of Engineers). 1998. *Evaluation of Dredged Material Proposed for Discharge in Waters of the U.S. Testing Manual*. EPA-823-B-98-004. U.S. Environmental Protection Agency, Office of Water. Washington, D.C.
78. Evans, S.M. and Nicholson, G.J. 2000. The Use of Imposex to Assess Tributyltin Contamination in Coastal Waters and Open Seas. *The Science of the Total Environment* 258:73-80.
79. Fairs, N.J., Quinlan, P.T., and Goad, L.J. 1990. Changes in Ovarian Unconjugated and Conjugated Steroid Titters During Vitellogenesis in *Penaeus monodon*. *Aquaculture* 89:83-99.
80. Farrell, A.P., Kennedy, C.J., Wood, A., Johnston, B.D., and Bennett, W.R. 1998a. Acute Toxicity of a Didecyldimethylammonium Chloride-Based Wood Preservative, Bardac 2280, to Aquatic Species. *Environmental Toxicology and Chemistry* 17:1552-1557.
81. Farrell, A.P., Stockner, E., and Kennedy, C.J. 1998b. A Study of the Lethal and Sublethal Toxicity of Polyphase P-100, an Antisapstain Fungicide Containing 3-Iodo-2-Propynyl Butyl Carbamate (IPBC), on Fish and Aquatic Invertebrates. *Archives of Environmental Contamination and Toxicology* 35:472-478.
82. Feix, M. and Hoch, M. 2002. Phylogeny and Evolution of Hormone Systems. *Anesthesiologie Intensivmedizin Notfallmedizin Schmerztherapie* 37(11):651-658.

83. Fingerman, M. 1997. Crustacean Endocrinology: A Retrospective, Prospective, and Introspective Analysis. *Physiological Zoology* 70(3):257-269.
84. Fingerman, M. 1987. The Endocrine Mechanisms of Crustaceans. *Journal of Crustacean Biology* 7(1):1-24.
85. Fingerman, M., Jackson, N.C., and Nagabhushanam, R. 1998. Hormonally-Regulated Functions in Crustaceans as Biomarkers of Environmental Pollution. *Comparative Biochemistry and Physiology C: Pharmacology Toxicology & Endocrinology* 120(3):343-350.
86. Fischer, A. and Dorresteijn, A. 2004. The Polychaete *Platynereis dumerilii* (Annelida): a Laboratory Animal With Spiralian Cleavage, Lifelong Segment Proliferation and a Mixed Benthic/Pelagic Life Cycle. *BioEssays* 26(3):314-325.
87. Ford, A.T., Fernandes, T.F., Read, P.A., Robinson, C.D., and Davies, I.M. 2004b. The Costs of Intersexuality: A Crustacean Perspective. *Marine Biology* 145:951-957.
88. Ford, A.T., Fernandes, T.F., Rider, S.A., Read, P.A., Robinson, C.D., and Davies, I.M. 2004a. Endocrine Disruption in a Marine Amphipod? Field Observations of Intersexuality and Demasculinisation. *Marine Environmental Research* 58:169-173.
89. Garnacho, E., Peck, L.S., and Tyler, P.A. 2000. Variations Between Winter and Summer in the Toxicity of Copper to a Population of the Mysid *Praunus flexuosus*. *Marine Biology* 137:631-636.
90. Garnacho, E., Peck, L.S., and Tyler, P.A. 2001. Effects of Copper Exposure on the Metabolism of the Mysid *Praunus flexuosus*. *Journal of Experimental Marine Biology and Ecology* 265(2):181-201.
91. Gaudron, S.M. and Bentley, M.G. 2002. Control of Reproductive Behaviour in the Scale Worm *Harmothoe imbricata* (Annelida : Polychaeta : Polynoidae). *Invertebrate Reproduction & Development* 41(1-3):109-118.
92. Gentile, J.H., Gentile, S.M., Hoffman, G., Heltshe, J.F., and Hairston, N.J. 1983. Effects of a Chronic Mercury Exposure on Survival, Reproduction and Population Dynamics of *Mysidopsis bahia*. *Environmental Toxicology and Chemistry* 2(1):61-68.
93. Gentile, S.M., Gentile, J.H., Walker, J., and Heltshe, J.F. 1982. Chronic Effects of Cadmium on Two Species of Mysid Shrimp *Mysidopsis bahia* and *Mysidopsis bigelowi*. *Hydrobiologia* 93(1-2):195-204.
94. Gleason, T.R. and Nacci, D.E. 2001. Risks of Endocrine-Disrupting Compounds to Wildlife: Extrapolating from Effects on Individuals to Population Response. *Human and Ecological Risk Assessment* 7(5):1027-1042.
95. Gonzalez, E.R. and Watling, L. 2002. Redescription of *Hyaella azteca* from Its Type Locality, Vera Cruz, Mexico (Amphipoda : Hyaellidae). *Journal of Crustacean Biology* 22(1):173-183.
96. Goodwin, W.T., Horn, D.H.S., Karlson, P., Koolman, J., Nakanishi, K., Robbins, W.E., Siddall, J.B., and Takemoto, T. 1978. Ecdysteroids: a New Generic Term. *Nature* 272:122.

97. Gorokhova, E. 2002. Moulting Cycle and Its Chronology in *Mysis mixta* and *Neomysis integer* (Crustacea, Mysidacea): Implications for Growth Assessment. *Journal of Experimental Marine Biology and Ecology* 278(2):179-194.
98. Gorokhova, E. and Hansson, S. 2000. Elemental Composition of *Mysis mixta* (Crustacea, Mysidacea) and Energy Costs of Reproduction and Embryogenesis under Laboratory Conditions. *Journal of Experimental Marine Biology and Ecology* 246:103-123.
99. Goto, T. and Hiromi, J. 2003. Toxicity of 17 α -ethynylestradiol and Norethindrone, Constituents of an Oral Contraceptive Pill to the Swimming and Reproduction of Cladoceran *Daphnia magna*, With Special Reference to Their Synergetic Effect. *Marine Pollution Bulletin* 47(1-6):139-142.
100. Grosholz, E.D. and Ruiz, G.M. 1996. Predicting the Impact of Introduced Marine Species: Lessons from the Multiple Invasions of the European Green Crab *Carcinus maenas*. *Biological Conservation* 78(1-2):59-66.
101. Grosholz, E.D. and Ruiz, G.M. 1995. Spread and Potential Impact of the Recently Introduced European Green Crab, *Carcinus maenas*, in Central California. *Marine Biology* 122(2):239-247.
102. Gross, M.Y., Maycock, D.S., Thorndyke, M.C., Morritt, D., and Crane, M. 2001. Abnormalities in Sexual Development of the Amphipod *Gammarus pulex* (L.) Found Below Sewage Treatment Works. *Environmental Toxicology and Chemistry* 20(8):1792-1797.
103. Hagger, J.A., Fisher, A.S., Hill, S.J., Depledge, M.H., and Jha, A.N. 2002. Genotoxic, Cytotoxic and Ontogenetic Effects of Tri-*n*-Butyltin on the Marine Worm, *Platynereis dumerilii* (Polychaeta : Nereidae). *Aquatic Toxicology* 57(4):243-255.
104. Hahn, T., Liess, M., and Schulz, R. 2001. Effects of the Hormone Mimetic Insecticide Tebufenozide on *Chironomus riparius* Larvae in Two Different Exposure Setups. *Ecotoxicology and Environmental Safety* 49(2):171-178.
105. Hahn, T., Schenk, K., and Schulz, R. 2002. Environmental Chemicals With Known Endocrine Potential Affect Yolk Protein Content in the Aquatic Insect *Chironomus riparius*. *Environmental Pollution* 120(3):525-528.
106. Hahn, T. and Schulz, R. 2002. Ecdysteroid Synthesis and Imaginal Disc Development in the Midge *Chironomus riparius* as Biomarkers for Endocrine Effects of Tributyltin. *Environmental Toxicology and Chemistry* 21(5):1052-1057.
107. Hansen, F.T., Forbes, V.E., and Forbes, T.L. 1999. Effects of 4-*n*-Nonylphenol on Life-History Traits and Population Dynamics of a Polychaete. *Ecological Applications* 9(2):482-495.
108. Hardege, J.D. 1999. Nereid Polychaetes as Model Organisms for Marine Chemical Ecology. *Hydrobiologia* 402:145-161.
109. Harmon, V.L. and Langdon, C.J. 1996. A 7-D Toxicity Test for Marine Pollutants using the Pacific Mysid *Mysidopsis intii*. 2. Protocol Evaluation. *Environmental Toxicology and Chemistry* 15(10):1824-1830.
110. Hill, M., Stabile, C., Steffen, L.K., and Hill, A. 2002. Toxic Effects of Endocrine Disrupters on Freshwater Sponges: Common Developmental Abnormalities. *Environmental Pollution* 117(2):295-300.

111. Hollister, T.A., Heitmuller, P.T., Parrish, P.R. , and Dyar, E.E. 1980 . Studies to Determine Relationships Between Time and Toxicity of an Acidic Effluent and an Alkaline Effluent to Two Estuarine Species. In: *Aquatic Toxicology*. Pp. 251-265. American Society for Testing and Materials. Philadelphia, Pennsylvania.
112. Huber, J.T. 1998. The Importance of Voucher Specimens, With Practical Guidelines for Preserving Specimens of the Major Invertebrate Phyla for Identification. *Journal of Natural History* 32(3):367-385.
113. Huberman, A. 2000. Shrimp Endocrinology. A Review. *Aquaculture* 191(1-3):191-208.
114. Hunt, J.W., Anderson, B.S., Phillips, B.M., Tjeerdema, R.S., Puckett, H.M., and de Vlaming, V. 1999. Patterns of Aquatic Toxicity in an Agriculturally Dominated Coastal Watershed in California. *Agriculture, Ecosystems & Environment* 75:75-91.
115. Hunt, J.W., Anderson, B.S., Phillips, B.M., Tjeerdema, R.S., Puckett, H.M., Stephenson, M., Tucker, D.W., and Watson, D. 2002. Acute and Chronic Toxicity of Nickel to Marine Organisms: Implications for Water Quality Criteria. *Environmental Toxicology and Chemistry* 21(11):2423-2430.
116. Hunt, J.W., Anderson, B.S., Turpen, S.L., Englund, M.A., and Piekarski, W. 1997. Precision and Sensitivity of a Seven-Day Growth and Survival Toxicity Test using the West Coast Marine Mysid Crustacean *Holmesimysis costata*. *Environmental Toxicology and Chemistry* 16(4):824-834.
117. Hurlbert, S.H. 1984. Pseudoreplication and the Design of Ecological Field Experiments. *Ecological Monographs* 54(2):187-211.
118. Hutchinson, T.H. 2002. Reproductive and Developmental Effects of Endocrine Disrupters in Invertebrates: in Vitro and in Vivo Approaches. *Toxicology Letters* 131(1-2):75-81.
119. Hutchinson, T.H., Brown, R., Brugger, K.E., Campbell, P.M., Holt, M., Lange, R., Mccahon, P., Tattersfield, L.J., and Van Egmond, R. 2000. Ecological Risk Assessment of Endocrine Disruptors. *Environmental Health Perspectives* 108(11):1007-1014.
120. Hutchinson, T.H., Jha, A.N., and Dixon, D.R. 1995. The Polychaete *Platynereis dumerilii* (Audouin and Milne-Edwards) - a New Species for Assessing the Hazardous Potential of Chemicals in the Marine-Environment. *Ecotoxicology and Environmental Safety* 31(3):271-281.
121. Hutchinson, T.H., Pounds, N.A., Hampel, M., and Williams, T.D. 1999a. Impact of Natural and Synthetic Steroids on the Survival, Development and Reproduction of Marine Copepods (*Tisbe battagliai*). *The Science of the Total Environment* 233(1-3):167-179.
122. Hutchinson, T.H., Pounds, N.A., Hampel, M., and Williams, T.D. 1999b. Life-Cycle Studies With Marine Copepods (*Tisbe battagliai*) Exposed to 20-Hydroxyecdysone and Diethylstilbestrol. *Environmental Toxicology and Chemistry* 18(12):2914-2920.
123. Ingersoll, C.G., Hutchinson, T.H., Crane, M., Dodson, S., DeWitt, T., Gies, A., Huet, M.-C., McKenney, C.L., Oberdörster, E., Pascoe, D., Versteeg, D.J., and Warwick, O. 1999. The Endocrinology of Invertebrates. In: *Laboratory toxicity tests for evaluating potential effects of endocrine disrupting compounds*, DeFur, P.L., Crane, M., Ingersoll, C., and Tetterfield, L. eds., Pp. 107–197. Society of Environmental Toxicology and Chemistry (SETAC) Press. Pensacola, Florida.

124. James, M.O. and Boyle, S.M. 1998. Cytochromes P450 in Crustacea. *Comparative Biochemistry and Physiology C-Toxicology & Pharmacology* 121(1-3):157-172.
125. Jarman, S.N., Nicol, S., Elliott, N.G., and McMinn, A. 2000. 28s rDNA Evolution in the Eumalacostraca and the Phylogenetic Position of Krill. *Molecular Phylogenetics and Evolution* 17(1):26-36.
126. Jobling, S., Casey, D., Rodgers-Gray, T., Oehlmann, J., Schulte-Oehlmann, U., Pawlowski, S., Baunbeck, T., Turner, A.P., and Tyler, C.R. 2003. Comparative Responses of Molluscs and Fish to Environmental Estrogens and an Estrogenic Effluent. *Aquatic Toxicology* 65(2):205-220.
127. Johnston, N.M. and Ritz, D.A. 2001. Synchronous Development and Release of Broods by the Swarming Mysids *Anisomysis mixta Australis*, *Paramesopodopsis rufa* and *Tenagomysis tasmaniae* (Mysidacea: Crustacea). *Marine Ecology Progress Series* 223:225-233.
128. Jonckheere, A.R. 1954. A Distribution-Free K-Sample Test Against Ordered Alternatives. *Biometrika* 59:467-471.
129. Kahl, M.D., Makynen, E.A., Kosian, P.A., and Ankley, G.T. 1997. Toxicity of 4-Nonylphenol in a Life-Cycle Test With the Midge *Chironomus tentans*. *Ecotoxicology and Environmental Safety* 38(2):155-160.
130. Kashian, D.R. and Dodson, S.I. 2004. Effects of Vertebrate Hormones on Development and Sex Determination in *Daphnia magna*. *Environmental Toxicology and Chemistry* 23(5):1282-1288.
131. Khan, A., Barbieri, J., Khan, S., and Sweeney, F. 1992. A New Short-Term Mysid Toxicity Test using Sexual Maturity as an Endpoint. *Aquatic Toxicology* 23(2):97-105.
132. Kostrouch, Z., Kostrouchova, M., Love, W., Jannini, E., Piatigorsky, J., and Rall, J.E. 1998. Retinoic Acid X Receptor in the Diploblast, *Tripedalia cystophora*. *Proceedings of the National Academy of Sciences of the United States of America* 95(23):13442-13447.
133. Kreeger, K.E., Kreeger, D.A., Langdon, C.J., and Lowry, R.R. 1991. The Nutritional Value of *Artemia* and *Tigriopus californicus* (Baker) for Two Pacific Mysid Species, *Metamysidopsis elongata* (Holmes) and *Mysidopsis intii* (Holmquist). *Journal of Experimental Marine Biology and Ecology* 148(2):147-158.
134. Kropp, R.K. 1982. Responses of Five Holothurian Species to Attacks by a Predatory Gastropod, *Tonna pernix*. *Pacific Science* 36:445-452.
135. Kuhn, A., Munns, W.R., Champlin, D., McKinney, R., Tagliabue, M., Serbst, J., and Gleason, T. 2001. Evaluation of the Efficacy of Extrapolation Population Modeling to Predict the Dynamics of *Americamysis bahia* Populations in the Laboratory. *Environmental Toxicology and Chemistry* 20(1):213-221.
136. Kuhn, A., Munns, W.R., Poucher, S., Champlin, D., and Lussier, S. 2000. Prediction of Population-Level Response from Mysid Toxicity Test Data using Population Modeling Techniques. *Environmental Toxicology and Chemistry* 19(9):2364-2371.
137. Lafont, R. 2000a. The Endocrinology of Invertebrates. *Ecotoxicology* 9(1-2):41-57.

138. Lafont, R. 2000b. Understanding Insect Endocrine Systems: Molecular Approaches. *Entomologia Experimentalis et Applicata* 97(2):123-136.
139. Lagadic, L. and Caquet, T. 1998. Invertebrates in Testing of Environmental Chemicals: Are They Alternatives? *Environmental Health Perspectives* 106:593-611.
140. Langdon, C.J., Harmon, V.L., Vance, M.M., Kreeger, K.E., Kreeger, D.A., and Chapman, G.A. 1996. A 7-D Toxicity Test for Marine Pollutants using the Pacific Mysid *Mysidopsis intii*. 1. Culture and Protocol Development. *Environmental Toxicology and Chemistry* 15(10):1815-1823.
141. Laufer, H., Ahl, J., Rotllant, G., and Baclaski, B. 2002. Evidence that Ecdysteroids and Methyl Farnesoate Control Allometric Growth and Differentiation in a Crustacean. *Insect Biochemistry and Molecular Biology* 32(2):205-210.
142. Laufer, H. and Biggers, W.J. 2001. Unifying Concepts Learned from Methyl Farnesoate for Invertebrate Reproduction and Post-Embryonic Development. *American Zoologist* 41(3):442-457.
143. Laufer, H., Borst, D., Baker, F.C., Carrasco, C., Sinkus, M., Reuter, C.C., Tsai, L.W., and Schooley, D.A. 1987a. Identification of a Juvenile Hormone-Like Compound in a Crustacean. *Science* 235:202-205.
144. Laufer, H., Landau, M., Homola, E., and Borst, D.W. 1987b. Methyl Farnesoate: Its Site of Synthesis and Regulation of Secretion in a Juvenile Crustacean. *Insect Biochemistry* 17:1129-1131.
145. Lawrence, A. and Poulter, C. 1996. The Potential Role of the Estuarine Amphipod *Gammarus duebeni* in Sub-Lethal Ecotoxicology Testing. *Water Science and Technology* 34(7-8):93-100.
146. LeBlanc, G.A., Campbell, P.M., den Besten, P., Brown, R.P., Chang, E.S., Coats, J.R., deFur, P.L., Dhadialla, T., Edwards, J., Riddiford, L.M., Simpson, M.G., Snell, T.W., Thorndyke, M., and Matsumura, F. 1999. The Endocrinology of Invertebrates. In: *Endocrine Disruption in Invertebrates: Endocrinology, Testing, and Assessment*, DeFur P.L., Crane, M., Ingersoll, C., and Tetterfield, L. eds. Pp. 23-106. Society of Environmental Toxicology and Chemistry (SETAC) Press. Pensacola, Florida.
147. LeBlanc, G.A. and McLachlan, J.B. 1999. Molt-Independent Growth Inhibition of *Daphnia magna* by a Vertebrate Antiandrogen. *Environmental Toxicology and Chemistry* 18(7):1450-1455.
148. Lee, R.F. 1991. Metabolism of Tributyltin by Marine Animals and Possible Linkages to Effects. *Marine Environmental Research* 32:29-35.
149. Lee, R.F. and Oshima, Y. 1998. Effects of Selected Pesticides, Metals and Organometallics on Development of Blue Crab (*Callinectes sapidus*) Embryos. *Marine Environmental Research* 46:479-482.
150. Leitz, T. 2001. Endocrinology of the Cnidaria: State of the Art. *Zoology-Analysis of Complex Systems* 103(3-4):202-221.
151. Leung, B. and Forbes, M.R. 1996. Fluctuating Asymmetry in Relation to Stress and Fitness: Effects of Trait Type as Revealed by Meta-Analysis. *Ecoscience* 3(4):400-413.
152. Leung, B., Forbes, M.R., and Houle, D. 2000. Fluctuating Asymmetry as a Bioindicator of Stress: Comparing Efficacy of Analyses Involving Multiple Traits. *American Naturalist* 155(1):101-115.

153. Lussier, S.M., Champlin, D., Livolsi, J., Poucher, S., and Pruell, R.J. 2000. Acute Toxicity of Para-Nonylphenol to Saltwater Animals. *Environmental Toxicology and Chemistry* 19(3):617-621.
154. Lussier, S.M., Gentile, J.H., and Walker, J. 1985. Acute and Chronic Effects of Heavy Metals and Cyanide on *Mysidopsis bahia* (Crustacea:Mysidacea). *Aquatic Toxicology* 7:25-35.
155. Lussier, S.M., Kuhn, A., Chammas, M.J., and Sewall, J. 1988. Techniques for the Laboratory Culture of *Mysidopsis* Species (Crustacea Mysidacea). *Environmental Toxicology and Chemistry* 7(12):969-977.
156. Lussier, S.M., Kuhn, A., and Comeleo, R. 1999. An Evaluation of the Seven-Day Toxicity Test With *Americamysis bahia* (Formerly *Mysidopsis bahia*). *Environmental Toxicology and Chemistry* 18(12):2888-2893.
157. Marcial, H.S., Hagiwara, A., and Snell, T.W. 2003. Estrogenic Compounds Affect Development of Harpacticoid Copepod *Tigriopus japonicus*. *Environmental Toxicology and Chemistry* 22(12):3025-3030.
158. Martin, M., Hunt, J.W., Anderson, B.S., Turpen, S.L., and Palmer, F.H. 1989. Experimental evaluation of the mysid *Holmesimysis costata* as a test organism for effluent toxicity testing. *Environmental Toxicology and Chemistry* 8:1003-1012.
159. Matthiessen, P. and Gibbs, P.E. 1998. Critical Appraisal of the Evidence for Tributyltin-Mediated Endocrine Disruption in Mollusks. *Environmental Toxicology and Chemistry* 17(1):37-43.
160. Matthiessen, P., Sheahan, D., Harrison, R., Kirby, M.F., Rycroft, R., Turnbull, A., Volkner, C., and Williams, R. 1995. Use of a *Gammarus pulex* Bioassay to Measure the Effects of Transient Carbofuran Runoff from Farmland. *Ecotoxicology and Environmental Safety* 30:111-119.
161. Mauchline, J. 1980. The Biology of Mysids and Euphausiids. *Advances in Marine Biology* 18:1-369.
162. McCahon, C.P. and Pascoe, D. 1988. Culture Techniques for Three Fresh-Water Macroinvertebrate Species and Their Use in Toxicity Tests. *Chemosphere* 17(12):2471-2480.
163. McGee, B.L. and Spencer, M. 2001. A Field-Based Population Model for the Sediment Toxicity Test Organism *Leptocheirus plumulosus*: II. Model Application. *Marine Environmental Research* 51(4):347-363.
164. McKenney, C.L. Jr. 1982. Interrelationships Between Energy Metabolism, Growth Dynamics, and Reproduction During the Life Cycle of *Mysidopsis bahia* as Influenced by Sublethal Endrin Exposure. In: *Physiological Mechanisms of Marine Pollutant Toxicity*, Vernberg W.B., Calabrese, A., Thurberg, F.P., and Vernberg, F.J. eds. Pp. 447-476. Academic Press. New York.
165. McKenney, C.L. Jr. 1985. Associations Between Physiological Alterations and Population Changes in an Estuarine Mysid During Chronic Exposure to a Pesticide. In: *Marine Pollution and Physiology: Recent Advances*. Pp. 397-418. University of South Carolina Press. Columbia, S.C.
166. McKenney, C.L. Jr. 1986. Influence of the Organophosphate Insecticide Fenthion on *Mysidopsis bahia* Exposed During a Complete Life-Cycle .1. Survival, Reproduction, and Age-Specific Growth. *Diseases of Aquatic Organisms* 1(2):131-139.

167. McKenney, C.L., Jr. 1987. *Optimization of Environmental Factors during the Life Cycle of Mysidopsis bahia*. Environmental Research Brief, EPA/600/M-87/004, U.S. EPA, Cincinnati, OH, 6 p.
168. McKenney, C.L. Jr. 1989. The Relationship Between Modified Energy Metabolism and Inhibited Growth and Reproduction in an Estuarine Mysid during Chronic Pesticide Exposure. *American Zoologist* 29(4):A63.
169. McKenney, C.L. Jr. 1994. Alterations in Growth, Reproduction, and Energy Metabolism of Estuarine Crustaceans as Indicators of Pollutant Stress. In: *Biological Monitoring of the Environment: A Manual of Methods*. Salanki J., Jeffrey, D., and Hughes, G.M. eds., Pp. 111-115. CAB International. Wallingford, England.
170. McKenney, C.L. Jr. 1996. The Combined Effects of Salinity and Temperature on Various Aspects of the Reproductive Biology of the Estuarine Mysid, *Mysidopsis bahia*. *Invertebrate Reproduction & Development* 29(1):9-18.
171. McKenney, C.L. Jr. 1998. Physiological Dysfunction in Estuarine Mysids and Larval Decapods With Chronic Pesticide Exposure. In: *Microscale Testing in Aquatic Toxicology: Advances, Techniques, and Practice*, Wells, P.G., Lee, K., and Blaise, C. eds. Pp. 465-476. CRC Press. Boca Raton, FL.
172. McKenney, C.L. Jr. 1999. Hormonal Processes in Decapod Crustacean Larvae as Biomarkers of Endocrine Disrupting Chemicals in the Marine Environment. In: *Environmental Toxicology and Risk Assessment: Standardization of Biomarkers for Endocrine Disruption and Environmental Assessment*. Henshel D.S., Black, M.C., and Harrass, M.C. eds. Pp. 119-135. American Society for Testing and Materials. West Conshohocken, PA.
173. McKenney, C.L., Jr. and Celestial, D.M. 1995. Interactions among Salinity, Temperature, and Age on Growth of the Estuarine Mysid *Mysidopsis bahia* Reared in the Laboratory Through a Complete Life Cycle: I. Body Mass and Age-specific Growth Rate. *Journal of Crustacean Biology* 15:169-178.
174. McKenney, C.L. Jr. and Celestial, D.M. 1996. Modified Survival, Growth and Reproduction in an Estuarine Mysid (*Mysidopsis bahia*) Exposed to a Juvenile Hormone Analogue Through a Complete Life Cycle. *Aquatic Toxicology* 35(1):11-20.
175. McKenney, C.L. Jr., Hamaker, T.L., and Matthews, E. 1991. Changes in the Physiological Performance and Energy Metabolism of an Estuarine Mysid (*Mysidopsis bahia*) Exposed in the Laboratory Through a Complete Life-Cycle to the Defoliant DEF. *Aquatic Toxicology* 19(2):123-135.
176. McKenney, C.L. Jr. and Matthews, E. 1990. Alterations in the Energy Metabolism of an Estuarine Mysid (*Mysidopsis bahia*) as Indicators of Stress from Chronic Pesticide Exposure. *Marine Environmental Research* 30(1):1-19.
177. McKenney, C.L. Jr., Weber, D.E., Celestial, D.M., and MacGregor, M.A. 1998. Altered Growth and Metabolism of an Estuarine Shrimp (*Palaemonetes pugio*) During and after Metamorphosis Onto Fenvalerate-Laden Sediment. *Archives of Environmental Contamination and Toxicology* 35(3):464-471.

178. McPeck, M.A. and Wellborn, G.A. 1998. Genetic Variation and Reproductive Isolation among Phenotypically Divergent Amphipod Populations. *Limnology and Oceanography* 43(6):1162-1169.
179. McTavish, K., Stech, H., and Stay, F. 1998. A Modeling Framework for Exploring the Population-Level Effects of Endocrine Disruptors. *Environmental Toxicology and Chemistry* 17(1):58-67.
180. Mees, J., Abdulkarim, Z., and Hamerlynck, O. 1994. Life-History, Growth and Production of *Neomysis-integer* in the Westerschelde Estuary (SW Netherlands). *Marine Ecology Progress Series* 109(1):43-57.
181. Mees, J., Fockedey, N., Dewicke, A., Janssen, C.R., and Sorbe, J.-C. 1995. Aberrant Individuals of *Neomysis integer* and Other Mysidacea: Intersexuality and Variable Telson Morphology. *Netherlands Journal of Aquatic Ecology* 29(2):161-166.
182. Meregalli, G., Pluymers, L., and Ollevier, F. 2001. Induction of Mouthpart Deformities in *Chironomus riparius* Larvae Exposed to 4-n-Nonylphenol. *Environmental Pollution* 111(2):241-246.
183. Molenock, J. 1969. *Mysidopsis bahia*, a New Species of Mysid (Crustacea: Mysidacea) from Galveston Bay, Texas. *Tulane Studies in Zoology and Botany* 15(3):113-116.
184. Moore, C.G. and Stevenson, J.M. 1994. Intersexuality in Benthic Harpacticoid Copepods in the Firth of Forth, Scotland. *Journal of Natural History* 28:1213-1230.
185. Moore, C.G. and Stevenson, J.M. 1991. The Occurrence of Intersexuality on Harpacticoid Copepods and its Relationship with Pollution. *Marine Pollution Bulletin* 22:72-74.
186. Moreau, X., Benzid, D., De Jong, L., Barthelemy, R.M., and Casanova, J.P. 2002. Evidence for the Presence of Serotonin in Mysidacea (Crustacea, Peracarida) as Revealed by Fluorescence Immunohistochemistry. *Cell and Tissue Research* 310(3):359-371.
187. Mu, X.Y. and LeBlanc, G.A. 2002a. Developmental Toxicity of Testosterone in the Crustacean *Daphnia magna* Involves Anti-Ecdysteroidal Activity. *General and Comparative Endocrinology* 129(2):127-133.
188. Mu, X.Y. and LeBlanc, G.A. 2002b. Environmental Antiectdysteroids Alter Embryo Development in the Crustacean *Daphnia magna*. *Journal of Experimental Zoology* 292(3):287-292.
189. Mu, X.Y. and LeBlanc, G.A. 2004. Synergistic Interaction of Endocrine-Disrupting Chemicals: Model Development using an Ecdysone Receptor Antagonist and a Hormone Synthesis Inhibitor. *Environmental Toxicology and Chemistry* 23(4):1085-1091.
190. Nagaraju, G.P.C., Suraj, N.J., and Reddy, P.S. 2003. Methyl Farnesoate Stimulates Gonad Development in *Macrobrachium malcolmsonii* (H. Milne Edwards) (Decapoda, Palaemonidae). *Crustaceana* 76:1171-1178.
191. Nates, S.F. and McKenney, C.L. 2000. Growth, Lipid Class and Fatty Acid Composition in Juvenile Mud Crabs (*Rhithropanopeus harrisi*) Following Larval Exposure to Fenoxycarb (R), Insect Juvenile Hormone Analog. *Comparative Biochemistry and Physiology C-Toxicology & Pharmacology* 127(3):317-325.

192. Nice, H.E., Morritt, D., Crane, M., and Thorndyke, M. 2003. Long-Term and Transgenerational Effects of Nonylphenol Exposure at a Key Stage in the Development of *Crassostrea gigas*. Possible Endocrine Disruption? *Marine Ecology Progress Series* 256:293-300.
193. Nimmo, D.R., Bahner, L.H., Rigby, R.A., Sheppard, J.M., and Wilson, A.J. Jr. 1977. *Mysidopsis bahia*: An Estuarine Species Suitable for Life-Cycle Toxicity Tests to Determine the Effects of a Pollutant. In: *Aquatic Toxicology and Hazard Evaluation*, Mayer F.L. and Hamelink, J.L. eds. Pp. 109-116. American Society for Testing and Materials. Philadelphia, PA.
194. Nimmo, D.R. and Hamaker, T.L. 1982. Mysids in Toxicity Testing - A Review. *Hydrobiologia* 93(1-2):171-178.
195. Nimmo, D.R., Hamaker, T.L., Mathews, E., and Moore, J.C. 1981. An Overview of the Acute and Chronic Effects of 1st and 2nd Generation Pesticides on an Estuarine Mysid *Mysidopsis bahia*. In: *Biological Monitoring Of Marine Pollutants; Symposium On Pollution And Physiology Of Marine Organisms*. Pp. 3-20. Academic Press, Inc.. New York, N.Y.
196. Nimmo, D.R., Hamaker, T.L., and Sommers, C.A. 1978b. *Entire Life-Cycle Toxicity Test using Mysids (Mysidopsis bahia) in Flowing Water*. EPA-600/9-78-010. U.S. Environmental Protection Agency, Environmental Research Laboratory, Gulf Breeze, FL. (ERL,GB X107).
197. Nipper, M.G. and Williams, E.K. 1997. Culturing and Toxicity Testing with the New Zealand Mysid *Tenagomysis novae-zealandiae*, with a Summary of Toxicological Research in this Group. *Australasian Journal of Ecotoxicology* 3:117-129.
198. Oberdörster, E. and Cheek, A.O. 2001. Gender Benders at the Beach: Endocrine Disruption in Marine and Estuarine Organisms. *Environmental Toxicology and Chemistry* 20(1):23-36.
199. Oberdörster, E., Rice, C.D., and Irwin, L.K. 2000. Purification of Vitellin from Grass Shrimp *Palaemonetes pugio*, Generation of Monoclonal Antibodies, and Validation for the Detection of Lipovitellin in Crustacea. *Comparative Biochemistry and Physiology C-Pharmacology Toxicology & Endocrinology* 127(2):199-207.
200. OECD (Organization for Economic Cooperation and Development). 1998. *Daphnia magna Reproduction Test*. OECD Guidelines for Testing of Chemicals. Technical Guidance Document 211. 21 pp.
201. OECD (Organization for Economic Cooperation and Development). 2000. *Guidance Document on Aquatic Toxicity Testing of Difficult Substances and Mixtures*. OECD Series on Testing and Assessment. Number 23. 53 pp.
202. OECD (Organization for Economic Cooperation and Development). 2001a. *Proposal of a New Guideline 218: Sediment-water Chironomid Toxicity Test Using Spiked Sediment*. OECD Guidelines for Testing of Chemicals. Draft Document February 2001. 21 pp.
203. OECD (Organization for Economic Cooperation and Development). 2001b. *Proposal of a New Guideline 219: Sediment-water Chironomid Toxicity Test Using Spiked Water*. OECD Guidelines for Testing of Chemicals. Draft Document February 2001. 21 pp.
204. OECD (Organization for Economic Cooperation and Development). 2003. *Draft Guidance Document on the Statistical Analysis of Ecotoxicity Data*.

205. OECD (Organization for Economic Cooperation and Development). 2003b. *Proposal for an Enhanced Test Guideline. Daphnia magna Reproduction Test*. Draft OECD Guidelines for Testing of Chemicals. Enhanced Technical Guidance Document 211. 21 pp.
206. OECD (Organization for Economic Cooperation and Development). 2004a. *Draft Proposal for a New Guideline: Mysid Two-generation Test Guideline*. 17 pp.
207. OECD (Organization for Economic Cooperation and Development). 2004b. *Proposal for a New Guideline: Calanoid Copepod Development and Reproduction Test with Acartia tonsa*. OECD Draft Guidelines for Testing of Chemicals. 39 pp.
208. OECD (Organization for Economic Cooperation and Development). 2004c. *Proposal for a New Guideline: Harpacticoid Copepod Development and Reproduction Test*. OECD Draft Guidelines for Testing of Chemicals. 34 pp.
209. Oehlmann, J. and Schulte-Oehlmann, U. 2003. Endocrine Disruption in Invertebrates. *Pure and Applied Chemistry* 75(11-12):2207-2218.
210. Oehlmann, J., Schulte-Oehlmann, U., Tillmann, M., and Markert, B. 2000. Effects of Endocrine Disruptors on Prosobranch Snails (Mollusca : Gastropoda) in the Laboratory. Part I: Bisphenol A and Octylphenol as Xeno-Estrogens. *Ecotoxicology* 9(6):383-397.
211. Olmstead, A.W. and LeBlanc, G.A. 2000. Effects of Endocrine-Active Chemicals on the Development of Sex Characteristics of *Daphnia magna*. *Environmental Toxicology and Chemistry* 19(8):2107-2113.
212. Olmstead, A.W. and LeBlanc, G.A. 2003. Insecticidal Juvenile Hormone Analogs Stimulate the Production of Male Offspring in the Crustacean *Daphnia magna*. *Environmental Health Perspectives* 111(7):919-924.
213. Olmstead, A.W. and LeBlanc, G.A. 2002. Juvenoid Hormone Methyl Farnesoate is a Sex Determinant in the Crustacean *Daphnia magna*. *Journal of Experimental Zoology* 293(7):736-739.
214. Palmer, A.R. 1994. Fluctuating Asymmetry Analysis: A Primer. Markow, T.A. ed. Pp. 335-364. Kluwer. Dordrecht.
215. Palmer, A.R. and Strobeck, C. 1986. Fluctuating Asymmetry: Measurement, Analysis and Patterns. *Annual Review of Ecology and Systematics* 17:391-421.
216. Pascoe, D., Carroll, K., Karntanut, W., and Watts, M.M. 2002. Toxicity of 17 α -ethinylestradiol and Bisphenol A to the Freshwater Cnidarian *Hydra vulgaris*. *Archives of Environmental Contamination and Toxicology* 43(1):56-63.
217. Pascoe, D., Keywords, T.J., Maund, S.J., Math, E., and Taylor, E.J. 1994. Laboratory and Field Evaluation of a Behavioural Bioassay - the *Gammarus pulex* (L.) Precopula Separation Test (GAPPS) Test. *Water Research* 28:369-372.
218. Pennak, R.W. 1989. Fresh-Water Invertebrates of the United States. Third Edition. John Wiley & Sons. New York.
219. Pernet, V. and Anctil, M. 2002. Annual Variations and Sex-Related Differences of Estradiol-17 β Levels in the Anthozoan *Renilla koellikeri*. *General and Comparative Endocrinology* 129(1):63-68.

220. Poulton, M. and Pascoe, D. 1990. Disruption of Precopula in *Gammarus pulex* (L.) - Development of a Behavioral Bioassay for Evaluating Pollutant and Parasite Induced Stress. *Chemosphere* 20:403-415.
221. Pounds, N.A., Hutchinson, T.H., Williams, T.D., Whiting, P., and Dinan, L. 2002. Assessment of Putative Endocrine Disrupters in an in Vivo Crustacean Assay and an in Vitro Insect Assay. *Marine Environmental Research* 54(3-5):709-713.
222. Price, W.W., Heard, R.W., and Stuck, L. 1994. Observations on the Genus *Mysidopsis* Sars, 1864 With the Designation of a New Genus, *Americamysis*, and the Descriptions of *Americamysis alleni* and *A. stucki* (Peracarida: Mysidacea: Mysidae), from the Gulf of Mexico. *Proceedings of the Biological Society of Washington* 107(4):680-698.
223. Quackenbush, L.S. 1986. Crustacean Endocrinology, a Review. *Canadian Journal of Fisheries and Aquatic Sciences* 43:2271-2282.
224. Quirt, J. and Lasenby, D. 2002. Cannibalism and Ontogenetic Changes in the Response of the Freshwater Shrimp *Mysis relicta* to Chemical Cues from Conspecific Predators. *Canadian Journal of Zoology* 80(6):1022-1025.
225. Reitsema, L.A. and Neff, J.M. 1980. A Recirculating Artificial Seawater System for the Laboratory Culture of *Mysidopsis almyra* (Crustacea; Pericaridea). *Estuaries* 3(4):321-323.
226. Richter, S. and Scholtz, G. 2001. Phylogenetic Analysis of the Malacostraca (Crustacea). *Journal of Zoological Systematics and Evolutionary Research* 39(3):113-136.
227. Ristola, T., Parker, D., and Kukkonen, J.V.K. 2001. Life-Cycle Effects of Sediment-Associated 2,4,5-Trichlorophenol on Two Groups of the Midge *Chironomus riparius* With Different Exposure Histories. *Environmental Toxicology and Chemistry* 20(8):1772-1777.
228. Roast, S.D., Thompson, R.S., Widdows, J., and Jones, M.B. 1998. Mysids and Environmental Monitoring: a Case for Their Use in Estuaries. *Marine and Freshwater Research* 49(8):827-832.
229. Roast, S.D., Widdows, J., and Jones, M.B. 1999. Scope for Growth of the Estuarine Mysid *Neomysis integer* (Peracarida: Mysidacea): Effects of the Organophosphate Pesticide Chlorpyrifos. *Marine Ecology Progress Series* 191:233-241.
230. Roast, S.D., Widdows, J., and Jones, M.B. 2000b. Disruption of Swimming in the Hyperbenthic Mysid *Neomysis integer* (Peracarida : Mysidacea) by the Organophosphate Pesticide Chlorpyrifos. *Aquatic Toxicology* 47(3-4):227-241.
231. Roast, S.D., Widdows, J., and Jones, M.B. 2001. Impairment of Mysid (*Neomysis integer*) Swimming Ability: an Environmentally Realistic Assessment of the Impact of Cadmium Exposure. *Aquatic Toxicology* 52:217-227.
232. Roast, S.D., Widdows, J., and Jones, M.B. 2000a. Mysids and Trace Metals: Disruption of Swimming as a Behavioural Indicator of Environmental Contamination. *Marine Environmental Research* 50(1-5):107-112.
233. Rotchell, J.M. and Ostrander, G.K. 2003. Molecular Markers of Endocrine Disruption in Aquatic Organisms. *Journal of Toxicology and Environmental Health-Part B-Critical Reviews* 6(5):453-495.

234. Ruiz, G.M., Fofonoff, P., Hines, A.H., and Grosholz, E.D. 1999. Non-Indigenous Species as Stressors in Estuarine and Marine Communities: Assessing Invasion Impacts and Interactions. *Limnology and Oceanography* 44(3):950-972.
235. Ruiz, G.M., Fofonoff, P.W., Carlton, J.T., Wonham, M.J., and Hines, A.H. 2000. Invasion of Coastal Marine Communities in North America: Apparent Patterns, Processes, and Biases. *Annual Review of Ecology and Systematics* 31:481-531.
236. Salzet, M. 2001. The Neuroendocrine System of Annelids. *Canadian Journal of Zoology-Revue Canadienne De Zoologie* 79(2):175-191.
237. Schulte-Oehlmann, U., Tillmann, M., Markert, B., Oehlmann, J., Watermann, B., and Scherf, S. 2000. Effects of Endocrine Disruptors on Prosobranch Snails (Mollusca : Gastropoda) in the Laboratory. Part II: Triphenyltin as a Xeno-Androgen. *Ecotoxicology* 9(6):399-412.
238. Segner, H., Carroll, K., Fenske, M., Janssen, C.R., Maack, G., Pascoe, D., Schafers, C., Vandenberg, G.F., Watts, M., and Wenzel, A. 2003. Identification of Endocrine-Disrupting Effects in Aquatic Vertebrates and Invertebrates: Report from the European Idea Project. *Ecotoxicology and Environmental Safety* 54(3):302-314.
239. Singer, M.M., George, S., Lee, I., Jacobson, S., Weetman, L.L., Blondina, G., Tjeerdema, R.S., Aurand, D., and Sowby, M.L. 1998. Effects of Dispersant Treatment on the Acute Aquatic Toxicity of Petroleum Hydrocarbons. *Archives of Environmental Contamination and Toxicology* 34(2):177-187.
240. Snedecor, G.W. and Cochran, W.G. 1980. *Statistical Methods*. The Iowa State University Press. Ames, Iowa.
241. Snyder, M.J. 2000. Cytochrome P450 Enzymes in Aquatic Invertebrates: Recent Advances and Future Directions. *Aquatic Toxicology* 48(4):529-547.
242. Snyder, M.J. and Mulder, E.P. 2001. Environmental Endocrine Disruption in Decapod Crustacean Larvae: Hormone Titrers, Cytochrome P450, and Stress Protein Responses to Heptachlor Exposure. *Aquatic Toxicology* 55(3-4):177-190.
243. Sommer, C. 1996. Ecotoxicology and Developmental Stability as an in Situ Monitor of Adaptation. *Ambio* 25:374-376.
244. Spencer, M. and McGee, B.L. 2001. A Field-Based Population Model for the Sediment Toxicity Test Organism *Leptocheirus plumulosus*: I. Model Development. *Marine Environmental Research* 51(4):327-345.
245. Steimle, F.W., Pikanowski, R.A., McMillan, D.G., Zetlin, C.A., and Wilk, S.J. 2000. Demersal Fish and American Lobster Diets in the Lower Hudson - Raritan Estuary. *NOAA Technical Memorandum NMFS-NE-161*:106 p.
246. Stephan, C.E. and Rogers, J.W. 1985. Advantages of using regression to calculate results of chronic toxicity tests. *Aquatic Toxicology and Hazard Assessment: Eighth Symposium*. American Society for Testing and Materials. Philadelphia, PA.

247. Subramoniam, T. 2000. Crustacean Ecdysteroids in Reproduction and Embryogenesis. *Comparative Biochemistry and Physiology C-Pharmacology Toxicology & Endocrinology* 125:135-156.
248. Sánchez, P. and Tarazona, J.V. 2002. Development of a Multispecies System for Testing Reproductive Effects on Aquatic Invertebrates. Experience With *Daphnia magna*, *Chironomus prasinus* and *Lymnaea peregra*. *Aquatic Toxicology* 60(3-4):249-256.
249. Tarrant, A.M., Atkinson, M.J., and Atkinson, S. 2004. Effects of Steroidal Estrogens on Coral Growth and Reproduction. *Marine Ecology Progress Series* 269:121-129.
250. Tarrant, A.M., Atkinson, M.J., and Atkinson, S. 2001. Uptake of Estrone from the Water Column by a Coral Community. *Marine Biology* 139(2):321-325.
251. Tarrant, A.M., Atkinson, S., and Atkinson, M.J. 1999. Estrone and Estradiol-17 Beta Concentration in Tissue of the Scleractinian Coral, *Montipora verrucosa*. *Comparative Biochemistry and Physiology A-Molecular and Integrative Physiology* 122(1):85-92.
252. Tarrant, A.M., Blomquist, C.H., Lima, P.H., Atkinson, M.J., and Atkinson, S. 2003. Metabolism of Estrogens and Androgens by Scleractinian Corals. *Comparative Biochemistry and Physiology B-Biochemistry & Molecular Biology* 136(3):473-485.
253. Tatarazako, N., Takao, Y., Kishi, K., Onikura, N., Arizono, K., and Iguchi, T. 2002. Styrene Dimers and Trimers Affect Reproduction of Daphnid (*Ceriodaphnia dubia*). *Chemosphere* 48(6):597-601.
254. Tatarazako, N., Oda, S., Watanabe, H., Morita, M., and Iguchi, T. 2003. Juvenile Hormone Agonists Affect the Occurrence of Male *Daphnia*. *Chemosphere* 53(8):827-833.
255. Terpstra, T.J. 1952. The Asymptotic Normality and Consistency of Kendall's Test Against Trend When Ties Are Present in One Ranking. *Indagationes Mathematicae* 14:327-333.
256. Thiel, M. 1997. Reproductive Biology of a Filter-Feeding Amphipod, *Leptocheirus pinguis*, with Extended Parental Care. *Marine Biology* 130(2):249-258.
257. Tillmann, M., Schulte-Oehlmann, U., Duft, M., Markert, B., and Oehlmann, J. 2001. Effects of Endocrine Disruptors on Prosobranch Snails (Mollusca : Gastropoda) in the Laboratory. Part III: Cyproterone Acetate and Vinclozolin as Antiandrogens. *Ecotoxicology* 10(6):373-388.
258. Touart LW. 1982. The effects of diflubenzuron on molting and regeneration of two estuarine crustaceans, the mysid shrimp *Mysidopsis bahia* and the grass shrimp *Palaemonetes pugio*. University of West Florida, Pensacola, FL.
259. Turpen, S., Hunt, J.W., Anderson, B.S., and Pearse, J.S. 1994. Population Structure, Growth, and Fecundity of the Kelp Forest Mysid *Holmesimysis costata* in Monterey Bay, California. *Journal of Crustacean Biology* 14(4):657-664.
260. UCSC (University of California Santa Cruz). 1998. Acute and Chronic Nickel Toxicity: Development of an Acute-to-Chronic Ratio for West Coast Marine Species, Final Report Results of Toxicity Testing with Topsmelt, Abalone, and Mysids. San Jose/Santa Clara Water Pollution Control Plant.

261. Vandenberg, G.F., Adriaens, D., Verslycke, T., and Janssen, C.R. 2003. Effects of 17 α -ethinylestradiol on Sexual Development of the Amphipod *Hyalella azteca*. *Ecotoxicology and Environmental Safety* 54(2):216-222.
262. Verslycke, T., De Wasch, K., De Brabander, H.F., and Janssen, C.R. 2002. Testosterone Metabolism in the Estuarine Mysid *Neomysis Integer* (Crustacea; Mysidacea): Identification of Testosterone Metabolites and Endogenous Vertebrate-Type Steroids. *General and Comparative Endocrinology* 126(2):190-199.
263. Verslycke, T. and Janssen, C.R. 2002. Effects of a Changing Abiotic Environment on the Energy Metabolism in the Estuarine Mysid Shrimp *Neomysis integer* (Crustacea : Mysidacea). *Journal of Experimental Marine Biology and Ecology* 279(1-2):61-72.
264. Verslycke, T., Poelmans, S., De Wasch, K., De Brabander, H.F., and Janssen, C.R. 2004b. Testosterone and Energy Metabolism in the Estuarine Mysid *Neomysis integer* (Crustacea : Mysidacea) Following Exposure to Endocrine Disruptors. *Environmental Toxicology and Chemistry* 23(5):1289-1296.
265. Verslycke, T., Poelmans, S., De Wasch, K., Vercauteren, J., Devos, C., Moens, L., Sandra, P., De Brabander, H.F., and Janssen, C.R. 2003b. Testosterone Metabolism in the Estuarine Mysid *Neomysis integer* (Crustacea; Mysidacea) Following Tributyltin Exposure. *Environmental Toxicology and Chemistry* 22(9):2030-2036.
266. Verslycke, T., Vercauteren, J., Devos, C., Moens, L., Sandra, P., and Janssen, C.R. 2003a. Cellular Energy Allocation in the Estuarine Mysid Shrimp *Neomysis integer* (Crustacea: Mysidacea) Following Tributyltin Exposure. *Journal of Experimental Marine Biology and Ecology* 288(2):167-179.
267. Verslycke, T.A., Fockede, N., McKenney Jr., C.L., Roast, S.D., Jones, M.B., Mees, J., and Janssen, C.R. 2004a. Mysid Crustaceans as Potential Test Organisms for the Evaluation Of Environmental Endocrine Disruption: A Review. *Environmental Toxicology and Chemistry* 23(5):1219-1234.
268. Volz, D.C. and Chandler, G.T. 2004. An Enzyme-Linked Immunosorbent Assay for Lipovitellin Quantification in Copepods: a Screening Tool for Endocrine Toxicity. *Environmental Toxicology and Chemistry* 23(2):298-305.
269. Volz, D.C., Wirth, E.F., Fulton, M.H., Scott, G.I., Block, D.S., and Chandler, G.T. 2002. Endocrine-Mediated Effects of UV-A Irradiation on Grass Shrimp (*Palaemonetes pugio*) Reproduction. *Comparative Biochemistry and Physiology C-Toxicology & Pharmacology* 133(3):419-434.
270. Watts, M.M. and Pascoe, D. 2000. Comparison of *Chironomus riparius* Meigen and *Chironomus tentans* Fabricius (Diptera : Chironomidae) for Assessing the Toxicity of Sediments. *Environmental Toxicology and Chemistry* 19(7):1885-1892.
271. Watts, M.M. and Pascoe, D. 1996. Use of the Freshwater Macroinvertebrate *Chironomus riparius* (Diptera: Chironomidae) in the Assessment of Sediment Toxicity. *Water Science and Technology* 34(7-8):101-107.
272. Watts, M.M., Pascoe, D., and Carroll, K. 2001a. Chronic Exposure to 17 α -ethinylestradiol and Bisphenol A-Effects on Development and Reproduction in the Freshwater Invertebrate *Chironomus riparius* (Diptera : Chironomidae). *Aquatic Toxicology* 55(1-2):113-124.

273. Watts, M.M., Pascoe, D., and Carroll, K. 2003. Exposure to 17 α -ethinylestradiol and Bisphenol A- Effects on Larval Moulting and Mouthpart Structure of *Chironomus riparius*. *Ecotoxicology and Environmental Safety* 54(2):207-215.
274. Watts, M.W., Pascoe, D., and Carroll, K. 2002. Population Responses of the Freshwater Amphipod *Gammarus pulex* (L.) to an Environmental Oestrogen 17 α -ethinyloestradiol. *Environmental Toxicology and Chemistry* 21(2):445-450.
275. Watts, M.W., Pascoe, D., and Carroll, K. 2001b. Survival and Precopulatory Behaviour of *Gammarus pulex* Exposed to Two Xenoestrogens. *Water Research* 35(10):2347-2352.
276. Weber, C.I. (ed.) 1993. *Methods for Estimating the Acute Toxicity of Effluents and Receiving Waters to Freshwater and Marine Organisms*. EPA/600/4-90/027F. U.S. Environmental Protection Agency, Office of Prevention, Pesticides and Toxic Substances. Washington, DC.
277. Williams, A. B. 1984. Shrimps, Lobsters, and Crabs of the Atlantic Coast of the Eastern United States, Maine to Florida. Smithsonian Institution Press. Washington, DC,.
278. Williams, D.A. 1971. A Test for Differences Between Treatment Means When Several Dose Levels Are Compared with a Zero Dose Control. *Biometrics* 27:103-117.
279. Wilson, E.O. (ed.). 1988. *Biodiversity*. National Academy Press. Washington DC.
280. Winkler, G., Dodson, J.J., Bertrand, N., Thivierge, D., and Vincent, W.F. 2003. Trophic Coupling across the St. Lawrence River Estuarine Transition Zone. *Marine Ecology Progress Series* 251:59-73.
281. Winkler, G. and Greve, W. 2002. Laboratory Studies of the Effect of Temperature on Growth, Moulting and Reproduction in the Co-Occurring Mysids *Neomysis integer* and *Praunus flexuosus*. *Marine Ecology Progress Series* 235:177-188.
282. Witt, J.D.S. and Hebert, P.D.N. 2000. Cryptic Species Diversity and Evolution in the Amphipod Genus *Hyaella* Within Central Glaciated North America: a Molecular Phylogenetic Approach. *Canadian Journal of Fisheries and Aquatic Sciences* 57:687-698.
283. Wortham-Neal, J. and Price, W.W. 2002. Marsupial Development of *Americamysis bahia* (Crustacea: Mysida). *Journal of Crustacean Biology* 22(1):98-112.
284. Zou, E. and Fingerman, M. 1997. Synthetic Estrogenic Agents Do Not Interfere with Sex Differentiation but Do Inhibit Molting of the Cladoceran *Daphnia magna*. *Bulletin of Environmental Contamination and Toxicology* 58(4):596-602.
285. Zulkosky, A.M., Ferguson, P.L., and McElroy, A.E. 2002. Effects of Sewage-Impacted Sediment on Reproduction in the Benthic Crustacean *Leptocheirus plumulosus*. *Marine Environmental Research* 54(3-5):615-619.