

For Official Use

English - Or. English

**ENVIRONMENT DIRECTORATE
CHEMICALS AND BIOTECHNOLOGY COMMITTEE**

AOP 360: Chitin synthase 1 inhibition leading to mortality

Short name: inhibition leading to mortality

The draft AOP 360 on Inhibition of Chitin synthase 1 inhibition leading to mortality was reviewed by the scientific journal Environmental Toxicology and Chemistry (ET&C) in line with the principles described in the OECD Guidance Document for the scientific review of AOPs [\[Link\]](#). The review panel conducted the assessment of both an AOP journal article format developed for publication in ET&C and the Wiki content of the AOP. The corresponding AOP journal article has been accepted by ET&C and is available on the SETAC website [\[Link\]](#). The reviewers comments and responses from the authors are available in the discussion pages of AOP 360 in the Wiki [\[Link\]](#). The EAGMST now recommends the submission of this AOP to WNT and WPHA for endorsement.

Nathalie Delrue, Administrator
Nathalie.delrue@oecd.org

JT03480228

Foreword

This Adverse Outcome Pathway (AOP) 360 on I Inhibition of Chitin synthase 1 inhibition leading to mortality Response, has been developed under the auspices of the OECD AOP Development Programme, overseen by the Extended Advisory Group on Molecular Screening and Toxicogenomics (EAGMST), which is an advisory group under the Working Party of the National Coordinators for the Test Guidelines Programme (WNT).

The AOP has been reviewed for compliance with the AOP development principles following the EAGMST coaching approach. The scientific review was subsequently conducted by the scientific journal Environmental Toxicology and Chemistry (ET&C), following the OECD AOP review principles outlined in the Guidance Document on the scientific review of AOPs. This AOP was endorsed by the WNT and the Working Party on Hazard Assessment (WPHA) on XXX.

Through endorsement of this AOP, the WNT and the WPHA express confidence in the scientific review process that the AOP has undergone and accept the recommendation of the EAGMST that the AOP be disseminated publicly. Endorsement does not necessarily indicate that the AOP is now considered a tool for direct regulatory application.

The OECD's Chemicals and Biotechnology Committee agreed to declassification of this AOP on XXX.

This document is being published under the responsibility of the OECD's Chemicals and Biotechnology Committee.

The outcome of the compliance check and of the scientific review are publicly available in the [AOP Wiki](#) at the following links: [[internal review](#)].

Authors:

Simon Schmid ^{1,2}, You Song ¹, and Knut Erik Tollefsen ^{1,2,3}

¹ Norwegian Institute for Water Research (NIVA), Section of Ecotoxicology and Risk Assessment, Økernveien 94, N-0579, Oslo, Norway

² Faculty of Environmental Science and Resource Management (MINA), Norwegian University of Life Sciences (NMBU), N-1432, Ås, Norway

³ Centre for Environmental Radioactivity (CERAD), Norwegian University of Life Sciences (NMBU), N-1432 Ås, Norway

Contact: Simon.Schmid@niva.no

Acknowledgements: This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 859891 and was supported by NIVA's Computational Toxicology Program, NCTP (www.niva.no/nctp).

Abstract

In order to grow and develop, arthropods need to shed their exoskeleton (or cuticle) periodically and replace it with a new one in a process called molting. Successful molting, and therefore a successful development necessitates stability and integrity of the cuticle to support muscular contractions involved in the shedding of the old cuticle. The integrity of the cuticle is largely dependent on the N-acetylglucosamine (GlcNAc) polymer chitin. Therefore, arthropods heavily rely on chitin synthesis as chitin is one of the main constituents of the cuticle. The cuticular chitin synthase (CHS-1) is the key enzyme in the biosynthetic pathway and arthropods are therefore especially dependent on its proper function. The present AOP describes the effects of chemical inhibition of the cuticular chitin synthase (CHS-1) on the molting process leading to increased mortality in arthropods. Inhibition of CHS-1 is the molecular initiating event and leads to a decreased chitin content in the arthropod cuticle which leaves the organism immature at the stage for ecdysis. This phenomenon can be described as premature molting. The organism eventually dies due to being stuck in the old cuticle or due to the consequences of a weak exoskeleton after ecdysis. The AOP is considered to be very consistent. Essentiality of key events was rated as high for every key event and the biological plausibility was rated as high for the whole AOP. However, there does not exist very much empirical evidence that allows to draw a representative conclusion on dose concordance along the AOP whereas time concordance can be supported by knockdown studies of CHS-1. Therefore, empirical evidence was considered to be moderate and the quantitative understanding was considered to be low. The overall confidence in the AOP was valued as moderate. The present AOP will guide assay development for further experimental studies by revealing data and knowledge gaps. One of its primary applications will also be providing guidance in screening strategies in order to identify chemicals directly interacting with CHS-1.

Table of contents


Foreword	2
Abstract	4
Background	7
Graphical Representation	8
Summary of the AOP	9
Events	9
Key Event Relationships.....	9
Stressors	9
Overall Assessment of the AOP	10
Domain of Applicability	10
Essentiality of the Key Events	11
Weight of Evidence Summary	12
Quantitative Consideration	13
Considerations for Potential Applications of the AOP	14
Appendix 1 - MIE, KEs and AO	19
List of MIEs in this AOP	19
Event: 1522: Inhibition, Chitin synthase 1	19
Evidence for Perturbation by Stressor.....	20
Domain of Applicability.....	20
Key Event Description	21
How it is Measured or Detected.....	21
List of Key Events in the AOP	23
Event: 1523: Decrease, Cuticular chitin content.....	23
Domain of Applicability.....	23
Key Event Description	24
How it is Measured or Detected.....	24
Event: 1524: Increase, Premature molting.....	28
Domain of Applicability.....	28
Key Event Description	29
How it is Measured or Detected.....	29
List of Adverse Outcomes in this AOP.....	31
Event: 350: Increase, Mortality.....	31
Domain of Applicability.....	31
Key Event Description	32
How it is Measured or Detected.....	32
Regulatory Significance of the AO	32
Appendix 2 - List of Key Event Relationships in the AOP	34
List of Adjacent Key Event Relationships.....	34
Relationship: 1742: Inhibition, CHS-1 leads to Decrease, Cuticular chitin content.....	34
Evidence Supporting Applicability of this Relationship	34
Key Event Relationship Description	35
Evidence Supporting this KER	35

Quantitative Understanding of the Linkage	35
Relationship: 1743: Decrease, Cuticular chitin content leads to Increase, Premature molting	39
Evidence Supporting Applicability of this Relationship	39
Key Event Relationship Description	40
Evidence Supporting this KER	40
Quantitative Understanding of the Linkage	40
Relationship: 1744: Increase, Premature molting leads to Increase, Mortality	43
Evidence Supporting Applicability of this Relationship	43
Key Event Relationship Description	44
Evidence Supporting this KER	44
Quantitative Understanding of the Linkage	44

Background

Arthropods (including insects, crustaceans and arachnids) need to shed their exoskeleton in order to grow and reproduce. This process, also called molting or ecdysis, is mediated by behavioural mechanisms which involve the skeletal muscles (Ayali 2009; Song et al. 2017a). In order to properly shed its cuticle, the organism needs to possess a newly synthesized cuticle that possesses a certain integrity to support this process. Since chitin is a major constituent of the cuticle, it contributes substantially to its integrity (Cohen 2001; Vincent and Wegst 2004). Chitin is synthesized from uridine diphosphate-N-Acetylglucosamine (UDP-GlcNAc) in a polymerization reaction by the transmembrane enzyme chitin synthase isoform 1 (CHS-1). CHS-1 is localized on the apical side in the cuticular epithelium. Since chitin and the process of chitin synthesis does not occur in vertebrates, it can and has been exploited for the design of pest controlling agents. Inhibitors of chitin synthesis may not only be of use for the control of unwanted arthropods and fungi, they may also pose a risk for beneficial arthropods such as insects and crustaceans. Disruption of chitin synthesis or the endocrine mechanisms controlling molting generally lead to a disruption of ecdysis (Merzendorfer et al. 2012; Song et al. 2017a; Song et al. 2017b). If the amount of chitin in the cuticle decreases, the affected organism may not be able to molt properly and will most probably die of starvation or suffocation (Camp et al. 2014; Song et al. 2017a). Alternatively, if molting is completed despite an immature cuticle, the organism may be deformed and die as a consequence of a weak cuticle. Therefore, the present AOP should build the basis of a mechanistic approach for the systematic evaluation and the risk assessment of chemicals interfering with chitin synthesis by directly inhibiting CHS-1.

Graphical Representation



Summary of the AOP

Events

Molecular Initiating Events (MIE), Key Events (KE), Adverse Outcomes (AO)

Sequence	Type	Event ID	Title	Short name
1	MIE	1522	Inhibition, Chitin synthase 1	Inhibition, CHS-1
2	KE	1523	Decrease, Cuticular chitin content	Decrease, Cuticular chitin content
3	KE	1524	Increase, Premature molting	Increase, Premature molting
4	AO	350	Increase, Mortality	Increase, Mortality

Key Event Relationships

Upstream Event	Relationship Type	Downstream Event	Evidence	Quantitative Understanding
Inhibition, Chitin synthase 1	adjacent	Decrease, Cuticular chitin content	Moderate	Low
Decrease, Cuticular chitin content	adjacent	Increase, Premature molting	Moderate	Low
Increase, Premature molting	adjacent	Increase, Mortality	Moderate	Low

Stressors

Name	Evidence
Polyoxin B	High
Polyoxin D	High
Nikkomycins	High
Captan	Moderate
Captafol	Moderate
Folpet	Moderate

Overall Assessment of the AOP

Domain of Applicability

Life Stage

Life Stage	Evidence
larvae	High
Juvenile	High
Adult	Moderate

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
Pieris brassicae	Pieris brassicae	High	NCBI
Anopheles gambiae	Anopheles gambiae	High	NCBI
Lucilia cuprina	Lucilia cuprina	High	NCBI
Tribolium castaneum	Tribolium castaneum	High	NCBI
Bombyx mori	Bombyx mori	High	NCBI
Anopheles quadrimaculatus	Anopheles quadrimaculatus	High	NCBI
Trichoplusia ni	Trichoplusia ni	High	NCBI
Artemia salina	Artemia salina	High	NCBI
Daphnia magna	Daphnia magna	High	NCBI
Hyalophora cecropia	Hyalophora cecropia	High	NCBI
Ostrinia nubilalis	Ostrinia nubilalis	High	NCBI
Bradysia hygida	Bradysia hygida	Moderate	NCBI
Mamestra brassicae	Mamestra brassicae	Moderate	NCBI
Chilo suppressalis	Chilo suppressalis	Moderate	NCBI
Locusta migratoria	Locusta migratoria	Moderate	NCBI
Nilaparvata lugens	Nilaparvata lugens	Moderate	NCBI
Aphis glycines	Aphis glycines	Moderate	NCBI
Lepeophtheirus salmonis	Lepeophtheirus salmonis	Moderate	NCBI
Panonychus citri	Panonychus citri	Moderate	NCBI
Grapholita molesta	Grapholita molesta	Moderate	NCBI
Ectropis obliqua	Ectropis obliqua	Moderate	NCBI
Tigriopus japonicus	Tigriopus japonicus	Moderate	NCBI

Sex Applicability

Sex	Evidence
Unspecific	Moderate

Taxonomic: Since the whole phylum of arthropods is dependent on the synthesis of chitin to molt successfully, it is extremely likely that the AOP is applicable to all arthropods. Effect data along the AOP exist from Dipteran, Lepidopteran and Coleopteran insect species as well as from Branchiopods and Anostracans of the crustacea. Although data is limited, KEs seem to be well conserved across taxa, as shown in available studies with specific stressors known to inhibit CHS and in studies where CHS-1 was knocked down by RNA interference. However, due to limited data availability, it was not possible to cover

whole taxa but rather single species in the assessment of KEs. Alignment of amino acid residues in the catalytic center of CHS-1 using the Sequence Alignment to Predict Across Species Susceptibility tool (SeqAPASS, <https://seqapass.epa.gov/seqapass>, LaLone et al. 2016), confirmed structural and functional conservation in various insect, arachnid and crustacean species, strengthening the evidence for the applicability domain to be the whole phylum of arthropods. However, taxonomic applicability may not only be defined by structural conservation of the protein sequence. So the evidence for the taxonomic applicability for species with support only from sequence alignment was judged as moderate, whereas evidence for species with support from sequence alignment and effect data was judged as high.

Life stage: The AOP is applicable for organisms undergoing continuous molt cycles. As insects do not molt in their adulthood, the AOP is only applicable for larval and pupal stages of insects. Crustaceans and arachnids grow and molt throughout their lifetime (Passano 1961; Uhl et al. 2015), which makes the AOP applicable to all life stages, where juvenile life stages might be more susceptible to chemical perturbations due to higher growth rate and therefore more frequent molting.

Sex: The AOP is applicable to all sexes.

Chemical: Substances known to trigger the MIE and leading to the AO are of the family of pyrimidine nucleosides (e.g. polyoxin D, polyoxin B and nikkomycin Z) (Osada 2019). There also exists evidence for phthalimides (captan, captafol and folpet) to inhibit CHS-1 activity and to decrease the cuticular chitin content *in vitro* (Cohen and Casida 1982; Gelman and Borkovec 1986). However, as these substances are known to covalently bind to thiol groups in proteins (Lukens and Sisler 1958), it is not clear if the inhibition is due to specific CHS-1 inhibition or due to unspecific protein binding.

Essentiality of the Key Events

The essentiality of all key events was considered as high. Essentiality evaluations were mainly based on specifically designed studies demonstrating the expected effect pattern predicted by the AOP to occur after knockdown of CHS-1.

Inhibition, Chitin synthase 1 (High): Knockdown of the cuticular chitin synthase leads to the expected pattern of effects described in this AOP. It decreases the cuticular chitin content and leads to premature molting associated mortality in insects (Arakane et al. 2005; X. Zhang et al. 2010; Li et al. 2017; Zhai et al. 2017). If the cuticular chitin content was not directly measured as endpoint, knockdown of the CHS-1 led directly to the occurrence of premature molting associated increase of mortality (Chen et al. 2008; X. Zhang et al. 2010; Wang et al. 2012; Yang et al. 2013; Shang et al. 2016; Mohammed et al. 2017; Wang et al. 2019; Ye et al. 2019; Ullah et al. 2020)

Decrease, Cuticular chitin content (High): Abolishment of the cuticular chitin synthesis through knockdown of CHS-1 leads to premature molting associated mortality (Arakane et al. 2005; X. Zhang et al. 2010; Li et al. 2017; Zhai et al. 2017). By knocking down the UDP-GlcNAc pyrophosphorylase (UAP), which catalyzes the last sugar conversion before the polymerization to chitin, it was shown that reduced chitin content leads to the same outcome as the knockdown of CHS-1. Namely premature molting and increased mortality (Arakane et al. 2011; Liu et al. 2013). Knockdown of trehalase genes, which constitutes the start of the chitin synthetic pathway and convert trehalose to glucose, leads to a similar pattern of effects, namely decreased cuticular chitin content and premature molting associated mortality (Chen et al. 2010; Shi et al. 2016).

Increase, Premature molting (High): Several studies show that premature molting is a direct consequence of decreased chitin synthesis and leads to increased mortality. The KE is consistently listed as cause for mortality when CHS-1 is knocked down throughout a number of studies (Arakane et al. 2005;

Chen et al. 2008; J. Zhang et al. 2010; X. Zhang et al. 2010; Wang et al. 2012; Yang et al. 2013; Shang et al. 2016; Li et al. 2017; Mohammed et al. 2017; Zhai et al. 2017; Wang et al. 2019; Ye et al. 2019).

Increase, Mortality (High): Increased mortality was observed in all of the abovementioned studies.

Weight of Evidence Summary

Biological Plausibility: The biosynthesis of chitin is well characterized and is conserved among arthropods. Although the exact mode of action of chitin synthases remains elusive, it is widely accepted and well established that the chitin synthase is the key enzyme in the pathway, polymerizing chitin using UDP-*N*-Acetylglucosamine as substrate (Merzendorfer and Zimoch 2003).

Arthropod cuticles mostly consist of chitin embedded into a matrix of cuticular proteins. It is therefore widely accepted that chitin contributes crucially to the quality and function of the cuticle (Reynolds 1987; Muthukrishnan et al. 2012). The molting process requires the new cuticle to be strong enough to withstand the stresses of ecdysis.

During ecdysis, arthropods pause food intake and growth. If ecdysis is initiated before the new cuticle is strong enough, the organism likely dies of starvation or growth arrest (Song, Villeneuve, et al. 2017). It was also reported that certain arthropods pause respiration during ecdysis, which may lead to suffocation (Camp et al. 2014).

Based on the well-established biological knowledge on the processes this AOP bases on, the biological plausibility for all KER was rated as high.

Empirical Evidence: Empirical evidence assessment was conducted on the basis of *in vitro* and *in vivo* experiments performed with stressors affecting key events throughout the AOP. Studies showed that the key events are affected by model stressors such as Polyoxin D and Nikkomycin Z, which are able to competitively inhibit CHS1 (Endo et al. 1970). Several studies provide evidence that polyoxin B, polyoxin D and nikkomycin Z trigger the MIE in cell free systems of coleopteran, lepidopteran and dipteran insect species (Cohen 1982; Turnbull and Howells 1982; Kuwano and Cohen 1984; Cohen and Casida 1990; Zhang and Yan Zhu 2013). Also the cuticular chitin content was shown to be decreased by polyoxin D and nikkomycin Z in lepidopteran and dipteran species as well as in the crustacean *Artemia salina* (Gijswijt et al. 1979; Calcott and Fatig 1984; Gelman and Borkovec 1986; Zhuo et al. 2014). The AO is supported by *in vivo* studies with polyoxin D and nikkomycin Z in dipteran insects and *Daphnia magna* (Tellam et al. 2000; Tellam and Eisemann 2000; Zhu et al. 2007; Zhang and Yan Zhu 2013; New Zealand Environmental Protection Authority 2015). A major data gap constitutes the absence of data covering the KE “Increase, premature molting”. This KE is mentioned in some studies but never assessed as an individual endpoint (Gijswijt et al. 1979; Tellam et al. 2000). Another major data gap is the lacking quantitative data for KERs. As endpoints were only measured as individual endpoints and not in sequence, it makes it nearly impossible to evaluate the dose for the KEs and KERs. However, data from studies where CHS-1 was knocked down are able to support temporal concordance for all KERs. Knockdown of CHS-1 led to decreased chitin content and subsequently to premature molting associated mortality (Arakane et al., 2005; Li et al., 2017). Based on the major data gaps and therefore the lacking information on dose concordance as well as the given time concordance, empirical evidence was evaluated to be moderate for the whole AOP.

Overall confidence in the AOP: Both, essentiality of KEs and the biological plausibility of the whole AOP were considered to be high. However, due to missing quantitative data and the lack of evidence for

dose concordance, empirical evidence was judged to be moderate. Therefore the overall confidence in the AOP was evaluated as moderate.

Quantitative Consideration

Quantitative data are limited for all KER and therefore the whole AOP. Therefore, predictions on the occurrence of downstream KE and the AO on the basis of the occurrence of upstream KEs is not readily feasible. Quantitative understanding of the AOP was therefore considered to be low.

Considerations for Potential Applications of the AOP

Arthropods are responsible for many functions in terrestrial as well as aquatic ecosystems and are therefore jointly responsible for ecosystem health (Seastedt and Crossley 1984; Losey and Vaughan 2006; LeBlanc 2007). Therefore, it is important to develop AOPs which enhance the mechanistic knowledge on chemicals, such as chitin synthesis inhibitors, which may pose a risk to non-target arthropods. Those AOPs will contribute to the systematic use of mechanistic data to preserve beneficial arthropod populations and ecosystem health.

The present AOP will help to guide future experimental studies by identifying data gaps. This will lead to the identification and development suitable bioassays in order to populate the AOP with (quantitative) experimental data which may allow for predictions of regulatory relevant endpoints on the basis of the occurrence of the MIE.

The present AOP may also guide screening strategies in order to identify chemicals inhibiting CHS-1. The identified substances may then be prioritized and undergo a thorough hazard assessment.

As there already exist approaches to assess mixture toxicity using the AOP framework (Altenburger et al. 2012; Beyer et al. 2014), the present AOP could be employed for the effect assessment of mixtures of chemicals that share the same KEs (e.g. AOP #361, aopwiki.org/aops/361, AOP #358, aopwiki.org/aops/358, and AOP #359, aopwiki.org/aops/359).

References

- Altenburger R, Scholz S, Schmitt-Jansen M, Busch W, Escher BI. 2012. Mixture toxicity revisited from a toxicogenomic perspective. *Environ Sci Technol*. 46(5):2508–2522. doi:10.1021/es2038036.
- Arakane Y, Baguinin MC, Jasarapura S, Chaudhari S, Doyungan A, Kramer KJ, Muthukrishnan S, Beeman RW. 2011. Both UDP N-acetylglucosamine pyrophosphorylases of *Tribolium castaneum* are critical for molting, survival and fecundity. *Insect Biochem Mol Biol*. 41(1):42–50. doi:10.1016/j.ibmb.2010.09.011. <http://dx.doi.org/10.1016/j.ibmb.2010.09.011>.
- Arakane Y, Muthukrishnan S, Kramer KJ, Specht CA, Tomoyasu Y, Lorenzen MD, Kanost M, Beeman RW. 2005. The *Tribolium* chitin synthase genes TcCHS1 and TcCHS2 are specialized for synthesis of epidermal cuticle and midgut peritrophic matrix. *Insect Mol Biol*. 14(5):453–463. doi:10.1111/j.1365-2583.2005.00576.x.
- Ayali A. 2009. The role of the arthropod stomatogastric nervous system in moulting behaviour and ecdysis. *J Exp Biol*. 212(4):453–459. doi:10.1242/jeb.023879.
- Beyer J, Petersen K, Song Y, Ruus A, Grung M, Bakke T, Tollefsen KE. 2014. Environmental risk assessment of combined effects in aquatic ecotoxicology: A discussion paper. *Mar Environ Res*. 96:81–91. doi:10.1016/j.marenvres.2013.10.008. <http://dx.doi.org/10.1016/j.marenvres.2013.10.008>.
- Calcott PH, Fatig RO. 1984. Inhibition of Chitin metabolism by Avermectin in susceptible Organisms. *J Antibiot (Tokyo)*. 37(3):253–259. doi:10.7164/antibiotics.37.253.
- Camp AA, Funk DH, Buchwalter DB. 2014. A stressful shortness of breath: Molting disrupts breathing in the mayfly *Cloeon dipterum*. *Freshw Sci*. 33(3):695–699. doi:10.1086/677899.
- Chen Jie, Tang B, Chen H, Yao Q, Huang X, Chen Jing, Zhang D, Zhang W. 2010. Different functions of the insect soluble and membrane-bound trehalase genes in chitin biosynthesis revealed by RNA interference. *PLoS One*. 5(4). doi:10.1371/journal.pone.0010133.
- Chen X, Tian H, Zou L, Tang B, Hu J, Zhang W. 2008. Disruption of *Spodoptera exigua* larval development by silencing chitin synthase gene A with RNA interference. *Bull Entomol Res*. 98(6):613–619. doi:10.1017/S0007485308005932.
- Cohen E. 1982. In vitro chitin synthesis in an insect: formation and structure of microfibrils. *Eur J Cell Biol*. 26(2):289–294.
- Cohen E. 2001. Chitin synthesis and inhibition: A revisit. *Pest Manag Sci*. 57(10):946–950. doi:10.1002/ps.363.
- Cohen E, Casida JE. 1982. Properties and inhibition of insect integumental chitin synthetase. *Pestic Biochem Physiol*. 17(3):301–306. doi:10.1016/0048-3575(82)90141-9.
- Cohen E, Casida JE. 1990. Insect and Fungal Chitin Synthetase Activity: Specificity of Lectins as Enhancers and Nucleoside Peptides as Inhibitors. *Pestic Biochem Physiol*. 37(3):249–253. doi:10.1016/0048-3575(90)90131-K.
- Endo A, Kakiki K, Misato T. 1970. Mechanism of action of the antifungal agent polyoxin D. *J Bacteriol*. 104(1):189–196. doi:10.1128/jb.104.1.189-196.1970.
- Gelman DB, Borkovec AB. 1986. The pharate adult clasper as a tool for measuring chitin synthesis and for identifying new chitin synthesis inhibitors. *Comp Biochem Physiol Part C, Comp*. 85(1):193–197. doi:10.1016/0742-8413(86)90073-3.
- Gijswijt MJ, Deul DH, de Jong BJ. 1979. Inhibition of chitin synthesis by benzoyl-phenylurea insecticides, III. Similarity in action in *Pieris brassicae* (L.) with Polyoxin D. *Pestic Biochem Physiol*. 12(1):87–94. doi:10.1016/0048-3575(79)90098-1.

- Kuwano E, Cohen E. 1984. The use of a *Tribolium* chitin synthetase assay in studying the effects of benzimidazoles with a terpene moiety and related compounds. *Agric Biol Chem.* 48(6):1617–1620. doi:10.1080/00021369.1984.10866362.
- LaLone, C.A., Villeneuve, D.L., Lyons, D., Helgen, H.W., Robinson, S.L., Swintek, J.A., Saari, T.W., Ankley, G.T., 2016. Sequence alignment to predict across species susceptibility (seqapass): A web-based tool for addressing the challenges of cross-species extrapolation of chemical toxicity. *Toxicol. Sci.* 153, 228–245. <https://doi.org/10.1093/toxsci/kfw119>
- LeBlanc GA. 2007. Crustacean endocrine toxicology: A review. *Ecotoxicology.* 16(1):61–81. doi:10.1007/s10646-006-0115-z.
- Li T, Chen J, Fan X, Chen W, Zhang W. 2017. MicroRNA and dsRNA targeting chitin synthase A reveal a great potential for pest management of the hemipteran insect *Nilaparvata lugens*. *Pest Manag Sci.* 73(7):1529–1537. doi:10.1002/ps.4492.
- Liu X, Li F, Li D, Ma E, Zhang W, Zhu KY, Zhang J. 2013. Molecular and functional analysis of UDP-N-acetylglucosamine Pyrophosphorylases from the Migratory Locust, *Locusta migratoria*. *PLoS One.* 8(8). doi:10.1371/journal.pone.0071970.
- Losey JE, Vaughan M. 2006. The economic value of ecological services provided by insects. *Bioscience.* 56(4):311–323. doi:10.1641/0006-3568(2006)56[311:TEVOES]2.0.CO;2.
- Lukens RJ, Sisler HD. 1958. 2-Thiazolidinethione-4-carboxylic acid from the reaction of captan with cysteine. *Science (80-).* 127(3299):650. doi:10.1126/science.127.3299.650.
- Merzendorfer H, Kim HS, Chaudhari SS, Kumari M, Specht CA, Butcher S, Brown SJ, Robert Manak J, Beeman RW, Kramer KJ, et al. 2012. Genomic and proteomic studies on the effects of the insect growth regulator diflubenzuron in the model beetle species *Tribolium castaneum*. *Insect Biochem Mol Biol.* 42(4):264–276. doi:10.1016/j.ibmb.2011.12.008. <http://dx.doi.org/10.1016/j.ibmb.2011.12.008>.
- Merzendorfer H, Zimoch L. 2003. Chitin metabolism in insects: structure, function and regulation of chitin synthases and chitinases. *J Exp Biol.* 206(24):4393 LP – 4412. doi:10.1242/jeb.00709. <http://jeb.biologists.org/content/206/24/4393.abstract>.
- Mohammed AMA, Diab MR, Abdelsattar M, Khalil SMS. 2017. Characterization and RNAi-mediated knockdown of Chitin Synthase A in the potato tuber moth, *Phthorimaea operculella*. *Sci Rep.* 7(1):1–12. doi:10.1038/s41598-017-09858-y. <http://dx.doi.org/10.1038/s41598-017-09858-y>.
- Muthukrishnan S, Merzendorfer H, Arakane Y, Kramer KJ. 2012. Chitin Metabolism in Insects. Elsevier B.V. <http://dx.doi.org/10.1016/B978-0-12-384747-8.10007-8>.
- New Zealand Environmental Protection Authority. 2015. Application for approval to import ESTEEM for release. <https://www.epa.govt.nz/assets/FileAPI/hsno-ar/APP202334/fbce9a39e6/APP202334-APP202334-Staff-Report-Final-updated.pdf>.
- Osada H. 2019. Discovery and applications of nucleoside antibiotics beyond polyoxin. *J Antibiot (Tokyo).* 72(12):855–864. doi:10.1038/s41429-019-0237-1. <http://dx.doi.org/10.1038/s41429-019-0237-1>.
- Passano LM. 1961. The regulation of crustacean metamorphosis. *Integr Comp Biol.* 1(1):89–95. doi:10.1093/icb/1.1.89.
- Reynolds SE. 1987. The cuticle, growth and moulting in insects: The essential background to the action of acylurea insecticides. *Pestic Sci.* 20(2):131–146. doi:10.1002/ps.2780200207.
- Seastedt TR, Crossley DA. 1984. Influence of on arthropods ecosystems. *Bioscience.* 34(3):157–161.
- Shang F, Xiong Y, Xia WK, Wei DD, Wei D, Wang JJ. 2016. Identification, characterization and functional analysis of a chitin synthase gene in the brown citrus aphid, *Toxoptera citricida* (Hemiptera, Aphididae). *Insect Mol Biol.* 25(4):422–430. doi:10.1111/imb.12228.

- Shi JF, Xu QY, Sun QK, Meng QW, Mu LL, Guo WC, Li GQ. 2016. Physiological roles of trehalose in *Leptinotarsa* larvae revealed by RNA interference of trehalose-6-phosphate synthase and trehalase genes. *Insect Biochem Mol Biol.* 77:52–68. doi:10.1016/j.ibmb.2016.07.012.
- Song Y, Evenseth LM, Iguchi T, Tollefsen KE. 2017b. Release of chitinase as an indicator of potential molting disruption in juvenile *Daphnia magna* exposed to the ecdysone receptor agonist 20-hydroxyecdysone. *J Toxicol Environ Heal - Part A Curr Issues.* 80(16–18):954–962. doi:10.1080/15287394.2017.1352215. <https://doi.org/10.1080/15287394.2017.1352215>.
- Song Y, Villeneuve DL, Toyota K, Iguchi T, Tollefsen KE. 2017a. Ecdysone Receptor Agonism Leading to Lethal Molting Disruption in Arthropods: Review and Adverse Outcome Pathway Development. *Environ Sci Technol.* 51(8):4142–4157. doi:10.1021/acs.est.7b00480.
- Tellam RL, Eisemann C. 2000. Chitin is only a minor component of the peritrophic matrix from larvae of *Lucilia cuprina*. *Insect Biochem Mol Biol.* 30(12):1189–1201. doi:10.1016/S0965-1748(00)00097-7.
- Tellam RL, Vuocolo T, Johnson SE, Jarmey J, Pearson RD. 2000. Insect chitin synthase. cDNA sequence, gene organization and expression. *Eur J Biochem.* 267(19):6025–6043. doi:10.1046/j.1432-1327.2000.01679.x.
- Turnbull IF, Howells AJ. 1982. Effects of several larvicidal compounds on chitin biosynthesis by isolated larval integuments of the sheep blowfly *Lucilia cuprina*. *Aust J Biol Sci.* 35(5):491–504. doi:10.1071/BI9820491.
- Uhl G, Zimmer SM, Renner D, Schneider JM. 2015. Exploiting a moment of weakness: Male spiders escape sexual cannibalism by copulating with moulting females. *Sci Rep.* 5(July):1–7. doi:10.1038/srep16928.
- Vincent JFV, Wegst UGK. 2004. Design and mechanical properties of insect cuticle. *Arthropod Struct Dev.* 33(3):187–199. doi:10.1016/j.asd.2004.05.006.
- Wang Y, Fan HW, Huang HJ, Xue J, Wu WJ, Bao YY, Xu HJ, Zhu ZR, Cheng JA, Zhang CX. 2012. Chitin synthase 1 gene and its two alternative splicing variants from two sap-sucking insects, *Nilaparvata lugens* and *Laodelphax striatellus* (Hemiptera: Delphacidae). *Insect Biochem Mol Biol.* 42(9):637–646. doi:10.1016/j.ibmb.2012.04.009. <http://dx.doi.org/10.1016/j.ibmb.2012.04.009>.
- Wang Z, Yang H, Zhou C, Yang WJ, Jin DC, Long GY. 2019. Molecular cloning, expression, and functional analysis of the chitin synthase 1 gene and its two alternative splicing variants in the white-backed planthopper, *Sogatella furcifera* (Hemiptera: Delphacidae). *Sci Rep.* 9(1):1–14. doi:10.1038/s41598-018-37488-5. <http://dx.doi.org/10.1038/s41598-018-37488-5>.
- Yang WJ, Xu KK, Cong L, Wang JJ. 2013. Identification, mRNA expression, and functional analysis of chitin synthase 1 gene and its two alternative splicing variants in oriental fruit fly, *Bactrocera dorsalis*. *Int J Biol Sci.* 9(4):331–342. doi:10.7150/ijbs.6022.
- Ye C, Jiang Y Di, An X, Yang L, Shang F, Niu J, Wang JJ. 2019. Effects of RNAi-based silencing of chitin synthase gene on moulting and fecundity in pea aphids (*Acyrtosiphon pisum*). *Sci Rep.* 9(1):1–10. doi:10.1038/s41598-019-39837-4. <http://dx.doi.org/10.1038/s41598-019-39837-4>.
- Zhai Y, Fan X, Yin Z, Yue X, Men X, Zheng L, Zhang W. 2017. Identification and Functional Analysis of Chitin Synthase A in Oriental Armyworm, *Mythimna separata*. *Proteomics.* 17(21):1–11. doi:10.1002/pmic.201700165.
- Zhang J, Liu X, Zhang Jianqin, Li D, Sun Y, Guo Y, Ma E, Zhu KY. 2010. Silencing of two alternative splicing-derived mRNA variants of chitin synthase 1 gene by RNAi is lethal to the oriental migratory locust, *Locusta migratoria manilensis* (Meyen). *Insect Biochem Mol Biol.* 40(11):824–833. doi:10.1016/j.ibmb.2010.08.001. <http://dx.doi.org/10.1016/j.ibmb.2010.08.001>.

Zhang X, Yan Zhu K. 2013. Biochemical characterization of chitin synthase activity and inhibition in the African malaria mosquito, *Anopheles gambiae*. *Insect Sci.* 20(2):158–166. doi:10.1111/j.1744-7917.2012.01568.x.

Zhang X, Zhang J, Zhu KY. 2010. Chitosan/double-stranded RNA nanoparticle-mediated RNA interference to silence chitin synthase genes through larval feeding in the African malaria mosquito (*Anopheles gambiae*). *Insect Mol Biol.* 19(5):683–693. doi:10.1111/j.1365-2583.2010.01029.x.

Zhu KY, Heise S, Zhang J, Anderson TD, Starkey SR. 2007. Comparative Studies on Effects of Three Chitin Synthesis Inhibitors on Common Malaria Mosquito (Diptera: Culicidae). *J Med Entomol.* 44(6):1047–1053. doi:10.1093/jmedent/44.6.1047.

Zhuo W, Fang Y, Kong L, Li X, Sima Y, Xu S. 2014. Chitin synthase A: A novel epidermal development regulation gene in the larvae of *Bombyx mori*. *Mol Biol Rep.* 41(7):4177–4186. doi:10.1007/s11033-014-3288-1.

Appendix 1 - MIE, KEs and AO

List of MIEs in this AOP

Event: 1522: Inhibition, Chitin synthase 1

Short Name: Inhibition, CHS-1

Key Event Component

Process	Object	Action
chitin synthase activity		decreased

AOPs Including This Key Event

AOP ID and Name	Event Type
Aop:342 - S-adenosylmethionine depletion leading to population decline (1)	Molecular Initiating Event
Aop:360 - Chitin synthase 1 inhibition leading to mortality	Molecular Initiating Event

Stressors

Name
Polyoxin B
Polyoxin D
Nikkomycins
Captan

Biological Context

Level of Biological Organization
Molecular

Cell term

Cell term
cuticle secreting cell

Organ term

Organ term
epithelium

Evidence for Perturbation by Stressor

Overview for Molecular Initiating Event

Stressors known to competitively inhibit CHS1 are polyoxin B, polyoxin D and Nikkomycin Z (Cohen and Casida 1982; Cohen and Casida 1990; Zhang and Yan Zhu 2013). There may also be stressors that inhibit CHS-1 in a non-competitive manner which may become apparent in further characterization efforts of this MIE. There is also a study that reports inhibition of CHS-1 by the phthalimide fungicide captan (Cohen and Casida 1982). However, it remains elusive if the observed inhibition is due to specific interaction with the enzyme or due to unspecific protein binding which is the predominant mode of action of phthalimides (Lukens and Sisler 1958).

Domain of Applicability

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
Anopheles gambiae	Anopheles gambiae	High	NCBI
Tribolium castaneum	Tribolium castaneum	High	NCBI
Trichoplusia ni	Trichoplusia ni	High	NCBI
Hyalophora cecropia	Hyalophora cecropia	High	NCBI
Bradysia hygida	Bradysia hygida	Moderate	NCBI
Mamestra brassicae	Mamestra brassicae	Moderate	NCBI
Chilo suppressalis	Chilo suppressalis	Moderate	NCBI
Locusta migratoria	Locusta migratoria	Moderate	NCBI
Nilaparvata lugens	Nilaparvata lugens	Moderate	NCBI
Aphis glycines	Aphis glycines	Moderate	NCBI
Lepeophtheirus salmonis	Lepeophtheirus salmonis	Moderate	NCBI
Panonychus citri	Panonychus citri	Moderate	NCBI
Grapholita molesta	Grapholita molesta	Moderate	NCBI
Ectropis obliqua	Ectropis obliqua	Moderate	NCBI
Tigriopus japonicus	Tigriopus japonicus	Moderate	NCBI

Life Stage Applicability

Life Stage	Evidence
larvae	High
Juvenile	High
Adult	Moderate

Sex Applicability

Sex	Evidence
Unspecific	Moderate

Taxonomic: Effect data for the occurrence of CHS1 inhibition exist from Dipteran, Lepidopteran and Coleopteran insect species. Sequence alignment of CHS1 protein sequences using the Sequence Alignment to Predict Across Species Susceptibility (SeqAPASS, <https://seqapass.epa.gov/seqapass>) tool, yielded

susceptibility predictions for various insect species, arachnids and crustacean taxa such as branchiopods, hexanauplia, malacostraca and merostomata. However, most of the protein sequences were not identified as CHS1. The alignment of amino acid residues believed to be critical for ligand binding were therefore carried out with sequences identified as CHS1. Evidence was rated as high for species with a susceptibility prediction and effect data. Evidence was rated as moderate when only alignment data were available. Although most of the sequences are not annotated as CHS1, all arthropods rely on the synthesis of cuticular chitin therefore it is extremely likely that this MIE is applicable to the whole phylum of arthropods.

Life stage: This MIE is applicable for organisms undergoing continuous molt cycles. Namely larval stages of insects and all life stages of crustaceans and arachnids.

Sex: The MIE is applicable to all sexes.

Chemical: Substances known to trigger inhibit CHS-1 are of the family of pyrimidine nucleosides (e.g. polyoxin D, polyoxin B and nikkomycin Z) (Cohen and Casida 1982; Kuwano and Cohen 1984; Cohen and Casida 1990; Zhang and Yan Zhu 2013; Osada 2019). There also exists evidence for the phthalimide captan to inhibit CHS-1 activity *in vitro* (Cohen and Casida 1982). However, as phthalimides are known to covalently bind to thiol groups in proteins (Lukens and Sisler 1958), it is not clear if the inhibition is due to specific CHS-1 inhibition or due to unspecific protein binding.

Key Event Description

Chitin synthases are essential enzymes for all organisms synthesizing chitin, for example arthropods and fungi (Latgé 2007; Merzendorfer 2011). Chitin synthases polymerize chitin and subsequently translocate chitin through the cell membrane (Merzendorfer 2006; Merzendorfer 2011). In arthropods, two isoforms of the chitin synthase are known, CHS1, which is responsible for the synthesis of cuticular chitin, and chitin synthase isoform 2, which synthesizes chitin in the midgut (Arakane et al. 2005). In this MIE, inhibition of CHS-1 is characterized. The biological state being measured is the activity of the enzyme. CHS-1 has an essential role in the cuticle biology, as it constitutes the last and most critical step in the chitin biosynthetic pathway by catalyzing the polymerization of UDP-GlcNAc to chitin (Merzendorfer and Zimoch 2003; Merzendorfer 2006).

How it is Measured or Detected

Since the purification or even recombinant production of CHS-1 has not been achieved yet, the most common way is to use crude enzyme preparations for CHS-1 activity assays. It is noteworthy that in crude enzyme preparations of whole organisms both CHS isoforms, CHS-1 and CHS-2, are present. However, the expression of CHS-1 was shown to be much higher than CHS-2 in *Anopheles gambiae* (Zhang et al. 2012), therefore the effect of CHS-2 may be regarded as negligible. Alternatively, the digestive tract of the respective organism could be removed before producing the enzyme preparation. Different ways exist to detect the activity of the enzyme. One can incubate the enzyme preparation with radioactively labelled chitin precursors (e.g. ¹⁴C-UDP-GlcNAc) and measure radioactivity in the formed chitin chains by scintillation counting (Cohen 1982; Cohen and Casida 1990). Chitin synthase activity can also be measured in a non-radioactive way after the addition of precursors to a crude enzyme extract. There, the detection of CHS-1 activity involves the binding of chitin chains to wheat germ agglutinin (WGA) which possesses specific chitin binding properties (Lucero et al. 2002; Zhang and Yan Zhu 2013). The assay builds on the principle of a sandwich-ELISA, where chitin binds to a layer of WGA. A second layer of WGA which is conjugated to horseradish peroxidase (HRP) is then added and subsequently incubated with a HRP substrate. The cleavage of the HRP substrate leads to color formation and the amount of chitin synthesized can be determined colorimetrically.

References

- Arakane Y, Muthukrishnan S, Kramer KJ, Specht CA, Tomoyasu Y, Lorenzen MD, Kanost M, Beeman RW. 2005. The *Tribolium* chitin synthase genes TcCHS1 and TcCHS2 are specialized for synthesis of epidermal cuticle and midgut peritrophic matrix. *Insect Mol Biol.* 14(5):453–463. doi:10.1111/j.1365-2583.2005.00576.x.
- Cohen E. 1982. In vitro chitin synthesis in an insect: formation and structure of microfibrils. *Eur J Cell Biol.* 26(2):289–294.
- Cohen E, Casida JE. 1982. Properties and inhibition of insect integumental chitin synthetase. *Pestic Biochem Physiol.* 17(3):301–306. doi:10.1016/0048-3575(82)90141-9.
- Cohen E, Casida JE. 1990. Insect and Fungal Chitin Synthetase Activity: Specificity of Lectins as Enhancers and Nucleoside Peptides as Inhibitors. *Pestic Biochem Physiol.* 37(3):249–253. doi:10.1016/0048-3575(90)90131-K.
- Kuwano E, Cohen E. 1984. The use of a *Tribolium* chitin synthetase assay in studying the effects of benzimidazoles with a terpene moiety and related compounds. *Agric Biol Chem.* 48(6):1617–1620. doi:10.1080/00021369.1984.10866362.
- Latgé JP. 2007. The cell wall: A carbohydrate armour for the fungal cell. *Mol Microbiol.* 66(2):279–290. doi:10.1111/j.1365-2958.2007.05872.x.
- Lucero HA, Kuranda MJ, Bulik DA. 2002. A nonradioactive, high throughput assay for chitin synthase activity. *Anal Biochem.* 305(1):97–105. doi:10.1006/abio.2002.5594.
- Lukens RJ, Sisler HD. 1958. 2-Thiazolidinethione-4-carboxylic acid from the reaction of captan with cysteine. *Science (80-).* 127(3299):650. doi:10.1126/science.127.3299.650.
- Merzendorfer H. 2006. Insect chitin synthases: A review. *J Comp Physiol B Biochem Syst Environ Physiol.* doi:10.1007/s00360-005-0005-3.
- Merzendorfer H. 2011. The cellular basis of chitin synthesis in fungi and insects: Common principles and differences. *Eur J Cell Biol.* 90(9):759–769. doi:10.1016/j.ejcb.2011.04.014. <http://dx.doi.org/10.1016/j.ejcb.2011.04.014>.
- Merzendorfer H, Zimoch L. 2003. Chitin metabolism in insects: structure, function and regulation of chitin synthases and chitinases. *J Exp Biol.* 206(24):4393 LP – 4412. doi:10.1242/jeb.00709. <http://jeb.biologists.org/content/206/24/4393.abstract>.
- Osada H. 2019. Discovery and applications of nucleoside antibiotics beyond polyoxin. *J Antibiot (Tokyo).* 72(12):855–864. doi:10.1038/s41429-019-0237-1. <http://dx.doi.org/10.1038/s41429-019-0237-1>.
- Zhang X, Yan Zhu K. 2013. Biochemical characterization of chitin synthase activity and inhibition in the African malaria mosquito, *Anopheles gambiae*. *Insect Sci.* 20(2):158–166. doi:10.1111/j.1744-7917.2012.01568.x.
- Zhang X, Zhang J, Park Y, Zhu KY. 2012. Identification and characterization of two chitin synthase genes in African malaria mosquito, *Anopheles gambiae*. *Insect Biochem Mol Biol.* 42(9):674–682. doi:10.1016/j.ibmb.2012.05.005. <http://dx.doi.org/10.1016/j.ibmb.2012.05.005>.

List of Key Events in the AOP

Event: 1523: Decrease, Cuticular chitin content

Short Name: Decrease, Cuticular chitin content

Key Event Component

Process	Object	Action
cuticle development	cuticle	decreased

AOPs Including This Key Event

AOP ID and Name	Event Type
Aop:343 - S-adenosylmethionine depletion leading to population decline (2)	Key Event
Aop:342 - S-adenosylmethionine depletion leading to population decline (1)	Key Event
Aop:360 - Chitin synthase 1 inhibition leading to mortality	Key Event
Aop:361 - Sulfonylureareceptor binding leading to mortality	Key Event

Stressors

Name
Polyoxin D
Nikkomycins
Captan
Captafol
Folpet

Biological Context

Level of Biological Organization
Tissue

Organ term

Organ term
cuticle

Domain of Applicability

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
Pieris brassicae	Pieris brassicae	High	NCBI
Lucilia cuprina	Lucilia cuprina	High	NCBI
Bombyx mori	Bombyx mori	High	NCBI
Artemia salina	Artemia salina	High	NCBI

Term	Scientific Term	Evidence	Links
Ostrinia nubilalis	Ostrinia nubilalis	High	NCBI

Life Stage Applicability

Life Stage	Evidence
larvae	High
Juvenile	High
Adult	Moderate

Sex Applicability

Sex	Evidence
Unspecific	Moderate

Taxonomic: Effect data for the occurrence of this KE exist from *Pieris brassicae*, *Lucilia cuprina*, *Bombyx mori*, *Artemia salina* and *Ostrinia nubilalis*, defining its taxonomic applicability. Most likely, this KE is applicable to the whole phylum of arthropods, as they all rely on chitin as part of their exoskeleton.

Life stage: This KE is applicable for organisms synthesizing chitin in order to grow and develop, namely larval stages of insects and all life stages of crustaceans and arachnids.

Sex: This KE is applicable to all sexes.

Chemical: Substances known decrease the cuticular chitin content are of the family of pyrimidine nucleosides (e.g. polyoxin D and nikkomycin Z) (Gijswijt et al. 1979; Turnbull and Howells 1982; Calcott and Fatig 1984; Zhuo et al. 2014; Osada 2019). There also exists evidence for phthalimides (captan, captafol and folpet) to decrease the cuticular chitin content *in vitro* (Gelman and Borkovec 1986). However, as these substances are known to covalently bind to thiol groups in proteins (Lukens and Sisler 1958), it is not clear if the inhibition is due to specific CHS-1 inhibition or due to unspecific protein binding.

Key Event Description

This key event describes the decrease in cuticular chitin content. Chitin is a major part of the arthropod cuticle and therefore also responsible for its integrity (Reynolds 1987; Muthukrishnan et al. 2012). The cuticle is the exoskeleton of arthropods and has manifold functions, it protects organisms from predators, loss of water, acts as a physical barrier against microbial pathogens and provides support for muscular function (Vincent and Wegst 2004). Hence, cuticular chitin is also indispensable for the development of arthropods, as an immaculate cuticle is required for proper molting and therefore also for the growth of an organism.

How it is Measured or Detected

Several ways to determine cuticular chitin are described in the literature. Some of them are based on the determination of amino sugars after digestion or hydrolysis of chitin. For example, after the digestion of chitin by a bacterial chitinase, the *N*-Acetylglucosamine (GlcNAc) amount can be determined colorimetrically by a modified Morgan-Elson assay (Reissig et al. 1955; Arakane et al. 2005).

Alternatively, one can also quantify glucosamine colorimetrically after deacetylation and hydrolysis of chitin (Lehmann and White 1975; Zhang and Zhu 2006).

There also exists an approach based on the detection of fluorescence after staining with calcofluor white. In this assay, no treatment of the samples is necessary, the detection is carried out in homogenates of the respective organisms as calcofluor white directly binds to chitin (Henriques et al. 2020).

Chitin can also be quantified using radioactively labelled precursors (e.g. ^{14}C -UDP-GlcNAc) which are incorporated into *in vitro* cultured integument pieces or into the cuticle of whole organisms (Gijswijt et al. 1979; Turnbull and Howells 1982; Calcott and Fatig 1984; Gelman and Borkovec 1986).

Another possibility is to use the non-radioactive assay developed to measure chitin synthase activity (Lucero et al. 2002; Zhang and Yan Zhu 2013). Instead of adding an enzyme extract and chitin precursors to the reaction, one could simply add homogenized chitin containing material to the reaction to quantify its chitin content.

References

- Arakane Y, Muthukrishnan S, Kramer KJ, Specht CA, Tomoyasu Y, Lorenzen MD, Kanost M, Beeman RW. 2005. The *Tribolium* chitin synthase genes TcCHS1 and TcCHS2 are specialized for synthesis of epidermal cuticle and midgut peritrophic matrix. *Insect Mol Biol*. 14(5):453–463. doi:10.1111/j.1365-2583.2005.00576.x.
- Calcott PH, Fatig RO. 1984. Inhibition of Chitin metabolism by Avermectin in susceptible Organisms. *J Antibiot (Tokyo)*. 37(3):253–259. doi:10.7164/antibiotics.37.253.
- Clarke KU. 1957. On the Increase in Linear Size During Growth in *Locusta Migratoria* L. *Proc R Entomol Soc London Ser A, Gen Entomol*. 32(1–3):35–39. doi:10.1111/j.1365-3032.1957.tb00361.x.
- Dall W, Smith DM, Press B. 1978. Water uptake at ecdysis in the western rock lobster. *J Exp Mar Bio Ecol*. 35(1960). doi:10.1016/0022-0981(78)90074-6.
- deFur PL, Mangum CP, McMahon BR. 1985. Cardiovascular and Ventilatory Changes During Ecdysis in the Blue Crab *Callinectes Sapidus* Rathbun. *J Crustac Biol*. 5(2):207–215. doi:10.2307/1547867.
- Ewer J. 2005. How the ecdysozoan changed its coat. *PLoS Biol*. 3(10):1696–1699. doi:10.1371/journal.pbio.0030349.
- Gelman DB, Borkovec AB. 1986. The pharate adult clasper as a tool for measuring chitin synthesis and for identifying new chitin synthesis inhibitors. *Comp Biochem Physiol Part C, Comp*. 85(1):193–197. doi:10.1016/0742-8413(86)90073-3.
- Gijswijt MJ, Deul DH, de Jong BJ. 1979. Inhibition of chitin synthesis by benzoyl-phenylurea insecticides, III. Similarity in action in *Pieris brassicae* (L.) with Polyoxin D. *Pestic Biochem Physiol*. 12(1):87–94. doi:10.1016/0048-3575(79)90098-1.
- Henriques BS, Garcia ES, Azambuja P, Genta FA. 2020. Determination of Chitin Content in Insects: An Alternate Method Based on Calcofluor Staining. *Front Physiol*. 11(February):1–10. doi:10.3389/fphys.2020.00117.
- Lee RM. 1961. The variation of blood volume with age in the desert locust (*Schistocerca gregaria* Forsk.). *J Insect Physiol*. 6(1):36–51. doi:10.1016/0022-1910(61)90090-7.
- Lehmann PF, White LO. 1975. Chitin Assay Used to Demonstrate Renal Localization and Cortisone-Enhanced Growth of *Aspergillus fumigatus* Mycelium in Mice. *Infect Immun*. 12(5):987–992.
- Lucero HA, Kuranda MJ, Bulik DA. 2002. A nonradioactive, high throughput assay for chitin synthase activity. *Anal Biochem*. 305(1):97–105. doi:10.1006/abio.2002.5594.
- Lukens RJ, Sisler HD. 1958. 2-Thiazolidinethione-4-carboxylic acid from the reaction of captan with cysteine. *Science (80-)*. 127(3299):650. doi:10.1126/science.127.3299.650.
- Muthukrishnan S, Merzendorfer H, Arakane Y, Kramer KJ. 2012. Chitin Metabolism in Insects. Elsevier B.V. <http://dx.doi.org/10.1016/B978-0-12-384747-8.10007-8>.
- Osada H. 2019. Discovery and applications of nucleoside antibiotics beyond polyoxin. *J Antibiot (Tokyo)*. 72(12):855–864. doi:10.1038/s41429-019-0237-1. <http://dx.doi.org/10.1038/s41429-019-0237-1>.
- Reissig JL, Strominger JL, Leloir LF. 1955. A modified colorimetric method for the estimation of N-acetylamino sugars. *J Biol Chem*.:959–966.
- Reynolds SE. 1987. The cuticle, growth and moulting in insects: The essential background to the action of acylurea insecticides. *Pestic Sci*. 20(2):131–146. doi:10.1002/ps.2780200207.
- Turnbull IF, Howells AJ. 1982. Effects of several larvicidal compounds on chitin biosynthesis by isolated larval integuments of the sheep blowfly *Lucilia cuprina*. *Aust J Biol Sci*. 35(5):491–504. doi:10.1071/BI9820491.

Vincent JFV, Wegst UGK. 2004. Design and mechanical properties of insect cuticle. *Arthropod Struct Dev.* 33(3):187–199. doi:10.1016/j.asd.2004.05.006.

Zhang J, Zhu KY. 2006. Characterization of a chitin synthase cDNA and its increased mRNA level associated with decreased chitin synthesis in *Anopheles quadrimaculatus* exposed to diflubenzuron. *Insect Biochem Mol Biol.* 36(9):712–725. doi:10.1016/j.ibmb.2006.06.002.

Zhang X, Yan Zhu K. 2013. Biochemical characterization of chitin synthase activity and inhibition in the African malaria mosquito, *Anopheles gambiae*. *Insect Sci.* 20(2):158–166. doi:10.1111/j.1744-7917.2012.01568.x.

Zhuo W, Fang Y, Kong L, Li X, Sima Y, Xu S. 2014. Chitin synthase A: A novel epidermal development regulation gene in the larvae of *Bombyx mori*. *Mol Biol Rep.* 41(7):4177–4186. doi:10.1007/s11033-014-3288-1.

Event: 1524: Increase, Premature molting**Short Name: Increase, Premature molting****Key Event Component**

Process	Object	Action
ecdysis, chitin-based cuticle		decreased

AOPs Including This Key Event

AOP ID and Name	Event Type
Aop:343 - S-adenosylmethionine depletion leading to population decline (2)	Key Event
Aop:342 - S-adenosylmethionine depletion leading to population decline (1)	Key Event
Aop:358 - Chitinase inhibition leading to mortality	Key Event
Aop:359 - Chitinase inhibition leading to mortality	Key Event
Aop:360 - Chitin synthase 1 inhibition leading to mortality	Key Event
Aop:361 - Sulfonylureareceptor binding leading to mortality	Key Event

Stressors

Name
Polyoxin D
Nikkomycins

Biological Context

Level of Biological Organization
Individual

Domain of Applicability**Taxonomic Applicability**

Term	Scientific Term	Evidence	Links
Pieris brassicae	Pieris brassicae	High	NCBI
Lucilia cuprina	Lucilia cuprina	High	NCBI

Life Stage Applicability

Life Stage	Evidence
larvae	High
Juvenile	High
Adult	Moderate

Sex Applicability

Sex	Evidence
Unspecific	Moderate

Taxonomic: Effect data for the occurrence of this KE exist from *Pieris brassicae* and *Lucilia cuprina*. However, all arthropods undergo molting, so it is highly likely that this KE is applicable to the whole phylum of arthropods.

Life stage: This KE is applicable for organisms that undergo molting in order to grow and develop, namely larval stages of insects and all life stages of crustaceans and arachnids.

Sex: This KE is applicable to all sexes.

Chemical: Substances known to induce premature molting are of the family of pyrimidine nucleosides (e.g. polyoxin D and nikkomycin Z) (Gijswijt et al. 1979; Tellam et al. 2000; Arakawa et al. 2008).

Key Event Description

This key event is measured on the level of the individual. In order to grow and develop, arthropods need to shed their exoskeleton periodically (molting) (Heming 2018). During molting, the newly secreted cuticle is subject to mechanical stress associated and therefore needs to possess enough structural and functional integrity. The ecdysis motor program, which constitutes the behavioral part of the cuticle shedding requires the newly secreted cuticle to possess a certain strength to support for muscular force in order to shed the old cuticle (Ewer 2005). Cuticular integrity is also important after ecdysis, as insects and crustaceans expand their new cuticle by increasing internal pressure by swallowing air and water, respectively. This happens in order to expand and provide stability to the new cuticle until it is hardened (tanned) (Clarke 1957; Lee 1961; Dall et al. 1978; deFur et al. 1985). If arthropods are not able to molt properly, the organism will eventually die. Premature molting describes the unsuccessful molting where the organism is not able to shed the old cuticle, but also other effects related to molting in an immature stage where the new cuticle is not mature enough for the molt, such as rupture of the new cuticle and associated desiccation, deformities, higher susceptibility to pathogens or impaired locomotion. Specific effects observed are animals stuck in their exuviae (Wang et al., 2019), and if molting can be completed despite an immature cuticle, animals might be smaller and die at subsequent molts (Arakawa et al., 2008; Chen et al., 2008; Mohammed et al., 2017).

How it is Measured or Detected

Premature molting can be determined by observation. No standardized tests for the endpoint of molting exist to date. However, during an OECD 202 *Daphnia* sp. Acute immobilization test (OECD 2004), the cumulative number of molts can be assessed as an additional endpoint. Molting can also be assessed during a OECD 211 *Daphnia* sp. Reproduction test (OECD 2012), as proposed previously (OECD 2003). One could even prolong the test to 96h to get a clearer result of this endpoint. Additionally, one could apply histopathological methods to monitor the maturity of the newly synthesized cuticle (e.g. thickness of procuticle).

References

- Arakawa T, Yukuhiro F, Noda H. 2008. Insecticidal effect of a fungicide containing polyoxin B on the larvae of *Bombyx mori* (Lepidoptera: Bombycidae), *Mamestra brassicae*, *Mythimna separata*, and *Spodoptera litura* (Lepidoptera: Noctuidae). *Appl Entomol Zool.* 43(2):173–181. doi:10.1303/aez.2008.173.
- Chen, X.; Tian, H.; Zou, L.; Tang, B.; Hu, J.; Zhang, W. Disruption of *Spodoptera Exigua* Larval Development by Silencing Chitin Synthase Gene A with RNA Interference. *Bull. Entomol. Res.* 2008, 98 (6), 613–619. <https://doi.org/10.1017/S0007485308005932>.
- Clarke KU. 1957. On the Increase in Linear Size During Growth in *Locusta Migratoria* L. *Proc R Entomol Soc London Ser A, Gen Entomol.* 32(1–3):35–39. doi:10.1111/j.1365-3032.1957.tb00361.x.
- Dall W, Smith DM, Press B. 1978. Water uptake at ecdysis in the western rock lobster. *J Exp Mar Bio Ecol.* 35(1960). doi:10.1016/0022-0981(78)90074-6.
- deFur PL, Mangum CP, McMahon BR. 1985. Cardiovascular and Ventilatory Changes During Ecdysis in the Blue Crab *Callinectes Sapidus* Rathbun. *J Crustac Biol.* 5(2):207–215. doi:10.2307/1547867.
- Ewer J. 2005. How the ecdysozoan changed its coat. *PLoS Biol.* 3(10):1696–1699. doi:10.1371/journal.pbio.0030349.
- Gijswijt MJ, Deul DH, de Jong BJ. 1979. Inhibition of chitin synthesis by benzoyl-phenylurea insecticides, III. Similarity in action in *Pieris brassicae* (L.) with Polyoxin D. *Pestic Biochem Physiol.* 12(1):87–94. doi:10.1016/0048-3575(79)90098-1.
- Heming BS. 2018. *Insect development and evolution*. Ithaca: Cornell University Press.
- Mohammed, A. M. A.; DIab, M. R.; Abdelsattar, M.; Khalil, S. M. S. Characterization and RNAi-Mediated Knockdown of Chitin Synthase A in the Potato Tuber Moth, *Phthorimaea Operculella*. *Sci. Rep.* 2017, 7 (1), 1–12. <https://doi.org/10.1038/s41598-017-09858-y>.
- Lee RM. 1961. The variation of blood volume with age in the desert locust (*Schistocerca gregaria* Forsk.). *J Insect Physiol.* 6(1):36–51. doi:10.1016/0022-1910(61)90090-7.
- OECD (2003), Proposal for an Enhanced Test Guideline. *Daphnia magna* Reproduction Test. Draft OECD Guidel. Test. Chem. Enhanc. Tech. Guid. Doc. 211 21.
- OECD (2004), *Test No. 202: Daphnia sp. Acute Immobilisation Test*, OECD Guidelines for the Testing of Chemicals, Section 2, OECD Publishing, Paris, <https://doi.org/10.1787/9789264069947-en>.
- OECD (2012), *Test No. 211: Daphnia magna Reproduction Test*, OECD Guidelines for the Testing of Chemicals, Section 2, OECD Publishing, Paris, <https://doi.org/10.1787/9789264185203-en>.
- Tellam RL, Vuocolo T, Johnson SE, Jarmey J, Pearson RD. 2000. Insect chitin synthase. cDNA sequence, gene organization and expression. *Eur J Biochem.* 267(19):6025–6043. doi:10.1046/j.1432-1327.2000.01679.x.
- Wang, Z.; Yang, H.; Zhou, C.; Yang, W. J.; Jin, D. C.; Long, G. Y. Molecular Cloning, Expression, and Functional Analysis of the Chitin Synthase 1 Gene and Its Two Alternative Splicing Variants in the White-Backed Planthopper, *Sogatella Furcifera* (Hemiptera: Delphacidae). *Sci. Rep.* 2019, 9 (1), 1–14. <https://doi.org/10.1038/s41598-018-37488-5>.

List of Adverse Outcomes in this AOP

Event: 350: Increase, Mortality

Short Name: Increase, Mortality

Key Event Component

Process	Object	Action
mortality		increased

AOPs Including This Key Event

AOP ID and Name	Event Type
Aop:4 - Ecdysone receptor agonism leading to incomplete ecdysis associated mortality	Key Event
Aop:331 - Formation of DNA photoproducts leading to growth inhibition (1)	Adverse Outcome
Aop:327 - Excessive reactive oxygen species production leading to mortality (1)	Adverse Outcome
Aop:328 - Excessive reactive oxygen species production leading to mortality (2)	Adverse Outcome
Aop:329 - Excessive reactive oxygen species production leading to mortality (3)	Adverse Outcome
Aop:330 - Excessive reactive oxygen species production leading to mortality (4)	Adverse Outcome
Aop:343 - S-adenosylmethionine depletion leading to population decline (2)	Adverse Outcome
Aop:342 - S-adenosylmethionine depletion leading to population decline (1)	Adverse Outcome
Aop:358 - Chitinase inhibition leading to mortality	Adverse Outcome
Aop:359 - Chitobiase inhibition leading to mortality	Adverse Outcome
Aop:360 - Chitin synthase 1 inhibition leading to mortality	Adverse Outcome
Aop:361 - Sulfonyleureareceptor binding leading to mortality	Adverse Outcome

Stressors

Name
Polyoxin D
Nikkomycins

Biological Context

Level of Biological Organization
Individual

Domain of Applicability

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
Lucilia cuprina	Lucilia cuprina	High	NCBI
Daphnia magna	Daphnia magna	High	NCBI

Life Stage Applicability

Life Stage	Evidence
All life stages	High

Sex Applicability

Sex	Evidence
Unspecific	High

Taxonomic: This AO is applicable to all living organisms.

Life stage: This AO is applicable to all life stages.

Sex: This AO is applicable to all sexes.

Chemical: Substances known to increase mortality in arthropods are of the family of pyrimidine nucleosides (e.g. polyoxin D and nikkomycin Z) (Gijswijt et al. 1979; Tellam et al. 2000; Arakawa et al. 2008).

Key Event Description

This key event is observed at the biological level of the individual and describes the increase of mortality of individuals upon exposure to a stressor.

How it is Measured or Detected

The AO can be detected by observation, for example by immobilization of the respective organisms. There exist guidelines for the characterization of this AO in arthropods. For example, the OECD 202 Daphnia sp. Acute immobilization test (OECD 2004) which can also be modified depending on the effect one expects.

Regulatory Significance of the AO

The Adverse Outcome is highly significant from a regulatory point of view. It is employed as regulatory endpoint in most studies assessing the toxicity of stressors.

References

Arakawa T, Yukuhiro F, Noda H. 2008. Insecticidal effect of a fungicide containing polyoxin B on the larvae of *Bombyx mori* (Lepidoptera: Bombycidae), *Mamestra brassicae*, *Mythimna separata*, and *Spodoptera litura* (Lepidoptera: Noctuidae). *Appl Entomol Zool.* 43(2):173–181. doi:10.1303/aez.2008.173.

Gijswijt MJ, Deul DH, de Jong BJ. 1979. Inhibition of chitin synthesis by benzoyl-phenylurea insecticides, III. Similarity in action in *Pieris brassicae* (L.) with Polyoxin D. *Pestic Biochem Physiol.* 12(1):87–94. doi:10.1016/0048-3575(79)90098-1.

OECD. 2004. Test No. 202: *Daphnia sp.* Acute Immobilisation Test. OECD OECD Guidelines for the Testing of Chemicals, Section 2. [accessed 2020 Mar 3]. https://www.oecd-ilibrary.org/environment/test-no-202-daphnia-sp-acute-immobilisation-test_9789264069947-en.

Tellam RL, Vuocolo T, Johnson SE, Jarmey J, Pearson RD. 2000. Insect chitin synthase. cDNA sequence, gene organization and expression. *Eur J Biochem.* 267(19):6025–6043. doi:10.1046/j.1432-1327.2000.01679.x.

Appendix 2 - List of Key Event Relationships in the AOP

List of Adjacent Key Event Relationships

Relationship: 1742: Inhibition, CHS-1 leads to Decrease, Cuticular chitin content

AOPs Referencing Relationship

AOP Name	Adjacency	Weight of Evidence	Quantitative Understanding
S-adenosylmethionine depletion leading to population decline (1)	adjacent		
Chitin synthase 1 inhibition leading to mortality	adjacent	Moderate	Low

Evidence Supporting Applicability of this Relationship

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
crustaceans	Daphnia magna	Moderate	NCBI
insects	insects	Moderate	NCBI

Life Stage Applicability

Life Stage	Evidence
larvae	High
Juvenile	High
Adult	Moderate

Sex Applicability

Sex	Evidence
Unspecific	Moderate

Taxonomic: Likely, this KER is likely applicable to the whole phylum of arthropods as they all depend on the synthesis of chitin.

Life stage: This KER is applicable for organisms synthesizing chitin in order to grow and develop, namely larval stages of insects and all life stages of crustaceans and arachnids.

Sex: This KER is applicable to all sexes.

Chemical: Substances inducing both, the inhibition of CHS-1 and the decrease in cuticular chitin content are of the family of pyrimidine nucleosides (e.g. polyoxin D, polyoxin B and nikkomycin Z) (Gijswijt et al. 1979; Cohen and Casida 1982; Turnbull and Howells 1982; Calcott and Fatig 1984; Kuwano and Cohen 1984; Cohen and Casida 1990; Zhang and Yan Zhu 2013; Zhuo et al. 2014; Osada 2019). The phthalimide

captan was also shown to induce CHS-1 inhibition and a decrease in cuticular chitin content (Cohen and Casida 1982; Gelman and Borkovec 1986). However, studies assessing both endpoints in sequence are lacking.

Key Event Relationship Description

Chitin in the arthropod cuticle is synthesized by the chitin synthase isoform 1 (CHS-1) which spans the plasma membrane on the apical plasma membrane of epithelial cells (Locke and Huie 1979; Binnington 1985; Merzendorfer and Zimoch 2003; Merzendorfer 2006). Since CHS-1 is the enzyme to polymerize chitin from UDP-*N*-Acetylglucosamine (UDP-GlcNAc) (Merzendorfer 2006), it is solely responsible for the content of chitin in the exoskeleton. Consequently, the inhibition of CHS-1 leads to a decrease in chitin content in the arthropod cuticle.

Evidence Supporting this KER

Biological Plausibility

The process of chitin synthesis in arthropods is well characterized. Although the exact mechanism of the polymerization reaction remains elusive, CHS-1 is known to be the key enzyme in the biosynthesis of chitin and therefore, responsible for the cuticular chitin content (Merzendorfer and Zimoch 2003; Merzendorfer 2006). Therefore, the biological plausibility of this KER can be regarded as high.

Empirical Evidence

Empirical evidence for the occurrence of both KEs, the inhibition of CHS-1 and the decrease in cuticular chitin content exist. For example, the occurrence of chitin synthase inhibition was characterized using cell free crude enzyme preparations *in vitro* from coleopteran, lepidopteran and dipteran insect species upon treatment with polyoxin B, polyoxin D and nikkomycin Z (Cohen and Casida 1982; Kuwano and Cohen 1984; Cohen and Casida 1990; Zhang and Yan Zhu 2013). The cuticular chitin content was characterized *in vivo* in *Artemia salina* or using cultured integumental tissue from lepidopteran and dipteran species after exposure to polyoxin D and nikkomycin Z as well as the phthalimides captan, captafol, and folpet (Gijswijt et al. 1979; Turnbull and Howells 1982; Calcott and Fatig 1984; Gelman and Borkovec 1986; Zhuo et al. 2014). Data from studies with specific stressors assessing both endpoints and therefore supporting dose concordance of the KER are lacking. However, results from studies where CHS-1 was knocked down by RNA interference support temporal concordance of the KER (Arakane et al. 2005, Li et al. 2017, Zhang X. et al. 2010). Given the support for temporal concordance and the lack of studies showing dose concordance, the empirical evidence for this KER was judged as moderate.

Uncertainties and Inconsistencies

The major uncertainty in this KER is the absence of studies which assess both endpoints, the inhibition of the chitin synthase and the decrease in cuticular chitin content after exposure to specific stressors.

Quantitative Understanding of the Linkage

Response-response relationship

Due to the lack of studies linking the inhibition of CHS-1 to the decrease in cuticular chitin content, it is not possible to describe the nature of the response-response relationship.

Time-scale

Due to the lack of studies assessing the inhibition of CHS-1 and the decrease in cuticular chitin content, it is not possible to make a statement on the timescale of the relationship. However, the expression of CHS-1 peaks at the time of ecdysis (Ampasala et al. 2011; Wang et al. 2012), indicating the highest rate of chitin synthesis at this timepoint. Hence it can be assumed that a decrease in chitin content in the newly synthesized cuticle should become apparent shortly after. In studies where CHS-1 was knocked down,

chitin contents were assessed after 3 and 7 days and found to be decreased (Arakane et al. 2005, Li et al. 2017, Zhang X. et al. 2010).

Known modulating factors

CHS is dependent on bivalent ions as cofactor such as Mg^{2+} or Mn^{2+} (Merzendorfer 2006). Both low and high levels of Mg^{2+} inhibited CHS activity *in vitro* (Zhang and Yan Zhu 2013).

Known Feedforward/Feedback loops influencing this KER

Upon knockdown of CHS-1 in the salmon louse *Lepeophtheirus salmonis*, upregulation of the UDP-GlcNAc pyrophosphorylase (UAP), which catalyzes the conversion of GlcNAc to UDP-GlcNAc, was observed (Braden et al. 2020). The knockdown of UAP also led to upregulation of CHS-1 demonstrating a clear dependence of the two enzymes. Most likely, the upregulation of UAP is a compensatory mechanism with the goal to restore homeostasis in absence of CHS-1. The exact regulation of the feedback, however, remains to be investigated.

References

- Ampasala DR, Zheng S, Zhang D, Ladd T, Doucet D, Krell PJ, Retnakaran A, Feng Q. 2011. An epidermis-specific chitin synthase cDNA in *Choristoneura fumiferana*: Cloning, characterization, developmental and hormonal-regulated expression. *Arch Insect Biochem Physiol.* 76(2):83–96. doi:10.1002/arch.20404.
- Arakane, Y.; Muthukrishnan, S.; Kramer, K. J.; Specht, C. A.; Tomoyasu, Y.; Lorenzen, M. D.; Kanost, M.; Beeman, R. W. The *Tribolium* Chitin Synthase Genes TcCHS1 and TcCHS2 Are Specialized for Synthesis of Epidermal Cuticle and Midgut Peritrophic Matrix. *Insect Mol. Biol.* 2005, 14 (5), 453–463. <https://doi.org/10.1111/j.1365-2583.2005.00576.x>.
- Binnington KC. 1985. Ultrastructural changes in the cuticle of the sheep blowfly, *Lucilia*, induced by certain insecticides and biological inhibitors. *Tissue Cell.* 17(1):131–140. doi:10.1016/0040-8166(85)90021-7.
- Braden L, Michaud D, Igboeli OO, Dondrup M, Hamre L, Dalvin S, Purcell SL, Kongshaug H, Eichner C, Nilsen F, et al. 2020. Identification of critical enzymes in the salmon louse chitin synthesis pathway as revealed by RNA interference-mediated abrogation of infectivity. *Int J Parasitol.* 50(10–11):873–889. doi:10.1016/j.ijpara.2020.06.007. <https://doi.org/10.1016/j.ijpara.2020.06.007>.
- Calcott PH, Fatig RO. 1984. Inhibition of Chitin metabolism by Avermectin in susceptible Organisms. *J Antibiot (Tokyo).* 37(3):253–259. doi:10.7164/antibiotics.37.253.
- Cohen E, Casida JE. 1982. Properties and inhibition of insect integumental chitin synthetase. *Pestic Biochem Physiol.* 17(3):301–306. doi:10.1016/0048-3575(82)90141-9.
- Cohen E, Casida JE. 1990. Insect and Fungal Chitin Synthetase Activity: Specificity of Lectins as Enhancers and Nucleoside Peptides as Inhibitors. *Pestic Biochem Physiol.* 37(3):249–253. doi:10.1016/0048-3575(90)90131-K.
- Gelman DB, Borkovec AB. 1986. The pharate adult clasper as a tool for measuring chitin synthesis and for identifying new chitin synthesis inhibitors. *Comp Biochem Physiol Part C, Comp.* 85(1):193–197. doi:10.1016/0742-8413(86)90073-3.
- Gijswijt MJ, Deul DH, de Jong BJ. 1979. Inhibition of chitin synthesis by benzoyl-phenylurea insecticides, III. Similarity in action in *Pieris brassicae* (L.) with Polyoxin D. *Pestic Biochem Physiol.* 12(1):87–94. doi:10.1016/0048-3575(79)90098-1.
- Kuwano E, Cohen E. 1984. The use of a *Tribolium* chitin synthetase assay in studying the effects of benzimidazoles with a terpene moiety and related compounds. *Agric Biol Chem.* 48(6):1617–1620. doi:10.1080/00021369.1984.10866362.
- Li, T.; Chen, J.; Fan, X.; Chen, W.; Zhang, W. MicroRNA and DsRNA Targeting Chitin Synthase A Reveal a Great Potential for Pest Management of the Hemipteran Insect *Nilaparvata lugens*. *Pest Manag. Sci.* 2017, 73 (7), 1529–1537. <https://doi.org/10.1002/ps.4492>.
- Locke M, Huie P. 1979. Apolysis and the Turnover of Plasmamembrane Plaques during Cuticle formation in an Insect. *Tissue Cell.* 11(2):277–291. doi:10.1016/0040-8166(79)90042-9.
- Merzendorfer H. 2006. Insect chitin synthases: A review. *J Comp Physiol B Biochem Syst Environ Physiol.* doi:10.1007/s00360-005-0005-3.
- Merzendorfer H, Zimoch L. 2003. Chitin metabolism in insects: structure, function and regulation of chitin synthases and chitinases. *J Exp Biol.* 206(24):4393 LP – 4412. doi:10.1242/jeb.00709. <http://jeb.biologists.org/content/206/24/4393.abstract>.
- Osada H. 2019. Discovery and applications of nucleoside antibiotics beyond polyoxin. *J Antibiot (Tokyo).* 72(12):855–864. doi:10.1038/s41429-019-0237-1. <http://dx.doi.org/10.1038/s41429-019-0237-1>.

Turnbull IF, Howells AJ. 1982. Effects of several larvicidal compounds on chitin biosynthesis by isolated larval integuments of the sheep blowfly *Lucilia cuprina*. Aust J Biol Sci. 35(5):491–504. doi:10.1071/B19820491.

Wang Y, Fan HW, Huang HJ, Xue J, Wu WJ, Bao YY, Xu HJ, Zhu ZR, Cheng JA, Zhang CX. 2012. Chitin synthase 1 gene and its two alternative splicing variants from two sap-sucking insects, *Nilaparvata lugens* and *Laodelphax striatellus* (Hemiptera: Delphacidae). Insect Biochem Mol Biol. 42(9):637–646. doi:10.1016/j.ibmb.2012.04.009. <http://dx.doi.org/10.1016/j.ibmb.2012.04.009>.

Zhang, X.; Zhang, J.; Zhu, K. Y. Chitosan/Double-Stranded RNA Nanoparticle-Mediated RNA Interference to Silence Chitin Synthase Genes through Larval Feeding in the African Malaria Mosquito (*Anopheles Gambiae*). Insect Mol. Biol. 2010, 19 (5), 683–693. <https://doi.org/10.1111/j.1365-2583.2010.01029.x>.

Zhang X, Yan Zhu K. 2013. Biochemical characterization of chitin synthase activity and inhibition in the African malaria mosquito, *Anopheles gambiae*. Insect Sci. 20(2):158–166. doi:10.1111/j.1744-7917.2012.01568.x.

Zhuo W, Fang Y, Kong L, Li X, Sima Y, Xu S. 2014. Chitin synthase A: A novel epidermal development regulation gene in the larvae of *Bombyx mori*. Mol Biol Rep. 41(7):4177–4186. doi:10.1007/s11033-014-3288-1.

Relationship: 1743: Decrease, Cuticular chitin content leads to Increase, Premature molting

AOPs Referencing Relationship

AOP Name	Adjacency	Weight of Evidence	Quantitative Understanding
S-adenosylmethionine depletion leading to population decline (2)	adjacent		
S-adenosylmethionine depletion leading to population decline (1)	adjacent		
Chitin synthase 1 inhibition leading to mortality	adjacent	Moderate	Low
Sulfonylureareceptor binding leading to mortality	adjacent	Moderate	Moderate

Evidence Supporting Applicability of this Relationship

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
crustaceans	Daphnia magna	Moderate	NCBI
insects	insects	Moderate	NCBI

Life Stage Applicability

Life Stage	Evidence
larvae	High
Juvenile	High
Adult	Moderate

Sex Applicability

Sex	Evidence
Unspecific	Moderate

Taxonomic: In all likelihood, this KER is applicable to the whole phylum of arthropods as they all depend on the synthesis of chitin and molting in order to develop.

Life stage: This KER is applicable for organisms synthesizing chitin and molting in order to grow and develop, namely larval stages of insects and all life stages of crustaceans and arachnids.

Sex: This KER is applicable to all sexes.

Chemical: Occurrence of a decrease in cuticular chitin content as well as premature molting was observed after treatment with the pyrimidine nucleosides polyoxin D, polyoxin B and nikkomycin Z (Gijswijt et al. 1979; Turnbull and Howells 1982; Calcott and Fatig 1984; Gelman and Borkovec 1986; Tellam et al. 2000; Arakawa et al. 2008; Zhuo et al. 2014). However, studies causally linking both endpoints are lacking.

Key Event Relationship Description

As the arthropod cuticle is a central part in the molting process, its proper composition is indispensable for a proper molt. The ecdysis motor program, the behavioral part of ecdysis, constitutes a distinct motor pattern to split and shed the old cuticle (Ayali 2009). As the cuticle supports muscular function (Vincent and Wegst 2004), it needs to possess a certain integrity in order to successfully molt. The integrity of the cuticle is also important after ecdysis as arthropods, such as insects and crustaceans, expand the new cuticle by swallowing air or water in order to build up pressure to split the old and expand the new exoskeleton and provide stability to the soft new cuticle (Clarke 1957; Lee 1961; Dall et al. 1978; deFur et al. 1985). The arthropod cuticle mostly consists of chitin embedded in and crosslinked with a matrix of proteins (Muthukrishnan et al. 2012). If the chitin content is too low, the cuticle may not possess enough integrity to support muscular function or withstand the beforementioned stresses of ecdysis, which leads to the organism being stuck in the old cuticle or the rupture of the new cuticle.

Evidence Supporting this KER

Biological Plausibility

The ecdysis motor program, the behavioral part of ecdysis, constitutes a distinct motor pattern to split and shed the old cuticle (Ayali 2009). As the cuticle supports muscular function (Vincent and Wegst 2004), it needs to possess a certain integrity in order to successfully molt. The integrity of the cuticle is also important after ecdysis as arthropods, such as insects and crustaceans, expand the new cuticle by swallowing air or water in order to build up pressure to expand the new exoskeleton and provide stability to the soft new cuticle (Clarke 1957; Lee 1961; Dall et al. 1978; deFur et al. 1985). The arthropod cuticle mostly consists of chitin embedded in and crosslinked with a matrix of proteins (Muthukrishnan et al. 2012). Given the well biological understanding of the processes, the biological plausibility can be regarded as high.

Empirical Evidence

The cuticular chitin content was characterized *in vivo* in *Artemia salina* or using cultured integumental tissue from lepidopteran and dipteran insect species after exposure to polyoxin D and nikkomycin Z as well as the phthalimides captan, captafol, and folpet (Gijswijt et al. 1979; Turnbull and Howells 1982; Calcott and Fatig 1984; Gelman and Borkovec 1986; Zhuo et al. 2014). The event of premature molting was not assessed as endpoint in studies involving specific stressors rather than mentioned after exposure to polyoxin D, polyoxin B and nikkomycin Z (Gijswijt et al. 1979; Tellam et al. 2000; Arakawa et al. 2008). However, results from studies where CHS-1 was knocked down by RNA interference support temporal concordance of the KER (Arakane et al. 2005, Li et al. 2017, Zhang X. et al. 2010). Given the support for temporal concordance and the lack of studies showing dose concordance, the empirical evidence for this KER was judged as moderate.

Uncertainties and Inconsistencies

The absence of studies (quantitatively) assessing premature molting constitutes a major data gap. A further data gap is the absence of studies which assess both, the decrease in cuticular chitin content and the increase in premature molting.

Quantitative Understanding of the Linkage

Response-response relationship

Due to the lack of studies linking the decrease in cuticular chitin content with the increase in premature molting, it is not possible to describe the nature of the response-response relationship.

Time-scale

Due to the nature of the process, premature molting onsets at the time of ecdysis after the decrease in cuticular chitin content.

References

- Arakane, Y.; Muthukrishnan, S.; Kramer, K. J.; Specht, C. A.; Tomoyasu, Y.; Lorenzen, M. D.; Kanost, M.; Beeman, R. W. The Tribolium Chitin Synthase Genes TcCHS1 and TcCHS2 Are Specialized for Synthesis of Epidermal Cuticle and Midgut Peritrophic Matrix. *Insect Mol. Biol.* 2005, 14 (5), 453–463. <https://doi.org/10.1111/j.1365-2583.2005.00576.x>.
- Arakawa T, Yukuhiro F, Noda H. 2008. Insecticidal effect of a fungicide containing polyoxin B on the larvae of *Bombyx mori* (Lepidoptera: Bombycidae), *Mamestra brassicae*, *Mythimna separata*, and *Spodoptera litura* (Lepidoptera: Noctuidae). *Appl Entomol Zool.* 43(2):173–181. doi:10.1303/aez.2008.173.
- Ayali A. 2009. The role of the arthropod stomatogastric nervous system in moulting behaviour and ecdysis. *J Exp Biol.* 212(4):453–459. doi:10.1242/jeb.023879.
- Calcott PH, Fatig RO. 1984. Inhibition of Chitin metabolism by Avermectin in susceptible Organisms. *J Antibiot (Tokyo).* 37(3):253–259. doi:10.7164/antibiotics.37.253.
- Clarke KU. 1957. On the Increase in Linear Size During Growth in *Locusta Migratoria* L. *Proc R Entomol Soc London Ser A, Gen Entomol.* 32(1–3):35–39. doi:10.1111/j.1365-3032.1957.tb00361.x.
- Dall W, Smith DM, Press B. 1978. Water uptake at ecdysis in the western rock lobster. *J Exp Mar Bio Ecol.* 35(1960). doi:10.1016/0022-0981(78)90074-6.
- deFur PL, Mangum CP, McMahon BR. 1985. Cardiovascular and Ventilatory Changes During Ecdysis in the Blue Crab *Callinectes Sapidus* Rathbun. *J Crustac Biol.* 5(2):207–215. doi:10.2307/1547867.
- Gelman DB, Borkovec AB. 1986. The pharate adult clasper as a tool for measuring chitin synthesis and for identifying new chitin synthesis inhibitors. *Comp Biochem Physiol Part C, Comp.* 85(1):193–197. doi:10.1016/0742-8413(86)90073-3.
- Gijswijt MJ, Deul DH, de Jong BJ. 1979. Inhibition of chitin synthesis by benzoyl-phenylurea insecticides, III. Similarity in action in *Pieris brassicae* (L.) with Polyoxin D. *Pestic Biochem Physiol.* 12(1):87–94. doi:10.1016/0048-3575(79)90098-1.
- Lee RM. 1961. The variation of blood volume with age in the desert locust (*Schistocerca gregaria* Forsk.). *J Insect Physiol.* 6(1):36–51. doi:10.1016/0022-1910(61)90090-7.
- Li, T.; Chen, J.; Fan, X.; Chen, W.; Zhang, W. MicroRNA and DsRNA Targeting Chitin Synthase A Reveal a Great Potential for Pest Management of the Hemipteran Insect Nilaparvata Lugens. *Pest Manag. Sci.* 2017, 73 (7), 1529–1537. <https://doi.org/10.1002/ps.4492>.
- Muthukrishnan S, Merzendorfer H, Arakane Y, Kramer KJ. 2012. Chitin Metabolism in Insects. Elsevier B.V. <http://dx.doi.org/10.1016/B978-0-12-384747-8.10007-8>.
- Tellam RL, Vuocolo T, Johnson SE, Jarmey J, Pearson RD. 2000. Insect chitin synthase. cDNA sequence, gene organization and expression. *Eur J Biochem.* 267(19):6025–6043. doi:10.1046/j.1432-1327.2000.01679.x.
- Turnbull IF, Howells AJ. 1982. Effects of several larvicidal compounds on chitin biosynthesis by isolated larval integuments of the sheep blowfly *Lucilia cuprina*. *Aust J Biol Sci.* 35(5):491–504. doi:10.1071/BI9820491.
- Vincent JFV, Wegst UGK. 2004. Design and mechanical properties of insect cuticle. *Arthropod Struct Dev.* 33(3):187–199. doi:10.1016/j.asd.2004.05.006.
- Zhang, X.; Zhang, J.; Zhu, K. Y. Chitosan/Double-Stranded RNA Nanoparticle-Mediated RNA Interference to Silence Chitin Synthase Genes through Larval Feeding in the African Malaria Mosquito (*Anopheles Gambiae*). *Insect Mol. Biol.* 2010, 19 (5), 683–693. <https://doi.org/10.1111/j.1365-2583.2010.01029.x>.

Zhuo W, Fang Y, Kong L, Li X, Sima Y, Xu S. 2014. Chitin synthase A: A novel epidermal development regulation gene in the larvae of *Bombyx mori*. Mol Biol Rep. 41(7):4177–4186. doi:10.1007/s11033-014-3288-1.

Relationship: 1744: Increase, Premature molting leads to Increase, Mortality

AOPs Referencing Relationship

AOP Name	Adjacency	Weight of Evidence	Quantitative Understanding
S-adenosylmethionine depletion leading to population decline (2)	adjacent		
S-adenosylmethionine depletion leading to population decline (1)	adjacent		
Chitinase inhibition leading to mortality	adjacent	Moderate	Low
Chitobiase inhibition leading to mortality	adjacent	Moderate	Low
Chitin synthase 1 inhibition leading to mortality	adjacent	Moderate	Low
Sulfonylureareceptor binding leading to mortality	adjacent	High	High

Evidence Supporting Applicability of this Relationship

Taxonomic Applicability

Term	Scientific Term	Evidence	Links
crustaceans	Daphnia magna	Moderate	NCBI
insects	insects	Moderate	NCBI

Life Stage Applicability

Life Stage	Evidence
larvae	High
Juvenile	Moderate
Adult	Moderate

Sex Applicability

Sex	Evidence
Unspecific	Moderate

Taxonomic: Likely, this KER is applicable to the whole phylum of arthropods as they all depend on molting in order to develop.

Life stage: This KER is applicable for organisms molting in order to grow and develop, namely larval stages of insects and all life stages of crustaceans and arachnids.

Sex: This KER is applicable to all sexes.

Chemical: Occurrence of premature molting and an increase in mortality observed after treatment with the pyrimidine nucleosides (e.g. polyoxin D, polyoxin B and nikkomycin Z) (Gijswijt et al. 1979; Tellam et al. 2000; Tellam and Eisemann 2000; Arakawa et al. 2008; New Zealand Environmental Protection Authority 2015). However, studies causally linking both endpoints are lacking.

Key Event Relationship Description

During molting, arthropods pause food uptake and in certain cases also respiration (Camp et al. 2014; Song et al. 2017a). If molting is disrupted and the organism is not able to shed the old exoskeleton, the organism may eventually die of starvation, suffocation or the rupture of the exoskeleton.

Evidence Supporting this KER

Biological Plausibility

In order to grow and develop, arthropods need to molt periodically (Heming 2018). Since molting is a determining point in arthropod development, the disruption of molting leads to increased mortality (Arakawa et al. 2008; Merzendorfer et al. 2012; Song et al. 2017a; Song et al. 2017b). During ecdysis, arthropods pause food intake and respiration (Camp et al. 2014; Song et al. 2017a). Therefore, if the molt cannot be completed, the organism may die of starvation or suffocation. Additionally, if the cuticle is immature, it may not withstand the stresses associated with ecdysis (Clarke 1957; Lee 1961; Dall et al. 1978; deFur et al. 1985), and the organism may die of desiccation or increased susceptibility to pathogens. Given the well understood biological processes, the biological plausibility of this KER was rated as high.

Empirical Evidence

The event of premature molting is not well characterized. It gets mentioned as cause of death in studies with *Pieris brassicae*, *Spodoptera litura*, *Bombyx mori* and *Lucilia cuprina* after treatment with polyoxin D, polyoxin B, polyoxin AL (a mixture of polyoxins) and nikkomycin Z (Gijswijt et al. 1979; Tellam et al. 2000; Arakawa et al. 2008). The increase in mortality was reported in studies with *Lucilia cuprina*, *Spodoptera litura* and *Bombyx mori* (Tellam et al. 2000; Tellam and Eisemann 2000; Arakawa et al. 2008). Evidence from studies which assess and link both endpoints, and therefore would support dose concordance, is lacking. However, results from studies where CHS-1 was knocked down by RNA interference support temporal concordance of the KER (Arakane et al. 2005, Li et al. 2017, Chen et al., 2008; Mohammed et al., 2017; Shang et al., 2016; Wang et al., 2012, 2019; Yang et al., 2013; Ye et al., 2019; Zhai et al., 2017; Zhang et al., 2010). Given the support for temporal concordance and the lack of studies showing dose concordance, the empirical evidence for this KER was judged as moderate.

Uncertainties and Inconsistencies

The absence of studies (quantitatively) assessing premature molting constitutes a major data gap. A further data gap is the absence of studies which assess both, increase in premature molting and the increase in mortality are lacking.

Quantitative Understanding of the Linkage

Response-response relationship

Due to the lack of studies linking the increase in premature molting with the increase in mortality, it is not possible to describe the nature of the response-response relationship.

Time-scale

Death occurs after premature molting. However, an exact time frame in which death occurs cannot be defined yet.

References

- Arakawa T, Yukuhiro F, Noda H. 2008. Insecticidal effect of a fungicide containing polyoxin B on the larvae of *Bombyx mori* (Lepidoptera: Bombycidae), *Mamestra brassicae*, *Mythimna separata*, and *Spodoptera litura* (Lepidoptera: Noctuidae). *Appl Entomol Zool.* 43(2):173–181. doi:10.1303/aez.2008.173.
- Camp AA, Funk DH, Buchwalter DB. 2014. A stressful shortness of breath: Molting disrupts breathing in the mayfly *Cloeon dipterum*. *Freshw Sci.* 33(3):695–699. doi:10.1086/677899.
- Chen, X.; Tian, H.; Zou, L.; Tang, B.; Hu, J.; Zhang, W. Disruption of *Spodoptera Exigua* Larval Development by Silencing Chitin Synthase Gene A with RNA Interference. *Bull. Entomol. Res.* 2008, 98 (6), 613–619. <https://doi.org/10.1017/S0007485308005932>.
- Mohammed, A. M. A.; DIab, M. R.; Abdelsattar, M.; Khalil, S. M. S. Characterization and RNAi-Mediated Knockdown of Chitin Synthase A in the Potato Tuber Moth, *Phthorimaea Operculella*. *Sci. Rep.* 2017, 7 (1), 1–12. <https://doi.org/10.1038/s41598-017-09858-y>.
- Clarke KU. 1957. On the Increase in Linear Size During Growth in *Locusta Migratoria* L. *Proc R Entomol Soc London Ser A, Gen Entomol.* 32(1–3):35–39. doi:10.1111/j.1365-3032.1957.tb00361.x.
- Dall W, Smith DM, Press B. 1978. Water uptake at ecdysis in the western rock lobster. *J Exp Mar Bio Ecol.* 35(1960). doi:10.1016/0022-0981(78)90074-6.
- deFur PL, Mangum CP, McMahon BR. 1985. Cardiovascular and Ventilatory Changes During Ecdysis in the Blue Crab *Callinectes Sapidus* Rathbun. *J Crustac Biol.* 5(2):207–215. doi:10.2307/1547867.
- Gijswijt MJ, Deul DH, de Jong BJ. 1979. Inhibition of chitin synthesis by benzoyl-phenylurea insecticides, III. Similarity in action in *Pieris brassicae* (L.) with Polyoxin D. *Pestic Biochem Physiol.* 12(1):87–94. doi:10.1016/0048-3575(79)90098-1.
- Heming BS. 2018. *Insect development and evolution*. Ithaca: Cornell University Press.
- Lee RM. 1961. The variation of blood volume with age in the desert locust (*Schistocerca gregaria* Forsk.). *J Insect Physiol.* 6(1):36–51. doi:10.1016/0022-1910(61)90090-7.
- Merzendorfer H, Kim HS, Chaudhari SS, Kumari M, Specht CA, Butcher S, Brown SJ, Robert Manak J, Beeman RW, Kramer KJ, et al. 2012. Genomic and proteomic studies on the effects of the insect growth regulator diflubenzuron in the model beetle species *Tribolium castaneum*. *Insect Biochem Mol Biol.* 42(4):264–276. doi:10.1016/j.ibmb.2011.12.008. <http://dx.doi.org/10.1016/j.ibmb.2011.12.008>.
- New Zealand Environmental Protection Authority. 2015. Application for approval to import ESTEEM for release. <https://www.epa.govt.nz/assets/FileAPI/hsno-ar/APP202334/fbce9a39e6/APP202334-APP202334-Staff-Report-Final-updated.pdf>.
- Shang, F.; Xiong, Y.; Xia, W. K.; Wei, D. D.; Wei, D.; Wang, J. J. Identification, Characterization and Functional Analysis of a Chitin Synthase Gene in the Brown Citrus Aphid, *Toxoptera Citricida* (Hemiptera, Aphididae). *Insect Mol. Biol.* 2016, 25 (4), 422–430. <https://doi.org/10.1111/imb.12228>.
- Song Y, Evenseth LM, Iguchi T, Tollefsen KE. 2017b. Release of chitobiase as an indicator of potential molting disruption in juvenile *Daphnia magna* exposed to the ecdysone receptor agonist 20-hydroxyecdysone. *J Toxicol Environ Heal - Part A Curr Issues.* 80(16–18):954–962. doi:10.1080/15287394.2017.1352215. <https://doi.org/10.1080/15287394.2017.1352215>.
- Song Y, Villeneuve DL, Toyota K, Iguchi T, Tollefsen KE. 2017a. Ecdysone Receptor Agonism Leading to Lethal Molting Disruption in Arthropods: Review and Adverse Outcome Pathway Development. *Environ Sci Technol.* 51(8):4142–4157. doi:10.1021/acs.est.7b00480.
- Tellam RL, Eisemann C. 2000. Chitin is only a minor component of the peritrophic matrix from larvae of *Lucilia cuprina*. *Insect Biochem Mol Biol.* 30(12):1189–1201. doi:10.1016/S0965-1748(00)00097-7.

Tellam RL, Vuocolo T, Johnson SE, Jarmey J, Pearson RD. 2000. Insect chitin synthase. cDNA sequence, gene organization and expression. *Eur J Biochem.* 267(19):6025–6043. doi:10.1046/j.1432-1327.2000.01679.x.

Wang, Z.; Yang, H.; Zhou, C.; Yang, W. J.; Jin, D. C.; Long, G. Y. Molecular Cloning, Expression, and Functional Analysis of the Chitin Synthase 1 Gene and Its Two Alternative Splicing Variants in the White-Backed Planthopper, *Sogatella Furcifera* (Hemiptera: Delphacidae). *Sci. Rep.* 2019, 9 (1), 1–14. <https://doi.org/10.1038/s41598-018-37488-5>.

Wang, Y.; Fan, H. W.; Huang, H. J.; Xue, J.; Wu, W. J.; Bao, Y. Y.; Xu, H. J.; Zhu, Z. R.; Cheng, J. A.; Zhang, C. X. Chitin Synthase 1 Gene and Its Two Alternative Splicing Variants from Two Sap-Sucking Insects, *Nilaparvata Lugens* and *Laodelphax Striatellus* (Hemiptera: Delphacidae). *Insect Biochem. Mol. Biol.* 2012, 42 (9), 637–646. <https://doi.org/10.1016/j.ibmb.2012.04.009>.

Yang, W. J.; Xu, K. K.; Cong, L.; Wang, J. J. Identification, mRNA Expression, and Functional Analysis of Chitin Synthase 1 Gene and Its Two Alternative Splicing Variants in Oriental Fruit Fly, *Bactrocera Dorsalis*. *Int. J. Biol. Sci.* 2013, 9 (4), 331–342. <https://doi.org/10.7150/ijbs.6022>.

Ye, C.; Jiang, Y. Di; An, X.; Yang, L.; Shang, F.; Niu, J.; Wang, J. J. Effects of RNAi-Based Silencing of Chitin Synthase Gene on Moulting and Fecundity in Pea Aphids (*Acyrtosiphon Pisum*). *Sci. Rep.* 2019, 9 (1), 1–10. <https://doi.org/10.1038/s41598-019-39837-4>.

Zhai, Y.; Fan, X.; Yin, Z.; Yue, X.; Men, X.; Zheng, L.; Zhang, W. Identification and Functional Analysis of Chitin Synthase A in Oriental Armyworm, *Mythimna Separata*. *Proteomics* 2017, 17 (21), 1–11. <https://doi.org/10.1002/pmic.201700165>.

Zhang, J. et al. Silencing of two alternative splicing-derived mRNA variants of chitin synthase 1 gene by RNAi is lethal to the oriental migratory locust, *Locusta migratoria manilensis* (Meyen). *Insect Biochem. Mol. Biol.* 40, 824–833 (2010).