

1 *Guidance Document on Laboratory Product Performance*
2 *Testing Methods for Bed Bug Biocide Products*

DRAFT

Table of contents

4	Laboratory Product Performance Testing Methods for Bed Bug Biocide Products	1
5	1. Scope of Guidance Document.....	3
6	2. Definitions	4
7	3. Development of protocols for bed bug studies	6
8	3.1. Scientific design of research.	6
9	3.2. Data collection and reporting.....	9
10	4. Specific Guidance	12
11	4.1. Specific guidance for laboratory studies for resistance ratio determination and	
12	characterization of bed bug strain susceptibility.....	12
13	4.2. Specific guidance for laboratory studies for forced exposure (no-choice) residual surface	
14	treatments.....	14
15	4.3. Specific guidance for laboratory studies to determine the product performance of insecticide	
16	products under practical use situation.....	15
17	4.4. Specific guidance for laboratory studies to determine the product performance of a repellent	
18	or attractant product.	17
19	4.5. Specific guidance for laboratory studies for testing indoor insecticide total release aerosols,	
20	space sprays, and insecticide vapor strip products.....	20
21	4.6. Specific guidance for laboratory studies for direct application testing of insecticide products. .	22
22	4.7. Specific guidance for laboratory studies for testing ovicidal products.	22
23	4.8. Specific guidance for laboratory studies for fumigant products against all bed bug life stages. .	24
24	4.9. Specific guidance for laboratory studies of insect growth regulators (IGRs).	25
25	5. References	27

29

1. Scope of Guidance Document

30 This guidance provides recommendations for the design and execution of laboratory studies
31 to evaluate the performance of biocidal products in any formulation such as a liquid,
32 aerosol, fog, or impregnated fabric intended to repel, attract, and/or kill bed bugs (*Cimex*
33 *lectularius*). It does not apply to repellent products applied to human skin. The guidance is
34 based upon the American Laboratory Product Performance Testing Methods for Bed Bug
35 Pesticide Products (US EPA 2017, OCSPP 810.3900) and incorporates information from
36 laboratory efficacy testing standards for biocidal products against bed bugs in the
37 framework of the German Infectious Diseases Protection Act (18, 46). Investigators should
38 ensure research is conducted in compliance with any applicable laws or regulations, which
39 are independent of and additional to those cited in this guidance.

40

DRAFT

41

2. Definitions

42 The following definitions are of special importance in understanding this guidance
43 document. They apply only in the context of this guidance and are not intended to be more
44 generally applicable.

45

46 1. A bed bug **attractant** is a substance that causes bed bugs to make oriented
47 movements towards its source.

48 2. A **biocide** is a product with an active substance or with a formulation containing
49 one or more active substances intended to destroy, deter or attract, render harmless,
50 or prevent the action of, or otherwise exert a controlling effect on any harmful
51 organism by chemical or biological means.

52 3. **Behavioural resistance** refers to the avoidance by arthropods of a toxin through
53 detection or recognition (22).

54 4. A product that **controls bed bugs** demonstrates that the insecticide application
55 shows a sufficient efficacy (e.g. mortality and/or knockdown/morbidity followed
56 by mortality and/or residual efficacy). See also the definition for “residual
57 efficacy.”

58 5. **Crossing** is the act of passage by a bed bug from an untreated surface to a treated
59 surface or over a treated surface to another untreated surface.

60 6. The **EC₅₀** of a dose response curve represents the effective concentration of an
61 insecticide needed to knockdown 50% of the test population, after a specified
62 exposure time.

63 7. The **EC₉₀** of a dose response curve represents the effective concentration of an
64 insecticide needed to knockdown 90% of the test population, after a specified
65 exposure time.

66 8. A **fumigant** is an insecticide that is applied in the gaseous state or that forms a gas.
67 Fumigants may be used to kill bed bugs indoors or in containers.

68 9. A bed bug **harborage** is a sheltered area or refuge for bed bugs.

69 10. A bed bug **host** is a mammal (especially humans) or bird that bed bugs bite to obtain
70 blood for their survival and reproduction.

71 11. **Host-seeking** of bed bugs is the behaviour of bed bugs actively seeking a host.

72 12. An **insecticide** is a substance that kills insect pests.

73 13. An **Insect Growth Regulator (IGR)** is an insecticide that inhibits the maturation of
74 a bed bug through its life cycle.

75 14. Bed bug **knockdown** refers to a bed bug that is rendered incapable of coordinated
76 movement or unable to right itself following exposure to a biocidal product. In
77 contrast to “moribund”, bed bugs considered as knocked down have the potential
78 to recover during evaluation observations.

79 15. **LD₅₀** is a measure of lethality of a given toxicant calculated as the dose of toxicant
80 needed to kill 50% of a test population.

- 81 16. **LD₉₀** is a measure of lethality of a given toxicant calculated as the dose of toxicant
82 needed to kill 90% of a test population.
- 83 17. **Moribund** refers to bed bugs that are on their backs with only slight muscle
84 twitching of the extremities. Bed bugs exhibiting this behaviour may not be
85 considered dead. In contrast to “knockdown”, bed bugs considered as moribund
86 will not recover.
- 87 18. **Mortality** refers to bed bug death. A dead bed bug is a bed bug that does not move,
88 even when poked or probed.
- 89 19. An **ovicidal product** is an insecticide product that kills bed bug eggs.
- 90 20. **Insecticide resistance** is a heritable decrease in the susceptibility of a pest strain or
91 population to a given insecticide. This change is revealed in the repeated failure of
92 a product to achieve the expected level of mortality when used according to the
93 label directions for that pest species (22).
- 94 21. A **positive control** is a treatment with a well-known effective biocide.
- 95 22. A **repellent** is a substance, causing bed bugs to avoid a treated substrate (e.g.
96 disrupting the host-seeking or shelter-seeking behaviour of bed bugs).
- 97 23. **Residual efficacy** refers to a surface or space treated with a biocidal product
98 continuing to provide the intended biocidal effect at an acceptable level for an
99 extended length of time after application. The product’s residues should be
100 effective for at least 24 hours post application.
- 101 24. **Resistance Ratio (RR)** is a quantitative expression of the resistance of a bed bug
102 strain to a specific active ingredient or product formulation. A resistance ratio (e.g.
103 RR50) is calculated by dividing a quantitative measure of the lethality of a given
104 insecticide (e.g. LD50 value) for a bed bug strain of unknown level of resistance
105 by the corresponding measure of lethality for a strain known to be susceptible to
106 the given insecticide. A resistance ratio equal to or greater than 100 is characteristic
107 of a resistance strain. It has to be acknowledged, that practical bed bug control may
108 fail even in bed bug populations with a resistance ratio < 100.
- 109 25. A **resistant bed bug** population/strain refers to bed bugs that survive an insecticide
110 dose known to kill or control bed bugs.
- 111 26. A **susceptible bed bug** population/strain refers to bed bugs that exhibit mortality
112 when exposed to an appropriate dose of an effective insecticide for bed bugs.
113 Ideally, susceptible bed bugs will not have a history of exposure to the biocidal
114 mode of action being tested.
- 115

116

3. Development of protocols for bed bug studies

117 The first major stage of bed bug product testing is the development of a study protocol.
 118 General considerations in developing a study protocol for bed bug studies include scientific
 119 design of the study, data collection, data analysis, and reporting. Each of these topics is
 120 discussed in more detail in the sub-sections below.

121 3.1. Scientific design of research.

122 To be scientifically justified, the development of a new test protocol should address an
 123 important research or testing issue that cannot be answered by existing data. In addition,
 124 the new test design should be likely to provide a definitive answer to the research question.
 125 A detailed description of the experimental design and bed bug rearing (h) should be
 126 included.

127 a. **Objectives.** In the case of claims that products kill and/or knockdown bed bugs,
 128 the objective of bed bug product performance testing is to determine that a
 129 product application made at the proposed label claim kills or knocks down the
 130 bed bugs. For claims that products attract or repel bed bugs, the objective of
 131 product performance testing is to determine the ability of a product to
 132 encourage or deter bed bugs to or from a pre-determined locale. For “control
 133 bed bugs” claims, the objective is to determine that the insecticide application
 134 shows a sufficient efficacy (e.g. mortality and/or knockdown/morbidity
 135 followed by mortality) and/or residual efficacy. In all cases the scientific
 136 objective should be stated clearly, and all treated bed bugs should be compared
 137 to control bed bugs that have received no treatment (5, 12, 18, 21, 29).

138 **Test materials and treatments.** Product performance should be tested using
 139 the end-use formulation and application rates as registered or as proposed for
 140 use. Test materials should be stored at ambient temperature and humidity for at
 141 least one day before use.

142 b. **Product treatments for product performance tests:**

143 i. **Products that target bed bug nymphs and adults.** Testing of products
 144 that target bed bug nymphs and adults should be conducted with both, bed
 145 bug nymphs and adults. Preferably, when nymphs are tested, the last two
 146 nymphal stages (J4 or J5) should be used.

147 ii. **Products that target bed bug development.** Testing of products that
 148 target bed bug development should be conducted with mixed nymphal
 149 stages or eggs as appropriate. A blood meal should be available to provide
 150 the nourishment needed for bed bugs to moult from one nymphal stage to
 151 the next. The blood meal should be identified.

152 iii. **Products that target bed bug eggs (ovicidal products).** Testing of
 153 products that target bed bug eggs should be conducted with the egg life
 154 stage only.

155 c. **Dose determination.** The test dose in bed bug product performance studies is
 156 the lowest application rate from a proposed product label. The dose will differ
 157 according to the surface type (sorptive/non-sorptive). The rate should be
 158 reported using metric system measurements, as mg or ml of test chemical/cm²
 159 for surface area treatments or volumetrically as ml of test chemical/m³ for space

160 spray, total release aerosols, and fumigant treatments. The amount of active
161 ingredient tested per unit area or volume should also be given.

162 d. **Testing conditions.** During bed bug product performance testing the
163 temperature should be kept at $22 \pm 4^{\circ}\text{C}$ (unless otherwise justified, e. g. realistic
164 use in cold conditions), with a relative humidity of 40-60%, and a photoperiod
165 ranging from L:D 12:12 h to L:D 16:8 h. The temperature during the test should
166 be kept as constant as possible because changes can affect the performance of
167 the product treatments.

168 e. **Choice of endpoints and measures.** Endpoints chosen for the study should be
169 appropriate for the specific objectives of proposed research and likely to
170 provide a robust answer to the research question. Generally, the endpoints
171 tested will be bed bug knockdown, morbidity, kill, repellency, or attraction
172 tested at the claimed labelled application rate and will determine whether or not
173 a product is efficacious. The selected endpoint should be included into the
174 protocol. Bed bugs should be removed and placed into a clean container within
175 a maximum of 4 hours or, in accordance with the claim, after onset of exposure
176 to the biocide application.

177 The exception in the product evaluation is the simulated-use test with a test
178 duration of 24 h (Section 4.3) and the simulated-use test for repellents with a
179 test duration of at least 8 h (Section 4.4.).

180 Besides product evaluation, laboratory studies can be done for resistance ratio
181 determination and characterization of bed bug strain susceptibility where bed
182 bugs should be exposed to the treated filter paper in each treatment for 24 hours
183 (Section 4.1.).

184 For knockdown evaluation, observations should be made within a few minutes
185 post exposure. For mortality evaluation, observations should be reported at 24,
186 48, 72, and 96 hours up to 8 days post-exposure (or according to the claim)
187 unless all bed bugs die or negative control mortality exceeds 10% (18). Data
188 will not be acceptable if control mortality exceeds 10%. Observations of
189 mortality occurring after 96 hours should be justified based on the mode of
190 action and application type. Survival of bed bugs beyond 8 days in negative
191 control replicates does not justify making observations after 8 days. The
192 number of dead, knocked down, moribund and live bed bugs in each replicate
193 should be recorded. A count on mortality should be separate from a count on
194 knockdown and morbidity. The percentage of bed bugs killed, moribund and
195 knocked down, exclusively, for each treatment at each test interval should be
196 recorded. Confidence limits around the reported mean or median values should
197 be reported. Evaluation of speed of kill or additional knockdown evaluations
198 should include more observations on the first day of the test (e.g. observations
199 2, 4 and 6 hours post exposure) to record the data needed to support the desired
200 claims. If a test recommends that a bed bug should be contained during
201 exposure to an insecticide, a bed bug should not be confined to a treated surface
202 for more than 4 hours or in accordance with the claim. For the evaluation of
203 products containing attractants or repellents, the endpoint is the minimum
204 efficacy on the label claim, e.g. percentage of repelled or attracted bed bugs in
205 a given time, respectively.

206 f. **Test organisms.** Testing should be conducted with a demonstrably susceptible
207 laboratory strain of the common bed bug, *Cimex lectularius*. If *Cimex*
208 *hemipterus* is claimed, this species should be tested. Tests should be conducted

209 with both, male and female bed bugs, in a balanced sex ratio. If strains collected
210 in the field are tested, their susceptibility to the given insecticide has to be
211 evidenced by bioassays and/or molecular methods before testing. Testing
212 should occur no later than the second lab-reared generation. If more generations
213 are needed to produce sufficient numbers of bed bugs for testing, it should be
214 indicated in the study report. Bed bugs should be blood fed and should be tested
215 preferably seven days after the last blood meal. Deviations of five to ten days
216 after the last blood meal are possible.

- 217 g. **Representative sampling.**
- 218 i. **Sample size.** The sample should be large enough (with a minimum of 50
219 bed bugs) to likely yield a definitive answer to the research question being
220 addressed, and its size should be justified statistically, taking into account
221 the specific characteristics of the proposed research and the necessary
222 accuracy and precision of the results.
- 223 ii. **Replication.** A minimum of five replicates of ten bed bugs each and
224 balanced (equal number of treated and control replicates) experimental
225 designs are recommended for most studies. Exceptions will be noted in the
226 guidance that follows in this document.

227 Other factors that may affect sample size and replication are the number of
228 treatments, the experimental design, and the heterogeneity of the sample bed
229 bug population (e.g. developmental stage, sex, insecticide susceptibility) and
230 the environment (different habitat population densities). The protocol should
231 fully describe how sample size and replication were determined.

- 232 h. **Bed bug rearing, handling, and maintenance.** When applicable, a description
233 of the bed bug laboratory colony rearing practices should be included.
234 Collection details and maintenance procedures for field-collected strains should
235 be described. Bed bug feeding can be conducted on animal as well as on human
236 hosts or alternatively artificially. Although bed bugs can be fed on small
237 mammals like rabbits, for animal welfare reasons *in vitro* feeding methods
238 should be preferred (1, 28). Different membrane feeding systems can be used
239 to offer the blood meal (1, 28). In addition, a Hemotek® system can be used. In
240 this case, a flat blood reservoir should be preferably used, in which erythrocytes
241 do not sink to the bottom of the feeder where bed bugs cannot reach them.
242 Different membranes like double stretched parafilm, collagen or silicone can
243 be used. Before feeding, bed bugs should be placed in a jar closed with gauze
244 or mesh screening which bed bugs can pierce but cannot escape.

- 245 i. **Negative control.** A negative control should be included in all testing. The
246 number of negative control replicates should equal the number of replicates for
247 each treatment. When appropriate, a negative control is typically treated with
248 diluent only or receives no treatment at all.

- 249 j. **Positive controls.** A positive control is not necessary. An appropriate positive
250 control, if available, is recommended for determining a resistance ratio.

- 251 k. **Statistical analysis plan.** Protocols should include a full description,
252 explanation, and justification for the statistical methods proposed to analyse
253 both resistance ratio determinations and product performance test results,
254 taking into account the specific study objectives and variables. A statistician
255 should be consulted when developing test protocols. Protocols should explicitly
256 describe the model to be used and demonstrate whether or not assumptions

257 underlying the model can be met for all proposed analyses. Restrictions on
 258 randomisation of any testing components should be documented clearly and
 259 should be accounted for correctly in the statistical analyses. Generally,
 260 generalized linear models (GLMs) (25) are recommended to fit models directly
 261 to non-normal (e.g., binomial – which describe much of the collected product
 262 performance data sets) data using a probit link or logit link function. GLMs do
 263 not involve transforming the response variable, thereby allowing the data to
 264 remain on the original scale of measurement. Generalized linear mixed-models
 265 (GLMM) (13, 14, 19, 21) may also be appropriate. Software for analysis using
 266 GLMs or GLMMs is available in many widely sold statistical analysis
 267 packages. If survival analyses (27), such as the Kaplan- Meier Estimator, are
 268 used, provide justification for use of the median value to characterise product
 269 performance and demonstrate that the underlying assumptions of these analyses
 270 have been met. Other analysis including assumption of the normal distribution
 271 should be described and justified (2, 9, 37, 40, 44).

272 1. **Quality assurance /Quality control (QA/QC) plan.** Protocols should provide
 273 for periodic quality assurance inspections that are adequate to ensure the
 274 integrity of the study and consistency with the provisions of OECD Principles
 275 of Good Laboratory Practice (GLP) and Compliance Monitoring
 276 ([ENV/MC/CHEM\(98\)17](#)).

277 m. **Quality Assurance (QA) oversight.** Product performance testing is subject to
 278 the OECD Principles of Good Laboratory Practice and Compliance Monitoring
 279 (GLP) ([ENV/MC/CHEM\(98\)17](#)). The GLP states that each testing facility
 280 should include an independent Quality Assurance (QA) unit and that the QA
 281 unit monitors execution of each protocol and documents its conduct in
 282 accordance with the GLP ([ENV/MC/CHEM\(98\)17](#)). The QA unit will inspect
 283 each study at intervals adequate to ensure the integrity of the study and maintain
 284 written and properly signed records of each periodic inspection.

285 n. **Protocol amendments.** Amendments are planned changes to the protocol and
 286 should be made before the study is executed. All amendments to the protocol
 287 should be noted in the written report.

288 o. **Deviations from protocol.** Even when executing the best-designed and most
 289 comprehensive protocols, unanticipated deviations from the protocol may
 290 occur. All such deviations from the protocol and their impact on the research
 291 should be fully reported in the study report.

292 3.2. Data collection and reporting

293 Study protocols should provide for collection and reporting of data covering all aspects of
 294 the research including the following elements:

295 a. **Study identification:** Title, identifying study number(s), sponsor, study
 296 director, investigators, name and location of the testing facility, and dates of the
 297 study should be reported.

298 b. **Approved or proposed label directions for use:** A copy of the proposed or
 299 approved product label should be included.

300 c. **Study objective(s):** The purpose of the study should be stated.

301 d. **Testing conditions:** Information on temperature, relative humidity, ambient
 302 light and photoperiod, and air flow (where applicable) should be reported.

- 303 e. **Testing system, including but not limited to:**
- 304 • Bed bug species tested, including identification of strains of susceptible and
305 field bed bug populations, where bed bug strains were collected/obtained;
306 development stage, age, and sex of bed bugs; and methods for preparation
307 of bed bugs for test (feeding/starving), and when appropriate, the blood
308 meal should be identified (20, 35, 42).
- 309 • Bed bug rearing, handling, and maintenance.
- 310 • Description of test chemical (i.e. product, % active ingredient, and
311 formulation to be tested). Negative control should also be described.
- 312 • Description of the experimental unit.
- 313 • Treatment application rate and method of application (rate should be
314 consistent with label instructions).
- 315 • Number of product treatments.
- 316 • Number of negative control replicates.
- 317 • Number of positive control replicates (where applicable).
- 318 • Number of replicates per treatment.
- 319 • Number of bed bugs per replicate for each treatment.
- 320 • Length of time for bed bug exposure period to each treatment.
- 321
- 322 f. **Data/Results reporting.** Report the following information:
- 323 i. **Protocol with amendments and study deviations from the protocol.** A
324 copy of the study protocol should be included with amendments and
325 deviations. Amendments and deviations should be justified, and described
326 together with their impact on the validity of the study.
- 327 ii. **Data and endpoints.** Knockdown, morbidity and mortality values should
328 be corrected for negative control knockdown, morbidity and/or mortality
329 with Abbott's Formula or the equivalent. Endpoints should be reported as
330 observed throughout the test, though total and percent knockdown,
331 morbidity and mortality should be reported at the final evaluation.
- 332 iii. **Amount of product applied.** The product rate should be reported using
333 metric system measurements, as mg or ml of test chemical/cm² for surface
334 area treatments or volumetrically as ml of test chemical/m³ for space spray,
335 total release aerosols, and fumigant treatments for each replicate. The
336 amount of active ingredient tested per unit area or volume should also be
337 given.
- 338 iv. **Report the following other data:**
- 339 • Test results on all aspects of the research.
- 340 • Copies of all raw data.
- 341 • Certification of the test chemical's identity and origin.
- 342 • Description of each product treatment and the negative controls.

- 343
- 344
- Note: when (national) guidelines are used, those should be referred to in the report.
- 345
- 346
- 347
- 348
- 349
- 350
- 351
- 352
- 353
- 354
- 355
- v. **Data analysis.** The report should include the statistical analysis plan. Refer to Section 3.1.1 for recommendations on data analyses.
- vi. **Study Conclusions.** The report should include a discussion of the study results and conclusions based on treatment endpoints. Conclusions should state why and how the study results do or do not support the tested hypothesis.
- vii. **Storage and Retention of Records and Materials.** The record-keeping provisions of OECD Principles of Good Laboratory Practice and Compliance Monitoring (GLP) ([ENV/MC/CHEM\(98\)17](#)) apply to records of any study conducted under the Good Laboratory Practices rule, in compliance with any applicable state laws or regulations.

356

4. Specific Guidance

4.1. Specific guidance for laboratory studies for resistance ratio determination and characterization of bed bug strain susceptibility.

359 Response ratio determination, commonly known as a “Resistance Ratio,” is useful in
360 characterising the magnitude of susceptibility or resistance to insecticides used in bed bug
361 control. Different bioassays are suitable for resistance ratio determination of bed bug strain
362 susceptibility (e.g. 39). This guidance describes the filter contact bioassays designed to
363 determine the resistance ratio (37, 46).

4.1.1. Study objective.

364 To estimate the susceptibility and magnitude of resistance of bed bug strains to insecticides
365 used in product testing.
366

4.1.2. Materials and methods

Filter contact bioassay in petri dishes

369 a) **Experimental units.** Place a piece of white filter paper on the bottom of a 6 or 10
370 cm glass petri dish and secure a screen over the top of the petri dish. An insecticide
371 concentration should be applied to filter paper in each replicate at a volume that
372 saturates the paper, generally at least 200 µl. Allow paper to dry before exposing
373 the bed bugs. Prepare an equal number of negative control dishes with paper treated
374 with the diluent only.

375 b) **Number of treatments.** Five concentrations of the active ingredient should be
376 prepared with the appropriate diluent. Active ingredient concentrations should be
377 prepared based on a logarithmic scale, i.e. 0.0001%, 0.001%, 0.01%, 0.1%, and
378 1.0%. Other concentrations may be used based on previous knowledge of bed bug
379 susceptibility to the insecticide being tested but a justification should be provided
380 (3, 7, 8, 17, 19, 31, 36, 43, 51, 53). If a product contains a synergist, use only the
381 insecticide component with a solution concentration based on the active ingredient,
382 not the synergist (33).

383 c) **Number of replicates.** Ten replicates per concentration with ten bed bugs each are
384 recommended. It is recommended to test male bed bugs only, because results on
385 females with sublethal concentrations are often more variable due to traumatic
386 insemination.

Filter contact bioassay in 24-well cell culture plates.

387 Since bed bugs show strong aggregation behaviour, it is often practical to place test
388 individuals in separate chambers.
389

390 a) **Experimental units.** Place pieces (ø 1.6 cm) of white filter paper on the bottom of
391 24-well cell culture plates. An insecticide concentration should be applied to filter
392 papers in each replicate at a volume that saturates the paper, generally at least 50
393 µl. Allow paper to dry completely (preferably in a laboratory fume hood) before
394 exposing the bed bugs.

- 395 b) **Number of treatments.** Five concentrations of the active ingredient should be
 396 prepared with the appropriate diluent. Active ingredient concentrations should be
 397 prepared based on a logarithmic scale, i.e. 0.0001%, 0.001%, 0.01%, 0.1%, and
 398 1.0%. Other concentrations may be used based on previous knowledge of bed bug
 399 susceptibility to the insecticide being tested but a justification should be provided
 400 (3, 7, 8, 17, 19, 31, 36, 43, 51, 53). If a product contains a synergist, use only the
 401 insecticide component with a solution concentration based on the active ingredient,
 402 not the synergist (33).
- 403 c) **Number of replicates.** Five replicates of separate 24-well plates per concentration
 404 should be conducted. On each of the five separate plates it is recommended to place
 405 18 individual bed bugs on treated filter papers and 6 bed bugs as negative controls
 406 (number of bed bugs tested per concentration: n=90 treated bed bugs and n=30 as
 407 negative controls, number of bed bugs tested in total: n=450 treated bed bugs and
 408 n=150 as negative controls. It is recommended to test male bed bugs only, because
 409 results on females with sublethal concentrations are often more variable due to
 410 traumatic insemination.

411 *The following specifications find application with all bioassays.*

- 412 a) **Bed bug exposure to treatments.** Bed bugs should be exposed to the treated filter
 413 paper in each treatment for 24 hours.
- 414 b) **Negative control.** The negative control should be treated with the diluent for
 415 insecticide solution preparation.
- 416 c) **Positive control.** An appropriate positive control, if available, should be used.
- 417 d) **Lethal dose (LD) or effective concentration (EC) values.** An analysis using
 418 GLMs is recommended to determine the LD or EC values for each bed bug strain
 419 tested. Use of a logit or probit analysis should be justified (45).
- 420 e) **Resistance ratios (RR).** Resistance ratios should be calculated and reported as
 421 follows:
- 422 $LD \text{ or } EC \text{ (lab or field strain)} / LD \text{ or } EC \text{ (susceptible strain)} = RR$
- 423 For example: $LD_{50} \text{ (lab or field strain)} / LD_{50} \text{ (susceptible strain)} = RR_{50}$
- 424 $LD_{90} \text{ (lab or field strain)} / LD_{90} \text{ (susceptible strain)} = RR_{90}$

425 **4.1.3. Reporting results.**

426 Refer to Section 3.2.f. of this guidance for guidance in reporting results and the data
 427 analysis. Report the resistance ratio values for each strain for each insecticide tested and
 428 the associated data analysis (30).

429 **4.1.4. Efficacy evaluation.**

430 Immediately after the 24 h exposure period efficacy evaluation should be performed. If
 431 further observations are necessary, transfer bed bugs to clean, untreated petri dishes or 24-
 432 well cell culture plates and evaluation observations should be reported as described under
 433 Section 3.1.e. A resistance ratio equal to or greater than 100 is characteristic of a resistant
 434 strain. It has to be acknowledged, that practical bed bug control may fail even in bed bug
 435 populations with a resistance ratio < 100.

436 **4.1.5. Study conclusions.**

437 Summarise and discuss the study outcome.

438

439 **4.2. Specific guidance for laboratory studies for forced exposure (no-choice)**
440 **residual surface treatments.**441 Different bioassays are suitable for determination of the residual product performance with
442 a forced exposure (no-choice) test design (e.g. 18).443 **4.2.1. Study objective.**444 To determine the residual product performance of an application made to different sorptive
445 and non-sorptive surfaces in a forced exposure (no-choice) test (18).446 **4.2.2. Materials and methods**447 a) **Experimental units.** The surfaces to be treated with the product should have
448 sorptive and non-sorptive properties that are representative of where bed bugs are
449 found. For example: unpainted/unfinished plywood; wallpaper, 100% cotton
450 sheeting stretched over a cardboard panel, commercial linoleum tile; laminate or
451 tiles. Surfaces should be pre-cut to 10 x 10 cm or larger panels (18). Cotton sheet
452 replicates should be affixed to the top of the panel to provide a flat, rigid surface
453 for treatment. An application should be made to each surface. An equal number of
454 negative control replicates should be established with the same surfaces. Petri
455 dishes can be used as container for the surfaces. A glass ring on the surfaces can
456 prevent bed bugs from escaping.457 b) **Application of product dilutions.** The claimed application rate for bed bug
458 control should be applied on each panel. A metered bench top sprayer is preferred
459 as the delivery device to ensure consistent application volume and even distribution
460 of spray particles. Generally, panels are sprayed from a distance of 30 cm above
461 the panel surface. Use of other heights should be justified. Panels should be stored
462 and exposed to ambient conditions at the test site to age residues. Panels should be
463 fully dried before exposing bed bugs. Measure the volume of spray applied and
464 calculate the weight of the active ingredient(s) delivered. The quantity of product
465 per square meter should be determined.466 c) **Ready-to-use product application.** Application to panel surfaces should be made
467 at the rate equivalent to the amount of product to be sprayed per unit area as
468 directed by the label. Generally, panels are sprayed from a distance of 30 cm above
469 the panel surface. Use of other heights should be justified. In product performance
470 testing, the amount of product delivered by a ready-to-use spray product (aerosol
471 or pump-spray) is described as the amount of product sprayed per second or as
472 number of pumps per unit area and should be determined before treatments can be
473 made. To determine the quantity sprayed per second, spray five panels of each
474 surface type for three seconds each for each treatment. The product container
475 should be weighed before and after each spray and the difference recorded. The
476 mean value of the five replicates should be determined and that result divided by
477 three to determine the average amount of product applied per second of spraying.
478 The same procedure should be conducted to evaluate dust product formulations
479 except that application should be made from a height of 15 cm or as directed by

480 the product label. Other modifications to the protocol needed to apply dusts should
481 be described and justified.

482 d) **Replication.** A replicate should consist of a treated or untreated panel, each with
483 10 to 12 bed bugs per each life stage confined to the panel. A minimum of 5
484 replicates should be used for each product treatment and negative control. For each
485 bed bug strain tested at each exposure time, prepare 15 treated (3 surfaces x 5
486 replicates per surface) and 15 untreated panels. Three hundred to 360 bed bugs are
487 necessary for each exposure time tested per strain (including the negative control
488 specimens).

489 e) **Bed bug exposure to product treatments.** Bed bugs should be exposed to treated
490 panels for no more than 4 hours or in accordance with the claim. After the exposure
491 period, transfer the bed bugs to a clean, untreated container for further observation
492 and evaluation. After an initial test 24 hours post application, the insecticide
493 residues should be tested regularly until the end of residual period claimed. Treated
494 panels should be retested though bed bugs should not be reused.

495 f) **Positive control.** A positive control is not necessary.

496 **4.2.3. Reporting results.**

497 Refer to Section 3.2.f. of this guidance for guidance on reporting results and data analysis.

498 **4.2.4. Efficacy evaluation.**

499 Efficacy is usually considered sufficient if – at the end of the test, in each life stage tested,
500 – 100% mortality rate, is achieved.

501 **4.2.5. Study conclusions.**

502 Summarize and discuss study outcomes for residual control of bed bugs.

503

504 **4.3. Specific guidance for laboratory studies to determine the product performance** 505 **of insecticide products under practical use situation.**

506 Different bioassays are suitable for determination of the residual product performance with
507 a simulated-use test design. This guidance describes the efficacy evaluation of residual
508 surface treatments in a test system designed to mimic a practical use situation (46).

509 **4.3.1. Study objective.**

510 To determine the residual product performance of an application made to different sorptive
511 and non-sorptive surfaces in a simulated-use test.

512 **4.3.2. Materials and methods**

513 a) **Experimental units.** The test arena should consist of three closed chambers joined
514 with round or rectangular connectors. In the first chamber, a sealed harborage
515 should be provided (harborage chamber). The design of the harborage should
516 enable the bed bugs to leave easily, e.g. a pocket made of towel paper and tape
517 opened with scissors. After a minimum of 1 h of acclimatisation, the harborage
518 should be opened. In the chamber connected to the harborage chamber, the treated
519 surface or textile is placed (insecticide chamber). The insecticide chamber is

520 connected to a third chamber containing a carbon dioxide (CO₂) source and a heat
521 source (host chamber). The end of the last connector should protrude
522 approximately 10 cm into the host chamber. A collecting vessel should be placed
523 under the open end of the connecting tube. The collecting vessel should contain
524 filter paper as a harborage. Bed bugs which have crossed the surface then fall into
525 the vessel, which prevents bed bugs from escaping the test arena. Bed bugs should
526 be removed 24 h post exposure or according to the claim for determination of
527 efficacy. Bed bugs that have stayed in the towel paper pocket are excluded from
528 evaluation.

529 Distance from the harborage to host chamber should be between 50 cm and 80 cm
530 and the amount of CO₂ should be then adjusted to 0.75 l/min. If deviating distances
531 are used, the amount of CO₂ has to be adjusted. A suction pump reaching in at the
532 top of the harborage chamber prevents an intoxication of the bed bugs due to CO₂.
533 It should be used a CO₂ flow rate that guarantees a minimum of 80% bed bugs
534 leaving the harborage. Inside the host chamber, temperature should be adjusted to
535 37°C ± 2°C. Connectors between the three chambers should be about 15 cm to 20
536 cm. Connectors should be lined with material (e.g. masking tape or paper) which
537 is not slippery for bed bugs. Inner walls of the collection vessel should be treated
538 with a substance which prevents bed bugs from escaping. The surfaces to be treated
539 with the product should have sorptive and non-sorptive properties:
540 unpainted/unfinished plywood; wallpaper, 100% cotton sheeting stretched over a
541 cardboard panel, commercial linoleum tile; laminate or tiles. Surfaces should be
542 pre-cut to 10 x 10 cm or larger panels (18). Cotton sheet replicates should be
543 affixed to the top of the panel to provide a flat, rigid surface for treatment. An
544 application should be made to each surface. An equal number of negative control
545 replicates should be established with the same surfaces.

546 b) **Application of product dilutions.** The claimed application rate for bed bug
547 control should be applied on each panel. A metered bench top sprayer is preferred
548 as the delivery device to ensure consistent application volume and even distribution
549 of spray particles. Generally, panels are sprayed from a distance of 30 cm above
550 the panel surface. Use of other heights should be justified. Panels should be stored
551 and exposed to ambient conditions at the test site to age residues. Panels should be
552 fully dried before exposing bed bugs. Measure the volume of spray applied and
553 calculate the weight of the active ingredient(s) delivered.

554 c) **Ready-to-use product application.** Application to panel surfaces should be made
555 at rates equivalent to the amount of product to be sprayed per unit area as directed
556 by the label. Generally, panels are sprayed from a distance of 30 cm above the
557 panel surface. Use of other heights should be justified. In product performance
558 testing, the amount of product delivered by a ready-to-use spray product (aerosol
559 or pump-spray) is described as the amount of product sprayed per second or per
560 number of pumps per unit area and should be determined before treatments can be
561 made. To determine the quantity sprayed per second, spray five panels of each
562 surface type for three seconds each for each treatment. The product container
563 should be weighed before and after each spray and the difference recorded. The
564 mean value of the five replicates should be determined and that result divided by
565 three to determine the average amount of product applied per second of spraying.
566 The same procedure should be conducted to evaluate dust product formulations
567 except that application should be made from a height of 15 cm or as directed by
568 the product label. Other modifications to the protocol needed to apply dusts should
569 be described and justified.

- 570 d) **Replication.** Tests should be conducted with 50 to 100 bed bugs (equal sex ratio
571 and per each life stage) in minimum five treated and negative control replicates per
572 surface type.
- 573 e) **Positive control.** A positive control is not necessary.
- 574 f) **Additional testing conditions.** Within the exposure period, darkness or red light
575 is obligatory.

576 **4.3.3. Reporting results.**

577 Refer to Section 3.2.f. of this guidance for guidance on reporting results and data analysis.

578 **4.3.4. Efficacy evaluation.**

579 Efficacy is usually considered sufficient if – in each life stage tested – 100% mortality rate
580 after 24 h exposure and until the end of residual period claimed, corrected according to
581 Abbott is achieved.

582 **4.3.5. Study conclusions.**

583 Summarize and discuss study outcomes for residual control of bed bugs.
584

585 **4.4. Specific guidance for laboratory studies to determine the product performance** 586 **of a repellent or attractant product.**

587 Different bioassays are suitable to determine the efficacy of a repellent or an attractant (11,
588 24, 38, 41, 49, 50). The following guidance describes approaches that may be used to assess
589 whether a product is a repellent or attractant.

590 **4.4.1. Study objective.**

591 To determine if a biocide product repels or attracts bed bugs.

592 **4.4.2. Materials and methods**

593 *Repellents.*

594 Simulated-use tests for volatile repellents and impregnated fabric.

- 595 a) The test arena described by (46) should consist of three closed chambers joined
596 with round or rectangular connectors. In the first chamber, a sealed harborage
597 should be provided (harborage chamber). The design of the harborage should
598 enable the bed bugs to leave easily e.g. a pocket made of towel paper and tape
599 opened with scissors. After a minimum of 1 h of acclimatisation the harborage
600 should be opened. In the chamber connected to the harborage chamber, the treated
601 surface or textile is placed (treated chamber). The treated chamber is connected to
602 a third chamber containing a CO₂ source and a heat source (host chamber). The
603 end of the last connector should protrude approximately 10 cm into the host
604 chamber. A collecting vessel should be placed under the open end of the connecting
605 tube. The collecting vessel should contain filter paper as a harborage. Bed bugs
606 which might have crossed the surface then fall into the vessel, which prevents bed
607 bugs from escaping the test arena. Bed bugs should be removed 8 h post exposure
608 or according to the claim for determination of efficacy. Bed bugs that have stayed

609 in the towel paper pocket are excluded from evaluation. Bed bugs that have stayed
610 in the first chamber outside the towel paper pocket are considered as “repelled”,
611 while specimen that are found on the treated surface or have fallen into the vessel
612 are considered as “not repelled”. Non-insecticidal efficacy should be demonstrated
613 with bed bugs that have been in contact with the freshly applied product. Mortality
614 of the insects should be monitored at the end of the test until 8 days post exposure.

615 Distance from the harborage to host chamber should be between 50 cm and 80 cm
616 and the amount of CO₂ should be then adjusted to 0.75 l/min. If deviating distances
617 are used, the amount of CO₂ has to be adjusted. A suction pump reaching in at the
618 top of the harborage chamber prevents an intoxication of the bed bugs due to CO₂.
619 It should be used a CO₂ flow rate that guarantees a minimum of 80% bed bugs
620 leaving the harbourage. Inside the host chamber, temperature should be adjusted
621 to 37°C ± 2°C. Connectors between the three chambers should be about 15 cm to
622 20 cm. Size of the treated surface should be no larger than 20 cm. Connectors
623 should be lined with material (e.g. masking tape or paper) which is not slippery for
624 bed bugs. Inner walls of the collection vessel should be treated with a substance
625 which prevents bed bugs from escaping.

626 b) The test arena described by (47) consists of a plastic tray (80 by 75 by 5 cm) and a
627 small bed or imitations with four legs (e.g. a stool or small table) placed in the
628 center. Onto the simulated bed, a CO₂ and additionally a heat source should be
629 placed to mimic a human host. Under each leg a bed bug interceptor (i.e. a double-
630 walled bed bug trap, where the insects are being trapped between both walls, which
631 form a ring around the bed leg) should be placed as collection vessel. The repellent
632 product should be applied on the outer wall of the interceptor according to the label
633 claim. If the test surface should be larger, the interceptors should be placed onto
634 the treated surfaces. A minimum of 100 ml/min of CO₂ should be released on the
635 top of the bed. A harborage should be placed in the center of the plastic tray right
636 under the bed. For acclimatisation, the harborage has to be closed. After a
637 minimum of 1 h of acclimatisation, the harborage should be opened. The design of
638 the harborage should enable the bed bugs to leave easily, e. g. a pocket made of
639 towel paper and tape opened with scissors. Test arena should be lined with material
640 (e.g. paper and masking tape) which enables normal bed bug movements. Inner
641 walls of the tray should be treated with a substance which prevents bed bugs from
642 escaping.

643 c) **Replication.** Tests should be conducted with 50 to 100 bed bugs (equal sex ratio)
644 in minimum five treated and negative control replicates. The exposure period
645 should be according to the label claim, but at least 8 h (to cover the natural bed bug
646 activity time over night).

647 d) **Positive control.** A positive control is not necessary.

648 e) **Additional testing conditions.** Testing under darkness or red light is obligatory.

649 Laboratory choice tests for volatile repellents.

650 a) A petri dish assay with a treated and untreated filter paper similar to the test setup
651 of (47) may be considered.

652 b) Also a version of the still-air olfactometer (10) as modified by (48) may be
653 considered. The still-air model should be adapted to provide a source such as CO₂
654 and heat to mimic a human host in the presence and absence of repellent as
655 alternative choices in the same arena. The negative control should consist of the

656 same arrangement in a separate arena provided with only CO₂ and heat and with
657 no repellent.

658 c) A Y-tube assay (15) may also be considered.

659 d) **Replication.** Ten bed bugs per replicate for a total of five replicates per trial and
660 negative control or 50 replications of one bed bug each per trial and negative
661 control should be used. Justify the choice of individuals or groups of ten.

662 e) **Positive control.** A positive control is not necessary.

663 f) **Additional testing conditions.** Testing under red light is recommended.

664 Laboratory choice tests for impregnated fabric.

665 Untreated and impregnated fabric should be tested together. Bed bugs should be placed
666 onto the untreated fabric, and their movement both towards or away from the treated fabric
667 should be observed and measured. A negative control arena with untreated fabric only
668 should also be included in the study (e.g. 11, 16, 23).

669 *Attractants.*

670 Choice tests are recommended for testing the product performance of attractant products
671 (48, 49). Depending on its use pattern (whether the product is intended to attract bed bugs
672 towards a harborage or “pull” them away from human hosts), the product performance of
673 an attractant product may be determined by comparing its effect on bed bugs to that of bed
674 bug aggregation cues or host cues such as CO₂ and heat. Therefore, in this experiment either
675 host cues such as CO₂ and heat or bed bug aggregation cues deposited in a harborage should
676 be presented as an alternative to the attractant formulation in the same arena. Observations
677 of bed bug location should be recorded at the end of the exposure period. The exposure
678 period should be according to the label claim, but at least 8 hours.

679 a) **Replication.** Ten bed bugs per replicate for a total of five replicates per trial and
680 negative control or 50 replications of one bed bug each per trial and negative
681 control should be used. Justify the choice of individuals or groups of ten.

682 b) **Positive control.** A positive control is not necessary.

683 c) **Additional testing condition.** Testing in the dark under red light is recommended
684 (49).

685 **4.4.3. Reporting results.**

686 Refer to Section 3.2.f. of this guidance for guidance on reporting results and data analysis.
687 Analysing data from replicates with one bed bug or replicates with ten bed bugs is likely to
688 differ and should be justified.

689 a) **Percent repellency.** For repellent testing, report the number of bed bugs that
690 avoided the host mimic and the ones that did not. Calculate percent repellency
691 corrected for negative control results. The exposure period should be according to
692 the label claim, but at least 8 h.

693 b) **Percent attractancy.** For attractant testing, report the number of bed bugs that
694 were attracted to the attractant and the number that were not. Calculate the
695 percentage of bed bugs found at each location at 15 minutes post exposure and
696 until the end of the claimed period in treated choice and negative control arenas.

697 c) **Mortality, morbidity and knockdown.** Non-insecticidal efficacy should be
698 demonstrated with bed bugs that have been in contact with the freshly applied

699 product. Mortality of the insects should be monitored at the end of the test until 8
700 days post exposure. All raw data should be reported.

701 **4.4.4. Efficacy evaluation.**

702 a) **Repellents.** For the claim “prevents bed bug bites” or “prevents the spreading of
703 bed bugs” 100 % repellency are required in each life stage tested.

704 The exposure period should be according to the label claim, but at least 8 h (to
705 cover the natural bed bug activity time over night).

706 b) **Attractants.** The efficacy of an attractant is usually considered sufficient when at
707 least 80% of the test individuals were attracted compared to the negative control
708 within the test period or according to the claim, from the beginning and until the
709 end of the claimed efficacy period.

710 The percentage of bed bugs found at each location at 15 minutes post exposure and
711 until the end of the claimed period has to be reported in treated choice and negative
712 control arenas.

713 **4.4.5. Study conclusions.**

714 Describe the product performance of the product treatment. Include a discussion on
715 negative control results, and the adequacy of the host or aggregation cues used in the study.

716

717 **4.5. Specific guidance for laboratory studies for testing indoor insecticide total** 718 **release aerosols, space sprays, and insecticide vapour strip products.**

719 This guidance applies to testing total release aerosols and space sprays including misters,
720 hand-held aerosol products and vaporizing strips on surfaces used for indoor applications
721 to control bed bugs. Different bioassays are suitable to determine the efficacy these
722 treatments.

723 **4.5.1. Study objective.**

724 To determine the performance of insecticide products intended for total release aerosols,
725 space sprays, and vapour strip treatments against bed bugs.

726 **4.5.2. Materials and methods**

727 a) **Experimental unit.** Testing should be conducted in a Peet-Grady chamber with a
728 volume of 6.12 cubic meters (1.83 x 1.83 x 1.83 m) or greater (52). The chamber
729 should have a window for observation. The wall, ceiling, and floor of the
730 room/chamber may be lined with plastic or other suitable materials to facilitate
731 cleaning. Test doses for aerosols and mister products should be delivered by an
732 automatic dispenser calibrated for the proper droplet size and application rate. At
733 the end of each replicate, the air in the chamber should be exhausted and any
734 surface residues washed off. Surfaces should be clean and dry before the next test.
735 Alternative product application methods may be considered, but should be
736 described and justified. Vaporizing strips should be hung from the ceiling in the
737 center of the room/chamber or applied according to label directions.

738 b) **Replication.** Tests should be conducted with in minimum five treated and five
739 negative control replicates per each exposure period. One half of the replicates will

740 be in the chamber treated with the insecticide product while the other half will be
741 the negative control.

742 c) **Cage Placement.** For each exposure period, twelve cages from each strain should
743 be used. Appropriate shelters, such as stacked egg cartons should be added to each
744 cage to serve as a harborage. Allot six cages to the product treatment and place in
745 the chamber, while the other six should be kept outside the chamber as a negative
746 control for each exposure period. Ten bed bugs (equal sex ratio) should be
747 transferred to each cage. In the chamber use balanced replicates representing three
748 heights (two at floor level, two at mid wall and two at ceiling level).

749 d) **Bed bug exposure to the treatments.** The chamber should be sealed and the
750 product application made. After the application is made, the test cages should be
751 left in place for two hours or according to the claim and removed after the
752 insecticide has been evacuated from the chamber. Knockdown, morbidity and
753 mortality should be recorded at two hours after application or according to the
754 claim. Bed bugs from each cage should be transferred to a clean, untreated
755 container after the assessments are made, but no later than 4 hours post treatment
756 or in accordance with the claim. Vaporizing strips should be assessed in a similar
757 manner following 24 hours of exposure.

758 e) **Negative control.** Negative control replicates should be placed in the lab under the
759 same abiotic conditions as the treatments. Negative control replicates should be
760 untreated because treating with diluent is impractical.

761 f) **Positive control.** A positive control is not necessary.

762 **4.5.3. Reporting results.**

763 Refer to Section 3.2.f of this guidance for guidance on reporting results and data analysis.
764 In addition, the following information should be reported.

765 a) **Data and endpoints.** Mortality data from the treated group should be corrected for
766 negative control mortality with Abbott's Formula or the equivalent.

767 b) **Mortality, morbidity and knockdown.** Report the number and percentage of bed
768 bugs killed, moribund and knockdown exclusively for each treatment replicate at
769 each observation interval. Report mean mortality data for each strain in each
770 treatment at each height level and all heights combined as corrected arithmetic
771 mean values. Confidence limits around the mean values should be reported.

772 c) **Data analysis.** The analysis should consider the effect of the treatment cage height
773 and bed bug strain effects on product performance.

774 **4.5.4. Efficacy evaluation.**

775 Efficacy is usually considered sufficient if – at the end of the test, in each life stage tested
776 – 100% mortality rate, corrected according to Abbott is achieved.

777 **4.5.5. Study conclusions.**

778 Discuss the mortality (total and percent) for each replicate and treatment.

779

780 **4.6. Specific guidance for laboratory studies for direct application testing of**
781 **insecticide products.**

782 Different bioassays are suitable for determination of a direct effect against bed bugs.

783 **4.6.1. Study objective.**

784 To determine the product performance of direct application of insecticide product
785 formulations against bed bugs.

786 **4.6.2. Materials and methods**

787 a) **Experimental unit.** Testing should be conducted with caged bed bugs. Typically,
788 a test cage unit is a 0.4 l squat plastic cup with a screened bottom that has the inside
789 lined with a lubricant to prevent bed bug escape. Other cage designs are acceptable
790 provided the spray does not pool in the cage after spraying. A metered bench top
791 sprayer is preferred as the delivery device to ensure consistent application volume
792 and even distribution of spray particles. Applications should be made at the
793 claimed label rate and should be made from 30 cm above the test cage.

794 b) **Replication.** Tests should be conducted with in minimum five treated and five
795 negative control replicates each with 10 bed bugs.

796 c) **Bed bug exposure to the treatments.** Bed bugs should be transferred to clean
797 containers (e.g. petri dishes) in less than 4 hours or in accordance with the claim
798 after product application. Containers should be stored under ambient test site
799 conditions.

800 d) **Positive control.** A positive control is not necessary.

801 **4.6.3. Reporting results.**

802 See Section 3.2.f. of this guidance for guidance on results reporting and data analysis.

803 **4.6.4. Efficacy evaluation.**

804 Efficacy is usually considered sufficient if – at the end of the test, in each life stage tested,
805 – 100% mortality rate, is achieved.

806 **4.6.5. Study conclusions.**

807 Discuss knockdown, morbidity and mortality (total and percent) of the product treatment.
808

809 **4.7. Specific guidance for laboratory studies for testing ovicidal products.**

810 Different bioassays are suitable for determination of an ovicidal effect against bed bug
811 eggs.

812 **4.7.1. Study objectives.**

813 To determine the product performance of insecticide products intended for use as ovicides.

814 **4.7.2. Materials and methods**815 *Treatments.*

816 I. **Direct application.** Testing should be conducted with bed bug eggs laid on
817 filter paper the night before the test. Therefore, gravid female bed bugs of the
818 same age should be separated from the colony, allowing them to lay eggs in a
819 separate chamber on a filter paper surface. For females, the same feeding
820 conditions/blood source like in the rearing should be used. Tissue culture plates
821 or small petri dishes are recommended as test containers. Pieces of egg-laden
822 filter paper should be cut into pieces that fit into the depressions on the plate
823 or in a petri dish. Twenty eggs should be allotted to each depression or well. A
824 metered bench top sprayer is preferred as the spray device to ensure consistent
825 application volume and even distribution of spray particles. Applications
826 should be made at the claimed label rate and should be made from 30 cm above
827 the test surface (or as directed on the label). For a dust formulation, application
828 should be made at the claimed label rate from a height of 15 cm or less (or as
829 directed on the label). Record the weight of formulation applied and the weight
830 of active ingredient delivered. An equal number of negative control eggs
831 should be included in the study design on the same type of plates.

832 II. **Contact with residual surface application.** Gravid female bed bugs of the
833 same age should be separated from the colony, allowing them to lay eggs in a
834 separate chamber on a filter paper surface. For females the same feeding
835 conditions/blood source like in the rearing should be used and the eggs used
836 should be collected every day to ensure they are the same age. A petri dish is
837 recommended as the test container. The surfaces to be treated with the product
838 should have sorptive and non- sorptive properties and should be representative
839 of where bed bugs are found. For example: unpainted/unfinished plywood;
840 wallpaper, 100% cotton sheeting stretched over a cardboard panel, commercial
841 linoleum tile; laminate or tiles. Surfaces should be treated with the product
842 formulation at the claimed label rate and the residues should be aged for the
843 desired time. Another piece of the respective surface should be left untreated
844 to serve as the negative control. One piece should be placed into each dish.
845 Collect eggs from the filter paper in the gravid female chamber carefully
846 without damaging the eggshell and transfer 20 eggs to each treated and
847 untreated surface in the petri dishes.

848 a) **Replication.** A minimum of ten treated and ten untreated replicates each with 20
849 eggs should be tested.

850 b) **Exposure time.** Plates or dishes should be stored in the laboratory. Eggs should be
851 exposed continuously for 14-30 days or according to the claim. Observations for
852 mortality and hatching should be made every 24 hours for up to 30 days. Eggs
853 should be examined microscopically to determine if egg hatch has taken place, and
854 the number of unhatched and hatched eggs should be recorded from the treated and
855 negative control groups.

856 c) **Positive control.** A positive control is not necessary.

857 **4.7.3. Reporting results.**

858 See Section 3.2.f. of this guidance for guidance on results reporting and data analysis.

859 **4.7.4. Data and endpoints.**

860 Mortality data from the treated group should be corrected for negative control mortality
861 with Abbott's Formula or the equivalent.

862 **Egg mortality.**

863 Report observations every 24 hours for a minimum of 14 but no more than 30 days post-
864 exposure. The percentage of unhatched and hatched eggs for each treatment at each
865 observation interval should be reported.

866 **4.7.5. Efficacy evaluation.**

867 Efficacy is usually considered sufficient if – at the end of the test – 100% mortality rate, is
868 achieved.

869 **4.7.6. Study conclusions.**

870 Discuss ovicidal performance of the product formulation treatment.

871

872 **4.8. Specific guidance for laboratory studies for fumigant products against all bed**
873 **bug life stages.**

874 Different bioassays are suitable for determination of an effect by fumigant products against
875 all bed bug life stages (e.g. 6).

876 **4.8.1. Study objective.**

877 To determine the product performance of a fumigant in the laboratory against all bed bug
878 life stages.

879 **4.8.2. Materials and Methods**

880 a) **Experimental unit (34).** Five, 3.8 liter sealed glass containers with tubing capable
881 of delivering and evacuating fumigant in a closed system should be used as the
882 fumigation chambers. Bed bugs should be placed in a separate ventilated glass vial
883 that should be wrapped in mattress padding and placed in the chamber before
884 fumigation.

885 b) **Product treatment.** Treatment should be at the claimed rate directed by the
886 product label. This rate should be monitored by chemical detection to ensure the
887 target dose was achieved.

888 c) **Bed bug life stage.** This test may be used to evaluate product performance against
889 all bed bug life stages.

890 d) **Replication.** Five treated and five negative control replicates of 10 bed bugs each
891 from the same life stage of the same strain should be tested, with the exception of
892 eggs where 20 eggs should be included per each of the ten treated and ten untreated
893 replicates.

894 e) **Exposure time.** Bed bugs should be exposed for 24 hours or according to the label
895 claim. After fumigation, transfer the bed bugs to clean containers. For experiments
896 with eggs and nymphs, observe for egg hatch and survival of nymphs for 14-30
897 days.

- 898 f) **Environmental conditions.** Replication and treatment should be repeated at 15°C
899 (59° F) and 25°C (77° F).
- 900 g) **Negative control.** Bed bugs in the control group should be placed in a separate
901 ventilated glass vial and should be wrapped in mattress padding but placed in an
902 untreated fumigation chamber for the same period of time as the treatment group.
- 903 h) **Positive control.** A positive control is not necessary.

904 **4.8.3. Reporting results.**

905 Refer to Section 3.2.f. of this guidance for guidance on reporting results and data analysis.
906 In addition, report the following information.

- 907 a) **Amount of product applied.** The amount of product, expressed as weight of
908 product per unit volume, should be reported for each replicate.
- 909 b) **Mortality.** Mortality should be reported as number of dead bed bugs and percent
910 mortality as stated in Section 3.1.e.
- 911 c) **Egg hatch and survival of emerging nymphs.** Report egg hatch (total and
912 percent) and length of time the emerging nymphs survive.

913 **4.8.4. Efficacy evaluation.**

914 Efficacy is usually considered sufficient if – at the end of the test, in each life stage tested,
915 – 100 % mortality rate, is achieved.

916 **4.8.5. Study conclusions.**

917 Report the application rates at which product performance was achieved.
918

919 **4.9. Specific guidance for laboratory studies of insect growth regulators (IGRs).**

920 Different bioassays are suitable for determination of an effect by IGRs against bed bugs.
921 This guidance is based on approaches to evaluate hormonal IGRs and chitin synthesis
922 inhibitors against bed bugs (4, 32).

923 **4.9.1. Study objective.**

924 To determine the product performance of an insect growth regulator in the laboratory
925 against bed bugs.

926 **4.9.2. Materials and Methods**

- 927 a) **Life stages.** Life stages should be evaluated separately and individuals in the
928 population should be the same age. These products may be tested against eggs,
929 nymphs, and adults (potential decreased fecundity and oviposition). If specific
930 stages are claimed, these will have to be tested.
- 931 b) **Replication.** A minimum of five treated replicates and five untreated replicates
932 each with a minimum of 10 specimens should be tested for every life stage for
933 every IGR product tested.
- 934 c) **Test chemical.** Testing should be performed with the whole product formulated
935 with IGRs in combination with other active ingredients. In addition, testing should

- 936 be performed also with the IGR alone to verify its effect. Testing should not be
937 performed with tank mixes (mixtures) containing IGRs.
- 938 d) **Direct spray testing.** Testing should be conducted as described in Section 4.6. of
939 this guidance. Evaluation observations should be reported up to 30 days post-
940 exposure or according to the claim.
- 941 e) **Confinement to treated surfaces.** Testing should be conducted as described in
942 Section 4.2. of this guidance but evaluated up to 30 days.
- 943 f) **Blood meal.** A blood meal is needed for a bed bug to molt from one instar to the
944 next and from the final nymph instar to the adult stage. A source of blood should
945 be available regularly throughout the testing period.

946 **4.9.3. Reporting results.**

947 See Section 3.2.f. of this guidance for guidance on results reporting and data analysis.
948 Additional data reporting is described below.

- 949 a) Report any abnormalities in bed bug development including deformities.
- 950 b) Report egg hatch success and development of hatching nymphs.
- 951 c) Report survivorship of all life stages.
- 952 d) If female bed bugs were tested, track egg production, hatching success of eggs,
953 developmental success and survivorship of nymphs.

954 **4.9.4. Efficacy evaluation.**

955 Efficacy is usually considered sufficient if, in each life stage tested, 100% of the bed bugs
956 do not develop to the next instar. In addition, if female bed bugs were tested neither eggs
957 should be produced nor nymphs should hatch.

958 **4.9.5. Study conclusions.**

959 Discuss results and describe whether or not IGR effects impacted bed bug survivorship and
960 development.

961

5. References

962

963 The following publications were consulted for supporting guidance recommendations.

964

965 1. Aak, A. and B. A. Rukke. Bed bugs, their blood sources and life history
966 parameters: a comparison of artificial and natural feeding. *Medical and*
967 *Veterinary Entomology*. 2014. 28: 50-9. doi: 10.1111/mve.12015.

968 2. Aitchison, J. The Statistical Analysis of Compositional Data. *Chapman and*
969 *Hall. London*. 1986. 416.

970 3. Bailer, A. J. and W. W. Piegorsch. Chapter 7 Dose-response modeling and
971 analysis. *Dose Response Modeling in Statistics for Environmental Biology*
972 *and Toxicology*. Chapman & Hall. London. 1997.

973 4. Bajomi, D., J. Szilagyi, R. Naylor and L. Takacs. S-Methoprene formulations
974 laboratory tests for efficacy against bed bugs. *Proceedings of the Seventh*
975 *Annual ICUP*. 2011. 249-256.

976 5. Barile, J., R. Nauen, G. Nentwig, R. Pospischil and B. Reid. Laboratory and
977 field evaluation of deltamethrin and bendiocarb to control *Cimex lectularius*
978 (Heteroptera: Cimicidae). In: *W. H. Robinson and D. Bajomi (eds).*
979 *Proceedings of the Sixth International Conference on Urban Pests. OOK-*
980 *Press Kft., H-8200 Veszprém, Pápai út 37/a. Hungary*. 2008. 105-111.

981 6. Bonds, J. A. S, M. Greer, J. Coughlin and V. Patel. Caged mosquito bioassay:
982 Studies on cage exposure pathways, effects of mesh on pesticide filtration,
983 and mosquito containment. *American Mosquito Control Association*. 2010.
984 26(1):50-56.

985 7. Busvine, J. R. Insecticide resistance in bed-bugs. *Bulletin of the World Health*
986 *Organization*. 1958. 19: 1041-1052.

987 8. Busvine, J. R. and J. Lien. Methods for measuring insecticide susceptibility
988 levels in bed-bugs, cone-nosed bugs fleas and lice. *Bulletin of the World*
989 *Health Organization*. 1961. 24: 502-517.

990 9. Cochran, W. G. and G. M. Cox. *Experimental Designs (Second edition)*. John
991 Wiley & Sons, Inc., NY. 1992.

992 10. Dawson, G., Griffiths, D., Janes, N., Mudd, A., Pickett, J., Wadhams and L.
993 & C. Woodcock, C. Identification of an aphid sex pheromone. *Nature*. 1987.
994 325: 614– 616.

995 11. Domingue, M. J., M. Kramer and M. F. Feldlaufer. Sexual dimorphism of
996 arrestment and gregariousness in the bed bug (*Cimex lectularius*) in response
997 to cuticular extracts from nymphal exuviae. *Physiological Entomology*. 2010.
998 35(3):203-213.

999 12. European Chemicals Agency. *Transitional Guidance on the Biocidal Products*
1000 *Regulation. Transitional Guidance on Efficacy Assessment for Product Type*
1001 *18, Insecticide, Acaricides & other Biocidal Products against Arthropods and*
1002 *Product Type 19, Repellents & Attractants*. 2016. 1-111.

1003 13. Faraway, J. J. *Extending the Linear Model with R: Generalized Linear, Mixed*

- 1004 Effects and Nonparametric Regression Models. *Chapman and Hall/CRC*.
1005 *Boca Raton, FL*. 2006.
- 1006 14. Gbur, E. E., W. W. Stroup, K. S. McCarter, S. Durham, L. J. Young, M.
1007 Christman, M. West and M. Kramer. Analysis of Generalized Linear Mixed
1008 models in the Agricultural and Natural Resources Sciences. *American Society*
1009 *of Agronomy*. Madison, WI. 2012.
- 1010 15. Geier, M. and J. Boeckh. A new Y-tube olfactometer for mosquitoes to
1011 measure the attractiveness of host odors. *Entomologia Experimentalis Et*
1012 *Applicata*. 1999. 92: 9-19.
- 1013 16. González-Morales, M. A., M. Terán and A. Romero. Behavioral responses of
1014 the common Bed Bug to essential oil constituents. *Insects*. 2021.12, 184.
- 1015 17. Gratz, N. A survey of bed-bug insecticide resistance in Israel. *Bulletin of the*
1016 *World Health Organization*. 1959. 20: 835-840.
- 1017 18. Habedank B., Snelinski B., Reinsch M. and J. Klasen. Advanced method for
1018 evaluation of the residual efficacy of products to control *Cimex lectularius*
1019 (Hemiptera: Cimicidae). Robinson W.H. and de Carvalho Campos A.E.
1020 (eds.): *Proceedings of the Seventh International Conference on Urban Pests,*
1021 *Instituto Biologico, Sao Paulo, SP Brasil*. 2011. 271-276.
- 1022 19. Halliday, W.R. and K. P. Burnham. Choosing the optimal diagnostic dose for
1023 monitoring insecticide resistance. *Journal of Economic Entomology*. 1990.
1024 83(4): 1151-1159
- 1025 20. Harlan, H. J., M. K. Faulde, and G. J. Baumann. Bedbugs. In: X. Bonnefoy,
1026 H. Kampen, and K. Sweeney (eds.) *Public Health Significance of Urban Pests.*
1027 *World Health Organization*. Copenhagen, Denmark. 2008. 131-151.
- 1028 21. Health and Safety Executive (HSE), Biocides and Pesticides Unit (BPU).
1029 Guidelines on the efficacy data for approval of non-agricultural pesticide
1030 products: insecticides and acaricides. Merseyside, UK. 2004.
- 1031 22. Insecticide Resistance Action Committee (IRAC). 2016. Resistance
1032 Definition. Available on-line at: <http://www.iraconline.org/about/resistance/>
- 1033 23. Krüger A., E. Schmolz, and A. Vander Pan. Methods for Testing Repellents
1034 Against Bed Bugs (Hemiptera: Cimicidae). *Journal of Economic Entomology*.
1035 2021. 114(1), 265–273
- 1036 24. Levinson, H. Z. and A. R. Bar Ilan. Assembling and alerting scents produced
1037 by the bedbug *Cimex lectularius*. *Experientia*. 1970. 27:102-103.
- 1038 25. McCullagh, P. and J. A. Nelder. *Generalized Linear Models* (Second edition).
1039 *Chapman and Hall*, NY. 1989.
- 1040 26. Miller, R. G., Jr. *Survival Analysis*. *John Wiley & Sons, Inc*. NY. 1981.
- 1041 27. Miller, D. M. and M. L. Fisher. Bed bug (Hemiptera: Cimicidae) response to
1042 fumigation using sulfuryl fluoride. IW. H. Robinson and D. Bajomi (eds).
1043 *Proceedings of the Sixth International Conference on Urban Pests. OOK-*
1044 *Press Kft., H-8200 Veszprém, Pápai út 37/a, Hungary*. 2008. 123-127.
- 1045 28. Montes, C., Cuadrillero, C. and D. Vilella. Maintenance of a Laboratory
1046 Colony of *Cimex lectularius* (Hemiptera: Cimicidae) Using an Artificial
1047 Feeding Technique. *Journal of Medical Entomology*. 2002. 39: 675–679. doi:
1048 10.1603/0022-2585-39.4.675.

- 1049 29. Moore, D. J. and D. M. Miller. Laboratory evaluations of insecticide product
1050 efficacy for control of *Cimex lectularius*. *Journal of Economic Entomology*.
1051 2006. 99(6): 2080-2086.
- 1052 30. Morgan, B. J. T. (editor). *Statistics in Toxicology*. Oxford University Press.
1053 NY. 1996.
- 1054 31. Myamba, J., C. A. Maxwell, A. Sidi and C. F. Curtis. Pyrethroid resistance in
1055 tropical bedbugs, *Cimex hemipterus*, associated with use of treated bed nets.
1056 *Medical and Veterinary Entomology*. 2002. 16:448-451.
- 1057 32. Naylor, R. D. Bajomi and C. Boase. Efficacy of (s)-Methoprene against *Cimex*
1058 *lectularius* (Hemiptera: Cimicidae) in: *W. H. Robinson and D. Bajomi (eds).*
1059 *Proceedings of the Sixth International Conference on Urban Pests*. OOK-
1060 Press Kft., H-8200 Veszprém, Pápai út 37/a, Hungary. 2008.
- 1061 33. Pennetier, C., V. Corbel, P. Boko, A. Odjo, R. N'Guessan, B. Lapied and J-M
1062 Hougard. Synergy between repellents and non-pyrethroid insecticides
1063 strongly extends the efficacy of treated nets against *Anopheles gambiae*.
1064 *Malaria Journal* 6:38. 2007. doi:10.1186/1475-2875-6-38.
- 1065 34. Phillips, T. W., M. J. Atkins, E. Thomas, J. Demarck and C. Wang.
1066 Fumigation of bed bugs (Hemiptera: Cimicidae): effective application rates
1067 for sulfuryl fluoride. *Journal of Economic Entomology*. 2014. 107(4): 1582-
1068 1589.
- 1069 35. Reinhardt, K. and M. T. Siva-Jothy. The biology of bed bugs (Cimicidae).
1070 *Annual Review of Entomology*. 2007. 52, 351-374.
- 1071 36. Robertson, J. L., R. M. Russell, H. K. Preisler and N. E. Savin (eds). *Bioassays*
1072 *with Arthropods*. Second edition. CRC Press. Boca Raton, FL. 2007.
- 1073 37. Romero, A., M. F. Potter, D.A. Potter and K. F. Haynes. Insecticide resistance
1074 in the bed bug: A factor in the pest's sudden resurgence? *Journal of Medical*
1075 *Entomology*. 2007. 44(2): 175-178.
- 1076 38. Romero, A., M. F. Potter and K. F. Haynes. Behavioral response of the bed
1077 bug to insecticide residues. *Journal of Medical Entomology*. 2009. 46(1): 51-
1078 57.
- 1079 39. Romero, A. and T. D. Anderson. High levels of resistance in the common Bed
1080 Bug, *Cimex lectularius* (Hemiptera:Cimicidae), to Neonicotinoid
1081 Insecticides. *Journal of Medical Entomology*. 2016. 53(3): 727-731.
- 1082 40. Schabenberger, O. and F. J. Pierce. *Contemporary Statistical Models for the*
1083 *Plant and Soil Sciences*. CRC Press. NY, NY. 2002.
- 1084 41. Singh, N., Wang C. and R. Cooper. Effect of trap design, chemical lure,
1085 carbon dioxide release rate, and source of carbon dioxide on efficacy of bed
1086 bug monitors. *Journal of Economic Entomology*. 2013. 105 (4): 1802-1811.
- 1087 42. Siva-Jothy, M. T. Trauma, disease and collateral damage: conflict in cimicids.
1088 *Philosophical Transactions of the Royal Society of London Biological*
1089 *Sciences*. 2006. 361:269-275.
- 1090 43. Steelman, C. D., A. L. Szalanski, R. Trout, J. A. McKern, C. Solorzano and J.
1091 W. Austin. Susceptibility of the bed bug *Cimex lectularius* L. (Heteroptera:
1092 Cimicidae) collected in poultry production facilities to selected insecticides.
1093 *Journal of Agricultural and Urban Entomology*. 2008. 25(1): 41-51.

- 1094 44. Stuart, A., and Ord and K. Kendall's Advanced Theory of Statistics Sixth
1095 Edition Volume 1. Distribution Theory. Wiley. London. 1998. 351.
- 1096 45. Throne, J. E., D. K. Weaver and J. E. Baker. Probit analysis: Assessing
1097 goodness-of- fit based on back-transformation and residuals. *Journal of*
1098 *Economic Entomology*. 1995. 88(5), 1513-1516.
- 1099 46. Vander Pan, A., E. Schmolz, J. Krücken and C. Kuhn. A Novel Simulated-
1100 Use Test for Determining the Efficacy of Insecticides Against Bed Bugs
1101 (Hemiptera: Cimicidae). *Journal of Economic Entomology*. 2019. 112(5),
1102 2345–2353
- 1103 47. Wang, C., Lü, L., Zhang, A. and C. Liu. Repellency of Selected Chemicals
1104 Against the Bed Bug (Hemiptera: Cimicidae). *Journal of Economic*
1105 *Entomology*. 2013. 2522-2529.
- 1106 48. Weeks, E. N. I., J. G. Logan, M. A. Birkett, J. A. Pickett and M. M. Cameron.
1107 Tracking bed bugs (*Cimex lectularius*): a study of the effect of physiological
1108 and extrinsic factors on the response to bed bug-derived volatiles. *The Journal*
1109 *of Experimental Biology*. 2013. 216: 460-469. doi:10.1242/jeb.074930
- 1110 49. Weeks, E. N. I., J. G. Logan, S. A. Gezan, C. M. Woodcock, M. A. Birkett,
1111 Pickett J. A. and M. M. Cameron. A bioassay for studying behavioural
1112 responses of the common bed bug, *Cimex lectularius* (Hemiptera: Cimicidae)
1113 to bed bug-derived volatiles. *Bulletin of Entomological Research*. 2011. 101:
1114 1–8. doi: 10.1017/S0007485309990599.
- 1115 50. White, G. B. Chapter 2: Terminology of Insect Repellents. pp. In: *M.*
1116 *Debboun, S. P. Frances, and D. Strickman (eds). Insect Repellents:*
1117 *Principles, Methods, and Uses*. CRC Press, Boca Raton, FL. 2007. 31-47.
- 1118 51. World Health Organization Pesticide Evaluation Scheme (WHOPES). Test
1119 procedures for insecticide resistance monitoring in malaria vectors, bio-
1120 efficacy and persistence of insecticide on treated surfaces. *World Health*
1121 *Organization*. Geneva, Switzerland. 1998.
- 1122 52. WHOPES. 2009. Guidelines for efficacy testing of household insecticide
1123 products – Mosquito coils, vaporizer mats, liquid vaporizers, ambient
1124 emanators and aerosols. *World Health Organization*. Geneva, Switzerland.
- 1125 53. WHO Vector Control and Biology Division. 1982. VI. Bed bugs.
1126 WHO/VBC/82.857, 9 pp. *World Health Organization*, Geneva, Switzerland.
1127